


Plant Population Ecology

- Populations
- Measuring Plant density
- Population growth models




Population

- Population = group of individuals of a species with the potential to interbreed in a defined geographic region



Population


- What is an individual?
- Branches (modular units)
- genets = sexual reproduction, genetically distinct individuals
- ramets = asexual reproduction, tillers, runners; can be physiologically distinct



Opuntia fulgida

- Clonal fragmentation
- ramets





Measuring Plant density

- #'s / unit area, volume, diameter at breast height

To experience quadrat sampling in plant populations, my ecology classes set out a transect in an upland hardwood forest in southern Indiana. We laid out three lines, each 110 meters long, and counted all trees taller than 25 cm within a swath of 1 meter on either side of the line. Each transect line is, in effect, a very long, thin quadrat with an area of 220 m² (0.022 hectare). We obtained the following results:


	<i>No. counted</i>			<i>Estimated no. per hectare</i>		
	Line A	Line B	Line C	Line A	Line B	Line C
Chestnut oak	20	28	18	909	1273	818
Sugar maple	5	4	7	227	182	318
American beech	13	15	16	591	682	727

Plant structure problems

- seed pool
- age structure
- age of reproduction
- population density usually not static



Trillium grandiflorum



Population structure






Figure 4-3 An idealized plant life history. Near the middle of the growing period, N is 5 or 6, depending on whether the vegetative daughter is considered an individual or part of the parent plant. (Copyright © 1977 by J. L. Harper, *Population Biology of Plants*.)



Age vs stage in plants


(A)



Population growth models

- **Exponential growth model**
- birth and death rates
- immigration/emigration
- $dN/dt = rN$
- assume I and E are 0
- examples of exponential growth

Fig. 1.2 (a) Geometric; and (b) logistic population growth.



Exponential growth

$$\frac{dN}{dt} = rN$$

$$N_t = N_0 e^{rt}$$

$$R_0 = N_{t+1}/N_t$$




Fig. 5.2 The exponential increase of two species. (a) *Pinus sylvestris* between 9500 and 9000 years ago, when this tree was invading Hockham Mere, Norfolk, England. The abundance of *P. sylvestris* is plotted as the density of pollen grains in peat samples, which it is assumed is correlated with the historical size of the tree population (from Bennett 1983). (b) *Avena fatua* infesting a barley crop at Boxworth Experimental Farm, Cambridge, England (from Selman 1970).

Logistic growth model

- actual populations plateau
- carrying capacity
- $dN/dt = rN (K-N/K)$
- different types of logistic growth curves
- growth rates differ in different phases
- Overshoot K
- K dependent on resource availability




Fig. 1/2. Population growth curve for the alga *Chlorella* in culture. This alga forms clumps and so experiences the effects of density when there are still available resources in the medium. The initially exponential growth rate declines after 8 days but if the clumping is prevented by shaking the exponential rate continues for four more days. (From Pearsall and Bengry, 1940) x and + unshaken, o shaken

Growth rates

Population growth rates of some aquatic plants in culture

Species	Relative growth rate (g/g/day)		Simple growth rate (g/beaker/day)		12 weeks "stable" population (g/beaker)	
	Phase I	(Rank)	Phase II	(Rank)	Phase III	(Rank)
<i>L. minor</i>	0.35	(1)	14.7	(2)	603	(2)
<i>L. gibba</i>	0.25	(3)	9.0	(4)	355	(4)
<i>L. polyrrhiza</i>	0.21	(4)	12.5	(3)	776	(1)
<i>S. natans</i>	0.28	(2)	18.5	(1)	524	(3)

Logistic growth






Figure 18.6 (a) A stylized logistic growth curve and (b–e) types of fluctuations that take place when the population overshoots K . (b) Chaotic fluctuation. The population fluctuates wildly with no regulation. Such fluctuations can lead to sudden extinction. (c) Stable-limit cycles. The population fluctuates about some equilibrium level, with fluctuations having a certain period and amplitude. (d) Oscillations decrease over time. After overshooting K , the population levels off and maintains itself at K through compensating birth and death rates. (e) The population strongly overshoots K and then crashes. It may drop very low, recover, and return to some lower equilibrium level, or it may go extinct.

Fig. 1.4 Growth of a moss population colonizing the Icelandic island of Surtsey. (From Fridrikson 1975)



Random variation in Population growth and decline

- Hurricanes
- Fire
- Pests



Matrix models

- Transition Matrix
- calculation of plant populations
- matrix multiplication

		This census		
		Seed	Rosette	Flower
Next census	Seed	a_{ss}	a_{rs}	a_{fs}
	Rosette	a_{sr}	a_{rr}	a_{fr}
	Flower	a_{sf}	a_{rf}	a_{ff}

Multiplication

Column matrix

$$\begin{bmatrix} N_s \\ N_r \\ N_f \end{bmatrix}$$

$$\begin{matrix} \mathbf{A} & \times & \mathbf{B}_1 & = & \mathbf{B}_2 & \text{(Matrix 4-3)} \\ \begin{bmatrix} a_{ss} & 0 & a_{fs} \\ a_{si} & a_{ii} & a_{fi} \\ 0 & a_{if} & a_{ff} \end{bmatrix} & \times & \begin{bmatrix} N_s \\ N_i \\ N_f \end{bmatrix} & = & \begin{bmatrix} (N_s a_{ss}) + 0 + (N_f a_{fs}) \\ (N_s a_{si}) + (N_i a_{ii}) + (N_f a_{fi}) \\ 0 + (N_i a_{if}) + (N_f a_{ff}) \end{bmatrix} \end{matrix}$$

- Row X column
- Sum of result
- Gives new column matrix

Site A (northeastern exposure)	Site B (southwestern exposure)	Site C (hilltop) A_{\max}
$A_{\text{site A}} = \begin{bmatrix} 0.672 & 0 & 0.561 \\ 0.018 & 0.849 & 0 \\ 0 & 0.138 & 0.969 \end{bmatrix}$	$A_{\text{site B}} = \begin{bmatrix} 0.493 & 0 & 0.561 \\ 0.013 & 0.731 & 0 \\ 0 & 0.234 & 0.985 \end{bmatrix}$	$A_{\text{site C}} = \begin{bmatrix} 0.434 & 0 & 0.560 \\ 0.333 & 0.610 & 0 \\ 0 & 0.304 & 0.956 \end{bmatrix}$

Stable age structure

