

1 LIST OF MAPS	- 1 -
2 EXECUTIVE SUMMARY	- 2 -
2 PURPOSE AND GOALS	- 3 -
2.1 PURPOSE	- 3 -
2.2 PROJECT GOALS	- 3 -
2.3 STRATEGY	- 4 -
2.4 ASSESSMENT GUIDELINES	- 4 -
3 HOW TO USE THIS REPORT	- 6 -
3.1 LIST OF ACRONYMS AND ABBREVIATIONS	- 6 -
3.2 DATA	- 7 -
3.2.1 USE RESTRICTIONS	- 7 -
3.2.2 USE OF EXISTING DATA	- 7 -
3.2.2.1 Scale of Data and Units of Comparison	- 7 -
3.2.3 ACCURACY & UNCERTAINTY	- 8 -
4 WATERSHED CHARACTERISTICS AND RESOURCES	- 8 -
4.1 LOCATION	- 8 -
4.2 SETTLEMENT	- 9 -
4.2.1 HISTORIC SETTLEMENT	- 9 -
4.2.2 CITIES	- 9 -
4.2.3 HUMAN POPULATION	- 11 -
4.3 TRANSPORTATION	- 12 -
4.3.1 ROADS	- 12 -
4.3.2 RAILROADS	- 13 -
4.3.3 AIRPORTS	- 13 -
4.3.4 PORTS	- 13 -
4.4 LAND COVER, VEGETATION AND LAND USE	- 14 -
4.4.1 LAND COVER	- 14 -
4.4.1.1 Historic Land Cover Conditions	- 15 -
Historic Wildfire Conditions	- 18 -
4.4.1.2 Current Land Cover Conditions	- 20 -
4.4.1.3 Current Wildfire Conditions	- 25 -
4.4.2 VEGETATION AND VEGETATION COMMUNITIES	- 26 -
4.4.2.1 Native Vegetative Diversity and Weeds	- 26 -
Data Availability	- 26 -
Species Status and Geographic Patterns	- 27 -
Weed Species	- 28 -

Important Habitats and Communities	- 29 -
4.4.2.2 Wetlands	- 34 -
4.4.2.3 Riparian Areas	- 39 -
4.4.3 LAND USE	- 40 -
4.4.4 HISTORIC LAND USE CONDITIONS	- 41 -
4.4.5 CURRENT LAND USE CONDITIONS	- 41 -
4.4.5.1 Agriculture	- 41 -
4.4.5.2 Mills	- 42 -
4.4.5.3 Mining	- 44 -
4.4.5.4 Landfills	- 45 -
4.4.5.5 Military Facilities	- 45 -
4.4.5.6 Wildlife Preserves	- 46 -
4.4.6 LAND OWNERSHIP	- 46 -
4.5 ZONING AND REGULATIONS	- 47 -
4.6 ECOREGIONS AND PHYSIOGRAPHIC PROVINCES	- 48 -
4.7 CLIMATE	- 50 -
4.7.1 TEMPERATURE	- 50 -
4.7.2 PRECIPITATION	- 50 -
4.8 SOILS	- 51 -
4.8.1 HYDRIC SOILS	- 51 -
4.8.2 ERODIBLE SOILS	- 56 -
4.9 LITHOLOGY	- 56 -
4.10 ELEVATION	- 59 -
4.11 AQUATIC RESOURCES	- 59 -
4.11.1 HISTORIC AQUATIC RESOURCE CONDITIONS	- 59 -
4.11.2 CURRENT AQUATIC RESOURCE CONDITIONS	- 61 -
4.11.2.1 Streams	- 61 -
4.11.2.2 Lakes and Reservoirs	- 62 -
4.11.2.3 Springs	- 62 -
4.12 FISH AND WILDLIFE	- 63 -
4.12.1 FISH	- 63 -
4.12.1.1 General Fish Habitat Summary	- 66 -
4.12.1.2 Historic Catch Records (steelhead, cutthroat)	- 69 -
4.12.1.3 Historic/Recent Juvenile and Spawner Surveys	- 70 -
4.12.1.4 Hatcheries	- 71 -
4.12.1.5 Endangered/Threatened Fish Species and Species of Concern	- 72 -
4.12.2 WILDLIFE	- 73 -
4.12.2.1 Insects	- 73 -
Data Availability	- 73 -
Species Status and Geographic Patterns	- 73 -
Important Habitats and Communities	- 74 -
Conservation and Restoration Potential	- 74 -
4.12.2.2 Amphibians and Reptiles	- 74 -
Data Availability	- 74 -

Species Status and Geographic Patterns	- 75 -
Pest Species	- 76 -
Important Habitats and Communities	- 76 -
Conservation and Restoration Potential	- 76 -
4.12.2.3 Birds	- 77 -
Data Availability	- 77 -
Species Status, Trends, and Geographic Patterns	- 78 -
Pest Species	- 81 -
Game Birds	- 82 -
Important Habitats and Communities	- 82 -
Conservation and Restoration Potential	- 82 -
4.12.2.4 Mammals	- 83 -
Data Availability	- 83 -
Species Status and Geographic Patterns	- 83 -
Pest Species	- 84 -
Important Habitats and Communities	- 85 -
Conservation and Restoration Potential	- 85 -
4.13 EXISTING WATERSHED ANALYSES	- 87 -
4.14 EXISTING PRESERVES AND WATERSHED RESTORATION PROJECTS	- 87 -
5 LAND COVER ANALYSIS	- 92 -
5.1 TERRESTRIAL ANALYSES	- 92 -
5.1.1 FORESTED AND NON-FORESTED AREAS	- 92 -
5.1.2 OAK SAVANNAH	- 93 -
5.2 WETLAND ANALYSES	- 93 -
5.3 RIPARIAN ANALYSES	- 93 -
5.3.1 STREAM CHANNEL PERCENT SHADE	- 99 -
5.4 INTERPRETATION AND RECOMMENDATIONS	- 100 -
5.4.1 RECOMMENDED ANALYSES	- 101 -
5.4.1.1 Locate Riparian Wetlands	- 101 -
5.4.1.2 Locate Riparian Planting Areas	- 101 -
6 AQUATIC ANALYSIS	- 102 -
6.1 STREAM CHANNEL MORPHOLOGY	- 102 -
6.1.1 DATA AVAILABLE AND DATA USED	- 102 -
6.1.1.1 DEM data	- 102 -
6.1.1.2 AHI data	- 102 -
6.1.1.3 GIS data	- 102 -
6.1.2 STREAM CONFINEMENT	- 102 -
6.1.3 STREAM GRADIENT	- 103 -
6.1.4 CHANNEL TYPES	- 103 -
6.1.5 SIDE CHANNEL/SECONDARY CHANNEL HABITAT	- 105 -

6.1.6 RIFFLES	- 105 -
6.1.7 PERCENT POOLS	- 105 -
6.1.8 CHANNEL WIDTHS PER POOL	- 106 -
6.1.9 CHANNEL WIDTH-TO-DEPTH RATIO	- 106 -
6.1.10 CHANNEL MODIFICATIONS	- 106 -
6.2 IN-STREAM STRUCTURE	- 106 -
6.2.1 POOL COMPLEXITY	- 106 -
6.2.2 LARGE WOODY DEBRIS	- 106 -
6.2.2.1 LWD Frequency	- 106 -
6.2.2.2 LWD Source Areas	- 106 -
6.3 SUBSTRATES	- 107 -
6.4 STREAM FLOW	- 107 -
6.4.1 GAGE DATA	- 107 -
6.4.2 PEAK FLOWS	- 110 -
6.4.2.1 Land Use	- 110 -
6.4.2.2 Roads	- 110 -
6.4.2.3 Rain-on-Snow	- 112 -
6.4.3 FLOODING	- 113 -
6.4.4 STREAM DYNAMICS	- 113 -
6.5 WATER USE	- 114 -
6.5.1 GROUND WATER	- 116 -
6.5.2 SURFACE WATER	- 119 -
6.5.2.1 Points of Diversion & Types of Water Use	- 119 -
6.5.2.2 Water Availability	125
6.5.3 INTERPRETATION AND RECOMMENDATIONS	127
6.6 WATER QUALITY	127
6.6.1 FACTORS AFFECTING WATER QUALITY	127
6.6.1.1 Point Source Pollution	128
6.6.1.2 Non-Point Source Pollution	129
6.6.1.3 Water Temperature	129
6.6.2 WATER QUALITY EVALUATION	131
303(d) Listed Streams	131
6.6.2.1 High Quality Waters	132
6.6.3 WATER QUALITY MONITORING	132
6.6.3.1 EPA's STORET and ODEQ's LASER	132
6.7 FISH BARRIERS	134
6.7.1 CULVERTS	135
6.7.2 DAMS	135
6.7.3 NATURAL BARRIERS: RAPIDS, FALLS	138
6.8 CHANNEL MODIFICATION ASSESSMENT	138
6.9 INTERPRETATION AND RECOMMENDATIONS	139
7 SALMONID HABITAT ANALYSIS	140

7.1.1 DISTRIBUTIONS	140
7.1.1.1 Dataset Discussion	140
ODFW Distribution Maps and Salmonid Core Areas	140
ODF Fish Limits Maps	141
7.1.2 MULTI-FACTOR ANALYSES OF SALMONID HABITAT	141
7.1.3 MULTI-FACTOR SALMONID HABITAT ANALYSES: SYNTHESIS	152
7.2 INTERPRETATION AND RECOMMENDATIONS	152
8 SURFACE EROSION	153
8.1.1 SOILS	153
8.1.2 EROSION BY LAND USE	153
8.1.3 ROADS	154
8.2 STREAMBANK EROSION	156
8.3 LANDSLIDES	158
8.3.1 ODF DEBRIS FLOW HAZARD MAPS	158
8.4 INTERPRETATION AND RECOMMENDATIONS	160
8.5 DATA COLLECTION AND MONITORING RECOMMENDATIONS	161
8.6 RECOMMENDATIONS FOR FUTURE ANALYSES	161
8.7 WATERSHED ENHANCEMENT RECOMMENDATIONS	161
8.7.1 ECOSYSTEM CONTEXT	161

1 LIST OF MAPS

Map 1. Map of the Luckiamute / Ash Creek Study Area

Map 2. 7th Field Hydrologic Unit Codes (HUC)

Map 3. Ecoregions

Map 4. Current Land Cover

Map 5. Presettlement Land Cover

Map 6. Digital Orthoquadrangle.

Map 7. Hydric Soils

Map 8. Aquatic Habitat Inventory Data (AHI)

Map 9. Distribution of Steelhead in the Luckiamute / Ash Creek Study Area

Map 10. Distribution of Coho in the Luckiamute / Ash Creek Study Area

Map 11. Distribution of Spring Chinook in the Luckiamute / Ash Creek Study Area

Map 12. 303(d) Listed Streams

2 EXECUTIVE SUMMARY

2 PURPOSE AND GOALS

2.1 PURPOSE

A primary purpose of this watershed assessment is to summarize current conditions in the Luckiamute/Ash Creek watersheds and to rank sub-basin so that actions can be planned “for protecting and improving fish habitat and water quality” (Watershed Professionals Network, 1999). Stream networks, together with the landscape they drain, are called watersheds. Watersheds come in all shapes and sizes. Regardless of size, all watersheds are defined by drainages. That is, precipitation falling anywhere within a watershed must eventually flow into a stream, river, lake, or aquifer. Water that has become part of a stream or river leaves the watershed at a specific point. Watersheds of different sizes are nested within one another. Large watersheds, such as the one that drains the Columbia River Basin, are made up of smaller watersheds, such as the Willamette River Basin. Ridge tops delineate watersheds.

In order to facilitate communication among interested groups, larger watersheds have been identified and delineated by governmental agencies such as the U.S. Geologic Survey. These watersheds are designated by unique identifier numbers called Hydrologic Unit Codes (HUC). Frequently watersheds are referred to as 5th, 6th or 7th field watersheds. The terms “fifth”, “sixth” and “seventh” field refer to the size of watersheds, with fifth field being the larger of the three. Fifth field watersheds, the size of watershed for which the OWEB manual was developed (Watershed Professionals Network, 1999), range in size from 40,000 to 120,000 acres and average about 60,000 acres (24,300 ha). The Luckiamute and Rickreall watersheds are examples of 5th field watersheds and each contain many nested 6th and 7th field watersheds. Watersheds are convenient ecological units because they represent bounded areas that often share similar properties like flora and fauna, climatic patterns, and disturbance regimes (see Appendix A: Ecological Processes).

In this report the study area was defined by the Luckiamute Watershed Council as the Luckiamute River Basin and part of the Rickreall Basin. We acknowledge that the Ash Creek (Rickreall Basin) is not a complete watershed.

2.2 PROJECT GOALS

According to the Oregon Watershed Enhancement Board (OWEB), Oregon Watershed Assessment Manual, a Watershed Assessment is a process for evaluating how well a watershed is working (Watershed Professionals Network, 1999). “An assessment can’t give us site-specific prescriptions for fixing problems, but it can, and should, tell us what we need to know to develop action plans and monitoring strategies for protecting and improving fish habitat and water quality” (Watershed Professionals Network, 1999). The outcome of this project include a summary of the condition of the Luckiamute (201,737.94 acre) and Ash Creek (33,887.36 acre) watersheds and results from which an action plan will be developed. This assessment was not intended to identify specific places in the watershed for specific actions, rather it was intended to: (1) summarize the current conditions of the watershed, both descriptively and quantitatively; to identify important data gaps; and (3) to suggest ways in which the watershed council can proceed in its action planning.

Luckiamute Watershed Council (LWC) established the following goals and guidelines for this Watershed Assessment. This assessment will address all major topics outlined I-X in the Oregon Watershed Assessment Manual for the Luckiamute and Ash Creek watersheds. In addition, the assessment will address major ecosystem elements and ecological processes in the watershed uplands and a set of key questions developed by the watershed council members. The LWC identified the following goals for this assessment to (1) produce a source of educational information about watershed resources, environmental and biological processes, and human land uses that affect watershed resources and processes; (2) produce results that will provide restoration recommendations and support development of an action plan that will prioritize future outreach, restoration and monitoring; and (3) identify significant gaps in the understanding of watershed conditions in order to guide future information gathering efforts.

2.3 STRATEGY

Our approach was to produce both descriptive and spatial summaries, which characterized the Luckiamute and Ash Creek watersheds using existing data. Wherever possible, we have used quantitative data that were collected using known protocols. For the GIS-based analyses (spatial summary), we used 1:24,000 uniform scale data that covered the entire study area whenever possible.

Our strategy entailed working closely with members of the LWC by meeting 1-2 times each month; and frequent communication with our LWC liaison, Chris Vandenberg. Rather than performing our watershed assessment and then delivering a final report to the group, we involved LWC council members in the synthesis of watershed data.

2.4 ASSESSMENT GUIDELINES

This assessment was organized around the procedures outlined in the OWEB watershed assessment manual and around a set of key questions developed by the LWC. The assessment team reviewed and acquired available data necessary to follow the OWEB protocols and to address the watershed council's key questions. The assessment team did not acquire data if it was not germane to questions asked by the Council or the OWEB manual. We included web links to key data and information sources so that the Council could update data sets or acquire new information, as necessary, during their action planning process. When data gaps were encountered, we made suggestions for filling those gaps.

Not all data are good data for answering all questions. Wherever possible, we presented the strengths and weaknesses of each data set as the data related to the questions being asked.

This report includes methodological steps that were used to summarize existing conditions in the watershed. This will assist the Watershed Council in repeating these analyses should better data sets become available in the future. We recommend that all summaries and results be critically reviewed before being used in action planning. Our presentation of data limitations and methods

should make the report summaries as ‘transparent’ as possible so that watershed council members can use this watershed assessment wisely.

We urge the Council member to keep in mind that the results of this assessment are based on observations and measurements made by many people using many methods at many times. In most cases, it is very difficult to know how representative the observations and data are of actual conditions in the watershed. In addition, it is important to remember that GIS maps are only computer representations of actual watershed conditions. By definition, GIS maps are much more simple than the phenomena that they represent – they are imperfect. Be a critical data user.

3 HOW TO USE THIS REPORT

This report is organized into two major sections, each with multiple related sub-sections. Section 4 presents descriptions of important watershed characteristics. Sections 5-8 present summaries or analyses of these watershed characteristics. The LWC requested that wherever possible, results be summarize for the Luckiamute and Ash Creek watersheds separately. To minimize redundancy we have referenced material from related sections to each other throughout the report; however, LWC asked that our recommendations be repeated both after each main section and again at the end of the report.

Wherever possible, each section contains a description of the resources; the ecological importance of those resources; a description of the data and methods used in the assessment; the general results; and, where appropriate, our team's recommendations. Since material on any given subject may appear in several places throughout the text, we recommend that the entire document be read.

We have supplied LWC with Identical versions "hard" (printed) and electronic copies of this report. The electronic copy consists of a PDF. PDF files are Adobe Acrobat Portable Document Format files, which can be easily viewed and printed from any computer using the free Adobe Acrobat Reader, available free on the World Wide Web at <http://www.adobe.com/products/acrobat/readstep.html>.

Throughout this report we reference data sources and other studies. Web-based information sources are referenced in the text. Reports and other published sources of information are listed in the Literature Cited section.

The LWC requested that English units be used instead of metric units.

3.1 LIST OF ACRONYMS AND ABBREVIATIONS

AHI	Aquatic Habitat Inventory
BLM	Bureau of Land Management
CD-ROM	Compact Disc, Read Only Memory
CLAMS	Coastal Landscape Analysis and Modeling Study
DEM	Digital elevation model (GIS representation of topography)
DLCD	Department of Land Conservation and Development
DLG	Digital Line Graph
DEQ	Department of Environmental Quality
DOQ	Digital Ortho Quad
EPA	U.S. Environmental Protection Agency
GIS	Geographic Information Systems
HUC	Hydrologic Unit Code
K	1,000 (used in scale descriptions, e.g. 1:100K = 1:100,000 scale)
LASAR	
LWC	Luckiamute Watershed Council
NRCS	Natural Resource Conservation Service (formerly SCS)
NMFS	National Marine Fisheries Service
OCSRI	Oregon's Coastal Salmon Restoration Initiative

ODFW	Oregon Department of Fish and Wildlife
ODF	Oregon Department of Forestry
ODOT	Oregon Department of Transportation
OSU	Oregon State University
OWEB	Oregon Watershed Enhancement Board
OWRD	Oregon Water Resources Department
PRISM	Parameter-Elevation Regressions on Independent Slopes Model
REO	Regional Ecosystems Office
SMORPH	A landslide risk model used in this assessment
STORET	EPA's STOrage and RETrieval database
USFS	U.S. Forest Service
USGS	U.S. Geologic Survey

3.2 DATA

3.2.1 USE RESTRICTIONS

Some of the data used in this report were given to us with the condition that they would not be distributed. Specifically, we agreed not to distribute data concerning rare, threatened and endangered species that we acquired from the Natural Heritage Program. These data will be kept in the LWC office. In addition, we obtained the Coastal Landscape Modeling and Analysis Study (CLAMS) land cover data from researchers at Oregon State University. These data cannot be distributed by LWC. Persons wishing to use the CLAMS data should contact the CLAMS directly. CLAMS researchers are in no way responsible for our use of or conclusions drawn from our use of their data. Other data sets may also have use restrictions; we recommend that the LWC coordinator be contacted before data are distributed.

3.2.2 USE OF EXISTING DATA

Wherever possible we used uniform scale data that covered the entire study area. In some instances, data were only available for portions of the study area or stream network (e.g., AHI data). In these cases, we used our judgment to determine if enough data were available to summarize the condition of a 7th field watershed. In cases where there were not enough data to rank a watershed, we left the 7th field unranked. In other cases where there was only partial coverage, we indicated such on the map and in the report. In some cases, as with the Aquatic Habitat Inventory data, river reaches were ranked.

3.2.2.1 Scale of Data and Units of Comparison

This watershed assessment was performed using existing 1:24,000 data sets, unless otherwise noted. This is the scale of a 7.5' USGS topographical map. In this assessment we provide descriptive information on species distributions, and general watershed characteristics and processes. With an eye toward action planning, we have used GIS to highlight areas for consideration by future monitoring and restoration planning. Since we relied almost exclusively on existing data, some of our prioritizations were constrained by those data sets. In particular, LWC requested that we prioritize stream reaches using the aquatic habitat inventory (AHI) data. Since the existing AHI data set did not cover the entire study area, we prioritized individual stream reaches, as defined by the data collection agency, in this assessment. Therefore, the units of comparison for the AHI summaries

are individual stream reaches. In other summaries, the 7th field watersheds are the units of comparison.

3.2.3 ACCURACY & UNCERTAINTY

The relationships between the various salmonid species and the watersheds they inhabit are extremely complex. Generally, we assume that “given enough research and the right models, or other analytical approaches, exact numbers can be determined for population size, components of population dynamics, and the responses of populations to given harvest levels... This assumption is nearly always erroneous” (Botkin *et al.*, 1993). Botkin *et al.* have identified three sources of environmental uncertainty: (1) incomplete information regarding the current state of a resource; (2) incomplete information on details of cause and effect relationships; and, (3) intrinsic unpredictability in nature. Since most population estimates are based on a relatively small sample, appropriate sampling methods and interpretation of results allow one to estimate the amount of uncertainty associated with each sample and to develop an understanding of causal relationships.

Both **accuracy** and **precision** are important considerations in making any measurement; generally, as accuracy and precision go up, so do the costs. Accuracy tells us how well our measurements reflect the condition of a variable (e.g., how many salmon there actually are in the watershed in which we are interested). Precision tells us how repeatable our measurement is time after time. You can have measurements that are precise and not accurate, ones that are accurate and not precise, and ones that are neither accurate nor precise. Statistics are used to assess accuracy and precision. Of course, the accuracy and precision of anecdotal observations cannot be known (in a statistical sense).

Making management decisions based on observations or measurements that do not accurately describe watershed conditions may produce unexpected results.

4 WATERSHED CHARACTERISTICS AND RESOURCES

Section 4 of this report describes the status of important watershed characteristics. More detailed summaries and analyses are presented in later sections.

4.1 LOCATION

The study area is located in the heart of the Willamette Valley approximately 62 mi south of Portland, Oregon and is composed of the Luckiamute, Ash Creek, American Bottom, and Duck Slough watersheds (Map 1). The study area stretches across Polk and Benton Counties, bounded to the west by the Coast Range Mountains and to the east by the Willamette River. The 235,500 acre (95,300 hectare) study area falls within two major watersheds, the Luckiamute and Rickreall 5th field HUCs. Fifty-seven 7th Field HUCs are contained within the study area (Map 2). The Luckiamute watershed consists of the Luckiamute River, Little Luckiamute River, Soap Creek, and their tributaries. At the southern end of the study area, Soap Creek flows into the Luckiamute River about 3mi before reaching the Willamette River near the Benton-Polk County line. Duck Slough and American Bottom, also in the Luckiamute watershed, are adjacent to the Willamette River just south of the town of Independence. Ash Creek, technically a part of the Rickreall watershed but added to the study area for this assessment, joins the Willamette River at the town of Independence. Ash Creek itself was

reportedly created by ditching and may not represent a natural drainage (M. Cairns, personal communication). Elevations in the watershed range from 150 feet above sea level at the Willamette River to 3,333 feet at Fanno Peak in the Coast Range (Taylor *et al.*, 2003).

4.2 SETTLEMENT

4.2.1 HISTORIC SETTLEMENT

Archeological information provides insight into where native people settled in the Willamette Valley (see Appendix B: Timeline). Artifacts and burial sites throughout the Willamette Valley reveal tools, food remains, and other indications of native culture and lifestyle. Hager's grove, near Salem has artifacts associated with charcoal-filled fire hearths and earth ovens. Other artifacts include narrow-pointed arrowheads from around 2500 BP, apparently used for arrows, and charred camas bulbs, hazelnuts and acorns (Aikens, 1992). Archeologists conclude that this site was used as a seasonal hunting site, probably used during midsummer or fall, where game was hunted and plants were collected (Aikens, 1992). With information from many sites like this, archeologists have reached some major conclusions about indigenous peoples in this area. Indigenous peoples of the Luckiamute Valley lived in small, independent groups, but belonged to the larger Kalapuyan family of peoples who occupied the Willamette Valley. In the Luckiamute Valley, there were probably six different bands (Ruby 1992) who were speakers of Central Kalapuyan, one of the three Kalapuyan languages (Aikens 1992). These people made seasonal camps within their individual ranges, so that groups could harvest various resources as they ripened or were most readily obtained (Aikens, 1992). Hunting occurred primarily in the fall, camas root was harvested throughout the summer, and fishing occurred mainly in spring, fall, and winter.

As in other places in the United States, many native peoples were killed by diseases introduced by European-American settlers. Even before 1812, when contact with fur-traders and explorers began, aboriginal populations were being decimated by European-American diseases such as smallpox and malaria (Whitlock, 19XX). Arrival of European-American settlers and the policies of the federal government further displaced aboriginal people from their homelands. The Luckiamute Indians were "relocated" twice and, in 1855, moved to a reservation designated by the U.S. government in Grand Ronde, just north of the Luckiamute homelands. The Luckiamute Indians lived on this reservation along with people from other tribes until the reservation was dissolved in 1957 (Ruby and Brown, 1992). The Donation Land Act of September 29, 1850 gave incentive for Americans to move to the West. Settlers were given land, provided that they live on and cultivate them. A man was offered 640 acres if married and 320 acres if single (Benton County Historical Museum: see Section 4.4.5.2: Mills).

4.2.2 CITIES

The Luckiamute/ Ash Creek study area includes the cities of Monmouth, Independence, Dallas, and Falls City, as well as the communities of Hoskins, Airlie, Kings Valley, Pedee, and Suver (Map 1). Major cities in the region, but outside of the study area, are Corvallis and Albany to the south and Salem to the northeast.

McArthur (1992) provides information on the naming of cities within the study area (see also Section 4.4.5.2: Mills):

Adair Village- Adair Village lies just south of the southern border of the Luckiamute Watershed on state highway 99W. Adair Village was named for Camp Adair, the military installment during World War II that resided at the same location. When Camp Adair was disbanded after the war, some of the buildings were sold to Oregon State University to be used as student and faculty housing. In 1947, a local government was established. The University gave up the lease after the post-war college enrollment boom subsided and Adair Village was run as a radar station by the Air Force from 1957 to 1969. The property was sold and individual houses were put on the market when the Air Force ceased operations, and Adair Village was incorporated in 1976 (see Section 4.4.5.5 for more information on Camp Adair).

Airlie- This town was established at the terminus of a narrow gauge line of the Oregon Railroad Company. The railroad track was removed in 1929, but the community of Airlie remained.

Buena Vista- The land for this community was donated from the land claim of Reason B. Hall. Buena Vista received its name in 1850, and was named thus because one of Hall's relatives fought in the battle of Buena Vista in Mexico.

Dallas- The town of Dallas was initially called 'Cynthia Ann'; it was settled in the 1840s but moved more than a mile south in 1856 due to an insufficient water supply. The town was later named for George Mifflin Dallas, a vice-president of the United States under Polk. Dallas was chosen over Independence as the county seat of Polk County after securing a narrow gauge railroad into the town at the cost of \$17,000 in 1878-1880.

Falls City- This town was named for the falls in the Little Luckiamute River, on the western edge of this community. The place originally served as a post office named Syracuse, which was established in February 1885. The name of the post office was changed to Falls City in October 1889.

Hoskins- The town of Hoskins was named after the fort, Fort Hoskins. The Fort was established on July 26, 1856 by the federal government. The Fort was originally established to oversee the "resettlement" of western Oregon native peoples to the newly established Siletz Indian Reservation. The fort was an important military post, but never a major military establishment. The location of the fort was on the Luckiamute River near the mouth of what is now Bonner Creek, on land that was probably owned by Rowland Chambers. The fort stayed at the original location until its closure on April 13, 1865. The present community of Hoskins is close to the site of this fort. The property is now largely wooded and undeveloped, with an old farmhouse (circa 1870) and the ruins of several farm buildings and a schoolhouse.

Independence- This city was founded by Elvin A. Thorp, from Missouri, who named it for Independence, Missouri. Thorp acquired the land for the community from a donation land claim.

Kings Valley- This community was named for the pioneer Nahum King who arrived in Oregon in 1845. A flourmill was built at this site by Rowland Chambers in 1853. Kings Valley post office was established on April 13, 1855.

Lewisville- Lewisville was established on the donation land claim of and named for David R. Lewis, a pioneer living around 1845.

Monmouth- This city was settled by a group of pioneers from Monmouth, Illinois who arrived in 1852. The same party gave 640 acres of land on which to establish a town and college. This college, originally known as Monmouth University, later became Christian College. In 1883, the Oregon Legislature passed a bill creating the Oregon State Normal School, which was later renamed the Oregon College of Education, then Western Oregon State College, and today is known as Western Oregon University.

Pedee- The Pedee community is near the mouth of Pedee Creek, a tributary to the Luckiamute River. Pedee creek was named by Colonel Cornelius Gilliam who came to Oregon in 1848 from North Carolina, home of its own famous Peedee River.

Suver- This community is named for the pioneer Joseph W. Suver who was born in Virginia in 1819 and settled on a donation land claim in the area in 1845.

4.2.3 HUMAN POPULATION

During the period of time from 1990 to 2000, Polk County’s population grew by about 22% and Benton County’s population by about 9%, with the state of Oregon growing by 18% for this same period (U.S. Census Bureau, 2003). Benton and Polk Counties show a similar pattern of growth throughout the late 19th and 20th Century. Benton County’s population grew from 3,065 in 1860 to 77,926 in 2000, and Polk County grew from 3,616 in 1860 to 63,679 in 2000. (U.S. Census Bureau Population Division, 1995; University of Virginia Geospatial and Statistical Data Center, 1998; U.S. Census Bureau, 2003).




Figure 4.2.3. Human population trends in Benton and Polk Counties from 1860-1990 (U.S. Census Bureau Population Division, 1995; University of Virginia Geospatial and Statistical Data Center, 1998; U.S. Census Bureau, 2003).

In the ten year span from 1990 to 2000, most of the cities in or near the study area saw an increase in population, with the exception of Adair Village. Table 4.2.3 contains the population and population change for several of these cities (U.S. Census Bureau, 2002). Significant increases were seen in all cities within the study area. Falls City's population increased by about 17%, Independence increased by approximately 27%, and Monmouth increased by almost 19%.

City	1990 Population	2000 Population	% Change
Adair Village ³	548	536	-2.2%
Albany ³	29,463	40,852	27.9%
Corvallis ³	44,757	49,322	9.3%
Dallas ²	9,422	12,459	24.4%
Falls City ¹	800	966	17.2%
Independence ¹	4,425	6,035	26.7%
Keizer ³	21,884	32,203	32.0%
Monmouth ¹	6,288	7,741	18.8%
Salem ³	107,786	136,924	21.3%

¹ City is within the study area, ²City is partially within the study area, ³City is near the study area and information is provided for comparison.

4.3 TRANSPORTATION

4.3.1 ROADS

Several major highways pass through the study area. Oregon state highway 99W runs in a north-south direction through the study area along the study area's eastern perimeter. Oregon State Hwy 223, otherwise known as Kings Valley Highway, crosses through the study area in a north-south direction in the western portion of the study area. The Monmouth Highway runs from Monmouth in an east-west direction between the two other major highways (Map 1).

Knowledge of the type and location of roads is important for a watershed assessment (see Sections 6.4.2.2 and 8.1.3). For example, roads located in floodplains and roads that cross streams can directly affect hydrologic patterns by constraining stream channels. Indirectly, road building replaces permeable soils with impervious surface so that instead of slowly infiltrating soils, water runs along road surfaces and enters the stream network over a short period of time. In extreme cases, roads have actually functioned as extensions of the stream network during storm events (Wemple, 1994). Since roads also act as barriers for many types of wildlife and transportation corridors for invasive species, roadless areas are also important to understand of how the watershed functions.

The Watershed Assessment Manual (Watershed Professionals Network, 1999) suggests that when impervious surfaces of roads cover 4% to 8% of a watershed's area, there is a moderate to high risk

of alteration to hydrologic peak flows. Of course, as water moves over a road surface it can transport pollutants and sediments to the stream network. A study by an independent group of scientists reported that roads could be a chronic source of sediments to streams (Independent Multidisciplinary Science Team, 1999).

In order to determine the risk of sediment (and pollutant) delivery and stream channel constraint that roads pose to each watershed, it is necessary to map and categorize roads throughout the basin. This would include classification of roads into paved and unpaved categories, and determination of road width. If road densities were to be calculated, a uniform-scale map of roads is necessary. There is a total of 1,432.2 miles of roads in the study area, with the majority of the roads with a known road surface having an aggregate surface (16.5%) (Table 4.3.1).

Road Surface	Length (mi)	% Total Length
Aggregate Base ASC - Aggregate Surface	236.5	16.5%
Bituminous	41.0	2.9%
Hard Surface	36.1	2.5%
Natural Unimproved	8.8	0.6%
Not Designated	3.4	0.2%
Not Known	1061.9	74.1%
Pit Run	44.5	3.1%
Total	1432.2	100.0%

4.3.2 RAILROADS

Railroads have been important in the area since the early 1900s. The Valley & Siletz Railroad, which was incorporated in 1912 by the Cobbs & Mitchell Lumber Company, passed through the watershed along the Luckiamute River linking Kings Valley and Pedee with Independence. It connected to the Southern Pacific Railroad at Independence. The primary goal of its construction was to move timber, but it was also used to move agricultural products. The line was purchased in late 1984/early 1985 by the Willamette Valley Railroad Company. The Willamette Valley Railroad Company continued to operate carrying cargo for the Mountain Fir Lumber Company until the lumber facility closed in May of 1992. This railroad is no longer in use (<http://www.pnwc-nrhs.org/rr-history/rr-history-VS.html>).

4.3.3 AIRPORTS

There are several airports in the study area. The Independence State Airport is located north of the City of Independence. McNary Field in Salem is the closest regional airport, and the closest international airport is located in Portland.

4.3.4 PORTS

The Buena Vista Ferry runs across the Willamette River from Independence/Buena Vista (<http://publicworks.co.marion.or.us/operations/ferries/bvinfo.asp>)

4.4 LAND COVER, VEGETATION AND LAND USE

Land cover, vegetation, and land use are related to and affect the placement and condition of one another (Fig 4.4). Land cover is the type of vegetation, vegetation communities, and land features present on the landscape. Vegetation communities are the groupings of specific plants, which commonly occur together, into functional units, such as wetlands and riparian areas. Plant species within a 'true' biological community interact with one another. Land use, or the way the landscape is used by humans, affects land cover because, through specific uses of the land, the vegetation and vegetation communities of the landscape are changed, as well as the land features present on the landscape. Land use can be affected by land cover if the land features are too extreme to be manipulated. For example, one would most likely leave a lava field in place rather than attempt to move it to farm in that area. Vegetation and vegetation communities are a component of land cover, and are also affected by it. For example, a type of land cover could be a rock outcropping that vegetation could not grow on.




Figure 5.4.1. Diagram showing relationship between Land Cover, Land Use and Vegetation.

The sections below detail land cover, vegetation and vegetation communities, and land use in the study area.

4.4.1 LAND COVER

Land cover is an important ecological characteristic to consider in a watershed assessment because land cover affects watershed characteristics such as potential for large wood delivery to a stream network, erosion or landslide potential, wildlife habitat potential, susceptibility to rain-on-snow events, riparian shade, amount of urbanization within the study area, and the location and amount of water present. Aquatic, riparian, wetland, and upland land cover classes are four major groupings that have a strong affect on the ecological processes and quality of the watershed. For this watershed analysis, we relied on two existing data sets which describe land cover/ land use for the Luckiamute and Ash Creek study area. For more information on the data sets used to describe these watershed features, please see Appendix C: Data Dictionary.

4.4.1.1 Historic Land Cover Conditions

Analysis of pollen and written records from early settlers and surveyors allow us to characterize the vegetation in the region from those earlier times.

Pollen studies show that after the peak of the last glaciation, the climate of the Willamette Valley shifted from cool and wet to warmer and drier. There was a time of transition between 9,000 and 7,000 yrs before present (BP) when trees which thrived in cool, wet conditions, such as Sitka spruce and white pine, declined in number. By 4,000 years BP, there was an even greater decrease in the abundance of cool-climate species, and an increase in the abundance of Douglas-fir and ponderosa pine, which thrived in warmer, drier conditions. White Oak, common to drier climates, increased greatly in abundance during this period (~4,000 BP). After 4,000 BP, the climate became slightly cooler and wetter again, leading to the establishment of forest conditions like those seen today, with Douglas-fir forests and some ponderosa pine on the surrounding hills, and oak and other deciduous species on the valley floor (Aikens, 1986).

Information regarding more recent history can be inferred from the archaeological and historical record. Archaeological information describing the Kalapuyan Indians' use of plants indicates that hazelnut, Oregon grape, salmonberry, elderberry, and ninebark were present in the study area before European settlement (Aikens, 1992).

Descriptions by European-American settlers also give a glimpse of what the Willamette Valley looked like in the years before present. The settlers began arriving in the Luckiamute valley in the mid 1800s, drawn by promises of free land, fertile soils, and a mild climate. One of the earliest European-American settlers to the Luckiamute valley, Anna King, described her impression of the landscape as follows:

“It is a beautiful country as far as I have seen... Soda springs are common and fresh water springs without number. It is now the 1st of April and not a particle of snow has fallen in the valley... There are thousands of strawberries, gooseberries, blackberries, whortleberries, currants, and other wild fruits but no nuts except filberts and a few chestnuts” (Aikens, 1992).

Early cadastral surveys in the valley show the hilly regions in the southwestern portion of the watershed as forested with fir and oak. As noted by Yamhill Basin Council (2001):

Oregon white oak occurs in two forms: forest and savanna. The majority of existing trees developed under forest conditions. These “forest-form” trees are relatively tall, seldom exceed 60 centimeters (23.62in.) in diameter measured at breast height (dbh), and have ascending branches clustered near the crown. Their crowns form a closed canopy. The average age of mature forest-form trees (in 1968) was 90 years with an age range of 47—135 years. Scattered through the forest and remaining in some fields are a few large relic Oregon white oaks developed by centuries of controlled burns in non-forest conditions. These “savanna-form” trees generally exceed one meter (39.37 in.) dbh and their boles are short in relation to the total height of the tree. They have massive branches and spreading crowns and are usually spaced so the crowns do not touch. There is an average of 17 savanna-form Oregon white oaks per hectare (2.471 acres) in remnant oak savanna forests of the region. In 1968, their annual growth ring indicated an age range of about 260—310 years. Other studies indicate Oregon white oak may live over 500 years and reach 90cm (35.43in) dbh at only 250 years of age.

Also written on the maps of the early cadastral surveys are comments that the lands have poor soil, and are unsuitable for settlement (Bureau of Land Management, 1882).

Historical accounts from the early 1800s and more recent analyses indicated that open prairie was the dominant feature in the Willamette Valley before settlement (Taft and Haig, 2003). Given hydric soils in this area, it is likely that a large portion of the prairie was wet prairie. Historic accounts written before the 1880s mentioned that the prairies in the winter and spring were “wet and muddy”, “covered with water” and “too wet to plow” (Taft and Haig, 2003). These conditions could have been one of the reasons early settlers chose donation land claims in the more hilly, forested areas rather than the valley lowlands, as described in (Barnhart, 1915). By 1871, after continued settlement and the introduction of railroads along the valley floor, farming activities began to increase in the valley lowlands (Taft and Haig, 2003). In order to improve the ability to farm this area, the land was drained through surface ditching between 1860 and 1880. By the early 1900s, more lands were drained by tile drains, as advocated by the State of Oregon (Taft and Haig, 2003). Thus, land use changes between 1800 and 1900 dramatically altered the extent of wet prairies and bottomlands, and facilitated the establishment or spread of several non-native plant species (see Section 4.4.2.1).

Analysis of the PNERC pre-settlement vegetation layer indicated that the coniferous forests dominated the western, higher elevations of the watershed and a large area along the south central boundary of the Luckiamute/ Ash Creek study area (Map 5). Coniferous forests covered approximately 31.9% (76,000 acres) of the study area. Interestingly, shrubby areas were just as common as coniferous forests. Upland and wetland shrubby areas covered much of the eastern, low areas in both the Luckiamute and Ash Creek watersheds covering about 76,114 acres (32.0% of the study area). Much (53,674 acres or ~22.5% of the study area) of the remaining area was dominated by herbaceous plant communities, wet prairies and natural grasslands. Mixed and deciduous forests (which included Oak Savannah) covered 7.3% and 5.8% of the study area, respectively.

We used the PNERC pre-settlement vegetation GIS layer to summarize vegetation in each of the 7th field watersheds in the study area (Table 4.4.4.1).

Table 4.4.1.1. Major habitat cover classes for each of the 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are the 7th field watershed IDs and the number of acres for the water, mixed forest, shrub, deciduous and coniferous forest and herbaceous cover classes. Derived from the PNERC pre-settlement vegetation data layer.

7th Field Watershed ID	Water	Forest (unspecified or mixed)	Shrub	Forest (deciduous)	Forest (conifer)	Herbaceous
17090003060204	0.0	1,554.8	2,026.4	151.0	1,725.8	1,837.1
17090003061101	0.0	140.2	1,258.9	66.6	4,373.8	144.9
17090003060302	0.0	427.7	149.0	0.0	1,856.7	346.1
17090003060303	0.0	637.4	282.4	5.2	2,012.9	455.0
17090003061102	0.0	0.0	3,152.7	3.6	71.1	1,177.0
17090003060301	0.0	651.8	756.7	138.6	316.4	1,564.7
17090003061302	41.4	0.0	1,904.6	37.8	0.0	2,212.0
17090003061303	2.5	0.0	1,612.6	396.0	0.0	905.2
17090003061103	0.0	2.3	1,155.6	454.5	189.9	1,310.2
17090003060304	0.7	1,239.1	1,111.5	2.5	2,195.6	1,368.5
17090003061201	2.9	961.7	1,010.3	12.4	1,157.4	370.8
17090003060305	0.0	1,577.9	1,093.3	114.8	323.8	1,142.8
17090003061202	0.0	0.0	1,865.5	35.6	0.0	607.7
17090003060203	0.5	1,094.4	1,509.8	69.1	3,994.2	621.7
17090003061301	3.8	0.0	2,578.7	104.4	0.0	1,908.2
17090003061204	0.0	0.0	1,456.9	108.9	0.0	1,793.9
17090003060501	0.0	721.1	556.4	654.8	269.6	472.7
17090003061005	155.3	0.0	1,449.5	746.6	0.0	995.9
17090003061203	0.0	0.0	2,180.7	63.7	0.0	1,502.3
17090003060202	0.0	0.0	78.8	178.7	5,422.3	0.0
17090003060504	1.8	832.3	2,922.3	392.2	499.1	959.2
17090003061003	103.5	0.0	1,271.3	783.2	0.0	721.4
17090003060403	0.0	831.4	0.0	0.0	1,507.5	132.1
17090007020404	260.1	0.0	795.4	110.5	0.0	835.7
17090003060404	0.0	5.2	763.7	1.1	5,193.5	479.0
17090003060503	19.6	0.0	4,669.4	1,224.0	0.0	2,516.6
17090003060401	0.0	1,592.8	1,590.3	11.0	1,820.3	1,098.2
17090003060201	0.0	0.0	0.0	0.0	3,267.7	0.0
17090003061002	0.2	0.0	996.5	12.8	0.0	1,415.9
17090003060502	0.0	1,311.8	2,612.3	322.9	195.1	1,066.1
17090007020501	319.7	0.0	48.2	1,219.1	0.0	980.3
17090003060102	0.0	0.0	0.0	0.0	5,677.9	0.0
17090003060402	0.0	2.7	687.4	0.0	3,969.0	21.8
17090003061001	74.3	0.0	2,510.3	299.0	0.0	1,579.5

Table 4.4.1.1. Major habitat cover classes for each of the 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are the 7th field watershed IDs and the number of acres for the water, mixed forest, shrub, deciduous and coniferous forest and herbaceous cover classes. Derived from the PNERC pre-settlement vegetation data layer.

7th Field Watershed ID	Water	Forest (unspecified or mixed)	Shrub	Forest (deciduous)	Forest (conifer)	Herbaceous
17090003060101	0.0	0.0	0.0	0.0	5,980.1	0.0
17090003060803	0.0	266.0	870.1	82.6	705.4	137.9
17090007020606	0.0	0.0	1,762.0	0.0	0.0	1,579.1
17090007020503	122.0	0.0	1,691.8	549.5	0.0	834.1
17090007020603	0.0	0.0	1,892.5	168.8	0.0	1,427.2
17090003060801	0.0	0.0	0.0	0.0	2,232.9	0.0
17090003060902	4.5	787.3	1,769.4	1,231.7	0.0	2,758.3
17090003060602	0.0	0.0	0.0	0.0	1,811.3	0.0
17090003060802	0.0	455.9	785.9	54.5	3,049.9	638.1
17090003060903	0.0	0.0	3,713.9	25.9	0.0	1,219.5
17090007020605	1.6	0.0	3,033.0	527.9	0.0	2,214.0
17090003060702	0.0	468.5	621.9	0.0	1,556.8	709.9
17090003060603	0.0	0.0	0.0	0.0	3,421.6	0.0
17090007020604	0.0	0.0	2,225.7	85.1	23.6	2,124.5
17090003060901	0.2	0.0	3,301.4	647.1	0.0	1,121.2
17090003060704	0.9	75.4	475.0	2,021.0	0.0	1,015.4
17090003060601	0.0	0.0	0.0	0.0	6,064.0	0.0
17090003060701	0.0	430.0	0.0	0.0	4,586.0	7.9
17090003060703	0.0	818.6	543.8	122.0	532.1	459.5
17090007020602	0.0	0.0	2,642.9	9.7	0.0	1,009.8
17090007020601	0.0	420.3	1,735.0	318.2	0.0	314.3
17090007020502	4.1	0.0	1,598.2	255.6	0.0	1,127.9
17090003061004	4.1	0.0	1,395.5	1.8	0.0	2,434.3

Historic Wildfire Conditions

Wildfires are part of the natural disturbance regime of the region. When considering the impact of fire on an ecosystem, it is important to consider the extent, frequency, timing, and intensity of the burn in relation to the natural range of variability. Fires create gaps important to wildlife and ecosystem processes in the forest canopy. Fires also impact watershed hydrology by altering watershed water storage capacity, erosion rates, nutrient and carbon mobilization, and soil permeability. Fire can also have an indirect impact on watersheds. For example, salvage operations that occurred in the Tillamook basin following the Tillamook Burn had a huge impact on stream network channel morphology because fallen trees were dragged out of the watershed using the streambeds. Much of what we know about wildfires comes from soil profiles (pollen & charcoal) and from the accounts of witnesses.

Climate, vegetation, and fire regimes were closely interlinked in prehistoric times, as they are today. Changes in climate affected which types of vegetation dominated the landscape, and how frequently fires burned through the area. When the climate was historically warmer and drier, fires appear to have been more frequent and drought and disturbance-adapted species more widespread (Whitlock and Knox, 2002).

Historically, fires in upland forests were likely governed solely by weather patterns and natural ignition. High intensity, stand-replacement fires occurred in the Coast Range at irregular intervals of 150-400 years. This is based on an analysis of vegetation in the area which showed that late-seral and old growth forests historically occupied 60 to 80% of the Coast Range landscape (Licata *et al.*, 1998). Very little is known about the frequency and extent of lower intensity fires (referred to as “under- burns”) in the northern Coast Range [(Walstad, 1990)cited in (Licata *et al.*, 1998)]. There are few detailed historic accounts of low-intensity fires and generally there is little physical evidence that can be used to determine the frequency and intensity that persists more than a few years following an under-burn. Licata *et al.* (1998) write,

“The influence of on-shore flow of marine air masses creates a predominantly cool and moist climate in the Coast Range, making the incidence of lightning strikes in this region one of the lowest in North America. This prevailing climatic condition is the primary reason for the infrequent nature of both major fires and underburns. It is hypothesized that human-caused ignitions played a more significant role in fire occurrence in the Coast Range compared with other areas of the state (Teensma *et al.*, 1991).”

Humans have been part of the Pacific Northwest landscape for at least 8,000 years and probably closer to 11,000 years (Aikens, 1993; Licata *et al.*, 1998; Whitlock and Knox., in press). Historical accounts and archeological records indicate that the Kalpuyan peoples periodically burned the meadows of the valley floor to facilitate hunting large game, clearing meadows for the harvest of camas, a major staple to their diet, and promoting growth of seed-producing grasses (Aikens, 1993; Williams, 2002). These burnings may have occurred as frequently as several times a year or in intervals of every 5-years (Licata *et al.*, 1998; Williams, 2002). In the spring there probably was no burning when Native Americans were concentrated on flood plain sites in the Willamette Valley wet prairies. During mid-summer (July and August), encampments probably shifted to drier prairie sites. Burning was sporadically initiated during July and August following the harvest of grass seeds, sunflower seeds, hazelnuts and blackberries (used as food) in limited areas. These burns would promote the re-growth of vegetation. Fire was used differently during the late summer. Burning facilitated the collection of tarweed and insects on the high prairies. In the fall, oak openings were burned following the harvest of acorns. The Kalapuya initiated large-scale communal drives for deer, which provided a winter’s supply of venison by burning areas along the valley edge. If late summer and fall fires were ignited prior to the onset of strong east winds, it seems likely that fires would have burned up into the higher elevations of the Coast Range (Ripple 1994, (Teensma *et al.*, 1991)). Pushed by a strong east wind following a very dry summer, it is not difficult to envision a late summer fire, started at a valley margin site, burning well into the interior of the Coast Range before weather conditions changed and halted its advance. Historic Fire Patterns Historic fire patterns, and their effects on the landscape pattern of the Coast Range, have become an item of considerable interest to many authors (Zybach 1988, Walstad *et al.* 1990, Teensma *et al.* 1991,

Agee 1993, Ripple 1994). However, there is still debate over how wide-spread these controlled fires were and how much of an impact they actually had on vegetation composition in the valley (Aikens, 1993; Williams, 2002; Whitlock and Knox., in press).

Logging practices during the past 150 years have shifted the vegetation towards younger seral stages (< 80 years-old). Direct changes to the vegetation through logging may have altered fire regimes. Early European-American settlers' activities may have also altered fire regimes through direct ignition. The settlers introduced new fire-ignition sources to the landscape including logging, mining, and road-building activities, as well as burning on a large scale to remove trees and to prepare agricultural lands (Licata *et al.*, 1998; Whitlock and Knox., 2002). Licata *et al.* (1998) reported that, "Rapid response to extinguish all fires and discontinuous arrangement of fuels, due to clear cutting, slash burning and road construction, has kept most fires small. Thus, there are hundreds of small (less than 100 acres) disturbance areas throughout the analysis area."

A very large wildfire, or series of fires, burned approximately 480,000 acres of the Central Coast range in the period between 1853 to 1868 (Licata *et al.*, 1998). According to a BLM spatial data set describing the fire history of the northwestern Oregon forests at 30-40 year intervals from 1850 to 1940 (U.S. Bureau of Land Management, 1996), an area 20,200 acres in size in the Luckiamute watershed study area had been burned prior to 1850 and had not been reforested as of 1850. Between 1850 and 1890, three areas sized 290, 600, and 1,120 acres in the Luckiamute watershed study area were burned and had not been reforested as of 1890. Between 1890 and 1920 there were two areas that had not recovered from a fire. Because these two areas are the 290 and 1,120 acre areas that were burned prior to 1890, it is possible that either they were the same areas burned prior to 1890 and had not yet been reforested, or they were both burned twice. The last data set describing fire history in 1940 showed no burned areas. These data sets did not extend into the Ash Creek study area.

4.4.1.2 Current Land Cover Conditions

We acquired current land cover/land use data for the study area from the Institute for a Sustainable Environment. Land cover/land use in the study area is dominated by forested and agricultural areas (Table 4.4.1.2a, Map 4).

Table 4.4.1.2a. Land cover/Land use for the study area as summarized from ec90 (Institute for a Sustainable Environment, 1999).	
Land Cover/Land Use	Area (acres)
Agriculture	72,721.5
Forested	133,144.2
Industrial/Commercial	281.8
Flooded/Marsh	704.3
Natural grassland	3,120.7
Natural shrub	15,572.2
Residential	1,692.8
Rural	3,002.6
Urban	654.8
Roads	2,601.7

Water	1,072.9
Bare/fallow	860.3
Topographic Shadow	14.0

Digital orthophotos are another important resource for understanding and interpreting land cover. For this project, digital orthophoto quads (DOQs) were obtained from the Oregon Geospatial Data Clearinghouse (formerly the State Service Center for GIS). This imagery was collected in 2000 and covers the entire study area with a resolution of 1m. A digital orthoquad is an aerial photograph, which has been corrected for the horizontal displacement that occurs when using aerial photography and has been spatially referenced for use in a GIS. Because the data are spatially referenced photographs, the DOQs give a bird’s eye view of the entire study area, in this case, as it looked in around 2000. Because they are spatially referenced, other GIS data layers, such as roads or vegetation, can be placed “over” the DOQs in a GIS for mapping or data verification (Fig. 5). These datasets can be very useful for detecting specific types of land cover such as riparian areas and infrastructure (see Section 5.3: Riparian Analysis)

We summarized existing land cover from the PNERC LULC_90 data set (Table 4.4.1.2b). We grouped 60 cover classes into nine major cover classes (and a background class). These data were generated from researchers using satellite imagery, aerial photographs, and other data sources. These data were last updated in 1999. We found that coniferous forests dominate the current Luckiamute and Ash Creek watersheds, covering ~76,153 acres or about 32% of the study area. Interestingly, current coniferous forests cover approximately the same extent as pre-settlement coniferous forests (see Section 4.4.1.1). Herbaceous cover classes, including crops, hay fields, pasture, hops and mint fields, as well as natural grasslands, are the second largest cover class in the study area covering 70,623 acres. Mixed and deciduous forests, shrubs and urban areas account for 42,550 acres, 19,606 acres, 18,185 acres, and 8,210 acres, respectively.

Table 4.4.1.2b. Current major habitat cover classes for each of the 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are the 7th field watershed IDs and the number of acres for the coniferous forest, herbaceous, mixed forest, deciduous forest, shrub, urban, water, non-vegetated open, and wetland cover classes. Derived from the PNERC LULC 90 vegetation data layer.

7th Field Watershed ID	Forest (conifer)	Herbaceous	Forest (unspecified or mixed)	Forest (deciduous)	Shrub	Urban	Water	Non-vegetated open areas	Wetland	Background
17090003060204	4,025.5	427.5	2,064.6	329.4	385.9	51.1	4.3	3.8	2.7	0.2
17090003061101	2,770.4	119.7	2,455.9	456.3	157.7	17.6	0.5	4.1	2.05	0.2
17090003060302	892.6	307.1	1,165.3	265.5	133.7	11.9	0.2	1.4	1.1	0.7
17090003060303	1,186.4	333.0	1,324.1	373.5	142.2	13.1	0.2	19.1	1.1	0.0
17090003061102	1,691.1	1,165.3	761.2	254.5	441.9	66.6	0.0	6.8	17.1	0.0
17090003060301	913.1	830.7	915.8	213.3	402.1	98.1	0.2	47.7	7.2	0.0
17090003061302	31.5	2,826.5	50.6	324.2	700.4	202.5	3.2	18.0	38.9	0.0
17090003061303	32.6	2,021.4	59.9	213.5	295.9	224.8	14.6	3.8	49.5	0.2
17090003061103	815.0	1,398.2	308.5	271.8	266.0	34.0	0.0	3.2	16.0	0.0
17090003060304	2,049.5	755.3	2,001.8	504.0	440.3	91.6	9.2	57.6	7.9	0.5
17090003061201	1,753.2	455.4	1,088.8	124.7	65.9	14.4	2.9	6.8	3.2	0.2
17090003060305	2,066.4	444.6	938.0	249.8	422.3	72.2	0.0	49.1	10.1	0.0
17090003061202	765.5	982.4	327.2	166.1	200.7	34.4	0.0	26.3	6.3	0.0
17090003060203	3,041.8	549.2	2,827.8	501.3	310.1	43.4	1.6	6.5	7.9	0.0
17090003061301	62.8	3,724.7	47.9	269.3	293.0	178.0	0.2	2.5	16.9	0.0
17090003061204	220.7	2,778.3	58.3	125.8	106.0	49.3	13.5	0.7	7.2	0.0
17090003060501	1,071.7	259.9	815.6	235.8	227.9	49.5	0.7	8.8	4.7	0.0
17090003061005	3.2	2,534.4	85.1	247.3	180.0	126.7	136.6	0.5	33.5	0.0
17090003061203	137.9	2,327.6	54.5	365.0	625.1	185.2	0.5	46.6	4.5	0.0
17090003060202	2,833.4	50.4	2,079.5	646.4	62.6	5.4	0.2	3.4	0.5	0.5
17090003060504	2,091.2	1,446.5	925.2	479.5	452.3	160.7	1.4	35.1	15.1	0.0
17090003061003	15.5	1,930.3	68.0	385.9	200.5	123.3	102.4	14.9	38.7	0.0

Table 4.4.1.2b. Current major habitat cover classes for each of the 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are the 7th field watershed IDs and the number of acres for the coniferous forest, herbaceous, mixed forest, deciduous forest, shrub, urban, water, non-vegetated open, and wetland cover classes. Derived from the PNERC LULC_90 vegetation data layer.

7th Field Watershed ID	Forest (conifer)	Herbaceous	Forest (unspecified or mixed)	Forest (deciduous)	Shrub	Urban	Water	Non-vegetated open areas	Wetland	Background
17090003060403	1,139.4	60.5	983.0	230.6	50.4	5.4	0.0	0.7	0.9	0.0
17090007020404	0.5	1,071.7	44.6	320.0	226.8	118.8	194.0	6.8	18.0	0.7
17090003060404	3,924.9	242.1	1,791.2	320.9	123.3	36.2	0.0	0.7	3.15	0.0
17090003060503	1,329.8	4,615.2	818.3	520.0	769.5	316.1	0.2	15.5	45.0	0.0
17090003060401	2,861.1	619.9	1,460.0	572.2	483.5	89.1	0.2	14.6	12.0	0.0
17090003060201	1,726.7	3.8	1,154.7	370.4	9.5	0.0	0.0	1.8	0	0.9
17090003061002	53.1	1,899.5	51.1	135.5	195.8	82.8	0.2	1.6	6.1	0.0
17090003060502	2,518.0	494.1	1,546.7	399.8	404.1	119.5	0.5	17.8	7.7	0.0
17090007020501	0.2	1,361.7	70.4	331.2	535.1	52.2	193.5	0.5	21.0	1.6
17090003060102	3,530.7	4.1	1,733.2	390.6	20.0	0.0	0.0	0.7	0.0	0.5
17090003060402	2,484.2	20.0	1,791.7	351.2	32.9	0.9	0.0	0.0	0.0	0.0
17090003061001	247.3	2,662.9	260.1	701.3	411.1	63.7	72.5	17.8	26.6	0.0
17090003060101	3,674.9	11.0	1,660.3	589.1	33.5	0.0	0.0	6.1	0.0	0.2
17090003060803	993.6	68.6	751.1	144.7	96.1	5.4	0.2	0.5	0.9	0.9
17090007020606	0.0	2,334.4	14.0	74.5	223.9	679.1	0.2	7.4	7.7	0.0
17090007020503	0.0	2,103.5	46.8	183.4	197.8	398.9	224.8	18.5	23.4	0.2
17090007020603	4.1	2,293.0	51.3	152.1	250.9	639.5	86.4	5.2	6.1	0.0
17090003060801	1,919.3	0.0	266.0	46.6	0.0	0.0	0.0	0.0	0.0	1.1
17090003060902	1,061.6	3,126.6	705.6	717.3	630.9	239.0	3.6	17.1	49.5	0.0
17090003060602	1,633.5	0.0	117.9	60.3	0.0	0.0	0.0	0.0	0.0	0.0
17090003060802	2,850.1	470.3	1,061.6	260.6	229.1	91.1	0.2	9.2	10.6	1.6
17090003060903	327.8	2,153.3	452.9	918.5	908.3	150.8	0.0	15.1	32.6	0.0
17090007020605	265.7	3,203.6	359.1	827.3	829.8	238.5	2.3	36.7	13.5	0.0
17090003060702	1,230.5	457.2	531.9	263.0	347.0	508.3	0.0	14.2	4.7	0.2

Table 4.4.1.2b. Current major habitat cover classes for each of the 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are the 7th field watershed IDs and the number of acres for the coniferous forest, herbaceous, mixed forest, deciduous forest, shrub, urban, water, non-vegetated open, and wetland cover classes. Derived from the PNERC LULC_90 vegetation data layer.

7th Field Watershed ID	Forest (conifer)	Herbaceous	Forest (unspecified or mixed)	Forest (deciduous)	Shrub	Urban	Water	Non-vegetated open areas	Wetland	Background
17090003060603	2,748.2	7.0	555.1	102.6	7.0	0.0	0.0	0.9	0	0.9
17090007020604	161.8	2,375.6	194.0	547.7	596.9	532.4	12.6	27.0	11.0	0.0
17090003060901	480.2	2,080.6	271.8	833.9	1,071.2	216.9	1.6	93.4	20.5	0.0
17090003060704	628.2	1,240.0	428.2	403.2	671.2	139.1	1.6	58.7	17.6	0.0
17090003060601	4,883.9	0.0	1,035.7	142.9	0.0	0.0	0.0	0.0	0.0	1.1
17090003060701	3,733.7	45.9	1,003.5	170.6	57.8	9.7	0.0	1.8	0.0	0.9
17090003060703	784.4	302.2	574.0	323.8	384.8	77.2	0.2	26.3	3.2	0.0
17090007020602	173.3	1,737.7	69.3	347.6	634.1	659.3	0.0	32.9	8.3	0.0
17090007020601	310.3	677.3	193.5	372.2	680.4	505.4	0.2	39.2	9.5	0.0
17090007020502	0.2	2,155.1	64.8	377.6	228.2	122.9	3.4	0.0	33.3	0.5
17090003061004	1.8	3,066.5	14.0	101.5	332.8	259.2	17.3	15.5	27.0	0.0

4.4.1.3 Current Wildfire Conditions

Today, the effects of logging on fuel structure and fire ignition continue to impact fire regimes in upland coniferous forests today. Another factor influencing both fuels and ignition, both of which influence wildfire behavior, is fire suppression. Fires in the coastal forests of Oregon are rare today because the months of dry fuels and ignition are not coincident (Long and Whitlockm, 2002).

According to a BLM dataset marking the initiation points of fires from 1980 to 1989, there have been 65 fires in the Luckiamute watershed and six fires in the Ash Creek watershed (Bureau of Land Management, 1991). In 1987, the Luckiamute watershed had the greatest number of fires (Chart 4.4.1.3) and at the same time 5,000 acres burned in the Rockhouse Creek fire, located in the adjacent Rickreall watershed (Mattson and Gallagher, 2001). The next year, 318 acres were burned in wildfires, a number far exceeding the total number of acres burned for the other years in this timeframe (Table 4.4.1.3).

From 1980 to 1989, the majority of the fires reported were assumed to have been caused by lightning (41 fires); with the next most likely cause being fire control activities, i.e., backfires (9); other causes included incendiary (4), campfire (2), smoking (2), railroads (1), and miscellaneous (6). Approximately the same number of fires had a flame length of 0-2 feet with a burning index of 1-20 (32 fires) and a flame length of 2-4 feet with a burning index of 21-40. Only three fires had a flame length from 4-6 feet and a burning index of 41-60.




Figure 4.4.1.3 . Number of wildfires in the Luckiamute watershed study area from 1980-1991 (Bureau of Land Management, 1991).

Table 4.4.1.3 . Number of acres burned from 1980 to 1989 in the Luckiamute watershed (U.S. Bureau of Land Management, 1991).	
Year	Area Burned (acres)
1980	2
1981	1
1982	1
1983	<1
1984	12
1985	4
1986	<1
1987	2
1988	318
1989	1

In the Ash Creek watershed, there was one fire reported in 1986, two in 1987, and three in 1989. The largest fire, burning 13 acres, was reported in 1987. For both 1986 and 1989 the total area burned was less than one acre. All of the fires reported during this time period were assumed to have been caused by lightning. One of the fires had a flame length of 0-2 feet with a burning index of 1-20 (32 fires), with the remainder of the fires having a flame length of 2-4 feet with a burning index of 21-40 (Bureau of Land Management, 1991).

4.4.2 VEGETATION AND VEGETATION COMMUNITIES

4.4.2.1 Native Vegetative Diversity and Weeds

“Native plant diversity” is the variety of plants that occur naturally within an area. “Weeds” may be defined as non-native (mostly) plant species that flourish and invade desirable crops or native plant communities, especially when soil, land cover, and natural runoff patterns are disturbed by humans or natural events. Noxious weeds (a legal term, ORS 570.505) applies to a specific list of weeds that are considered injurious to public health, agriculture, recreation, wildlife, or any public or private property. Native plant communities – as well as croplands, roadsides, and residential areas -- are under constant siege by weeds. The proportion of local plant diversity that consists of weed species has not been tallied, but some of the more troublesome species are listed in Table 4.4.2.1.

Data Availability

No comprehensive surveys have been undertaken of plant diversity in the Luckiamute / Ash Creek study area, or of the distribution and prevalence of particular weeds, or of the extent of weed control activities. However, the Oregon State University Herbarium has

compiled a list of vascular plants that can be queried by county (http://oregonstate.edu/dept/botany/herbarium/db/vasc_plant.html). Also, the University of Montana has compiled a list of noxious weeds that can be queried by county (<http://invader.dbs.umt.edu/>), known as the Invaders Database System. Plant lists are available for the McDonald-Dunn Forest -- woody species only (OSU College of Forestry 1990), and Beazell County Park. Lists may be available for a few additional areas where jurisdictional wetland determinations have been made (see Section 4.4.2.2), for the Delbert Hunter Arboretum in Dallas, and/or for other areas visited by local botanists. Non-systematic records of rare plants are maintained and updated by the Oregon Natural Heritage Information Center (ORNHC).

Species Status and Geographic Patterns

Probably well over 1,000 kinds of native plants occur in the Luckiamute Watershed, but the status and distribution of most of these within the watershed is unknown. In the Beazell Memorial Forest alone, 246 plant species (185 of them native) were found during spring of 2001 (Kaye 2001). As of 1990, 151 woody plant species had been reported from the McDonald-Dunn Forest (Oregon State University College of Forestry, 1990). The OSU Herbarium contains specimens of about 650 species from Polk and Benton Counties.

Of particular concern are plants that are rare or believed to have declined severely or disappeared entirely from the region in recent years. The ORNHIC database includes 11 such species, subspecies (ssp.), or varieties (var.) that have been reported specifically from the Luckiamute Watershed (Table 4.4.2.1a). It is virtually certain that other rare or declining species are present in the watershed but are not included in this table because of the lack of a survey covering all watershed lands.

Scientific Name	Common Name	Listing Status*	Habitat
<i>Anemone oregana</i> var. <i>felix</i>	Bog anemone	G4T2 S1; 2	moist woods, montane meadows
<i>Cimicifuga elata</i>	Tall bugbane	G3 S3; 1; ODA=C	See: http://www.appliedeco.org/reports.html
<i>Delphinium pavonaceum</i>	Peacock larkspur	G1 S1; 1 Fed=SOC; ODA=LE	unmowed prairies, fencerows
<i>Erigeron decumbens</i> var. <i>decumbens</i>	Willamette daisy	G4T1 S1; 1 Fed=LE; ODA=LE	See: http://www.epa.gov/fedrgstr/EPA-SPECIES/1998/January/Day-27/e1851.htm and http://www.appliedeco.org/reports.html
<i>Erythronium elegans</i>	Coast range fawn-lily	G1 S1; 1 Fed=SOC; ODA=LT	upland prairie
<i>Horkelia congesta</i> ssp. <i>congesta</i>	Shaggy horkelia	G4 T2 S2; 1 Fed=SOC; ODA=C	See: http://www.appliedeco.org/reports.html

Scientific Name	Common Name	Listing Status*	Habitat
<i>Lathyrus holochlorus</i>	Thin-leaved peavine	G2 S2; 1 Fed=SOC	upland prairie
<i>Lupinus sulphureus</i> ssp <i>kincaidii</i>	Kincaid's lupine	G5T2 S2; 1 Fed=LT; ODA=LT	See: http://www.epa.gov/fedrgstr/EPA-SPECIES/1998/January/Day-27/e1851.htm
<i>Sidalcea nelsoniana</i>	Nelson's checker mallow	G3G4 S1; 2	upland prairie
<i>Wolffia borealis</i>	Dotted water-meal	G5 S1; 2	sloughs
<i>Lobaria linita</i>	(type of lichen)	G4 S1; 2	mature Douglas-fir forest with old-growth structural components

* LE= legally listed as Endangered; LT= legally listed as Threatened; SOC= Species of Concern (ODA= Oregon Dept. of Agriculture). These species are protected only on state and federal lands.
G= global status (ONHP), S= state status (ONHP). 1= critically imperiled; 2= imperiled and vulnerable to extinction; 3= rare but not immediately imperiled, 4= not rare but of long-term concern, 5= widespread and secure
Number following the semicolon: 1= extinction threatened or presumed, 2= extirpation from Oregon threatened or presumed, 3= insufficient information, 4= of concern but not immediately imperiled

In the Willamette Valley, many of the rarest native plants grow only in natural wetlands, riparian areas, oak woodlands, or prairie (Titus *et al.*, 1996). Table 4.4.2.1a lists species that often characterize oak woodlands and prairies. For additional information on composition of threatened native plant and animal communities of the Willamette Valley (especially prairies), see [Wilson \(1998a, b\)](#).

Weed Species

Weeds comprise only a small percentage of all plant species, but can severely affect most others. Weed species listed as “noxious” by the Oregon Department of Agriculture which are known to occur (or are likely to occur) in the Luckiamute Watershed are shown in Table 4.4.2.1b. Note that some of these species have positive as well as negative attributes. For example, Scotch broom has colorful flowers, may control erosion on steep slopes, and benefits soil productivity by fixing nitrogen. Fruits of the Himalayan blackberry are used by wildlife as well as people, and dense thickets formed by this species may provide sensitive breeding birds with isolation and small mammals with cover.

Scientific Name	Common Name	Habitat
<i>Abutilon theophrasti</i>	Velvetleaf	croplands, lawns, pastures
<i>Alopecurus myosuroides</i>	Blackgrass	croplands, lawns, pastures
<i>Carduus pycnocephalus</i>	Italian thistle	croplands, lawns, pastures
<i>Centaurea pratensis</i>	Meadow knapweed	croplands, lawns, pastures

Table 4.4.2.1b. Plants known or likely to occur in the Luckiamute Watershed and legally designated as “noxious weeds” by Oregon Department of Agriculture.

Scientific Name	Common Name	Habitat
<i>Cirsium arvense</i>	Canada thistle	croplands, lawns, pastures
<i>Convolvulus arvensis</i>	Field bindweed	riparian, woodland edge
<i>Cuscuta pentagona</i>	Field dodder	croplands, lawns, pastures
<i>Cytisus scoparius</i>	Scotch broom	woodland edge, clearcuts
<i>Daphne laureola</i>	Spurge laurel	woodland edge
<i>Digitaria sanguinalis</i>	Large crabgrass	croplands, lawns, pastures
<i>Hedera helix</i>	English ivy	riparian, woodlands
<i>Lythrum salicaria</i>	Purple loosestrife	wetlands, riparian
<i>Orobanche minor</i>	Small broomrape	woodland edges
<i>Phalaris aquatica</i>	Aquatic canarygrass	wetlands, riparian
<i>Phalaris arundinacea</i>	Reed canarygrass	wetlands, riparian
<i>Polygonum cuspidatum</i>	Japanese knotweed	riparian, woodland edge
<i>Rubus discolor</i>	Himalayan blackberry	riparian, woodland edge, lawns
<i>Senecio jacobaea</i>	Tansy ragwort	croplands, lawns, pastures
<i>Silybum marianum</i>	Milk thistle	croplands, lawns, pastures
<i>Ulex europaeus</i>	Gorse	woodland edge, clearcuts

Additional species, although not officially listed as noxious weeds, can be problematic in even small amounts when they contaminate ryegrass harvested for grass seed, due to strict standards for purity of ryegrass sold commercially. Some landowners consider mistletoe (*Phoradendron villosum*), a plant that parasitizes mostly oaks, to be undesirable, so kill it or harvest it for sale. However, several studies have documented the outstanding value of its berries to wintering wildlife, especially bluebirds (this is also true of madrone). Also, some Luckiamute weeds highly damaging to native plant communities have not yet made it onto official agency lists of noxious weeds due to administrative delays or limited information. One example is false-brome, *Brachypodium sylvaticum*, See:

- <http://www.appliedeco.org/FBWG.htm> and
- http://www.nativeseednetwork.org/about/feature_detail.php?id=387

Information on noxious weeds and their control also is available from several sources, such as:

- http://www.nwcb.wa.gov/weed_info/contents.html
- <http://invader.dbs.umt.edu/>
- <http://tncweeds.ucdavis.edu/>
- <http://www.cof.orst.edu/resfor/cameron/demo5.php>

Important Habitats and Communities

Under Oregon’s Natural Heritage Program (ORNHP), the BLM’s Little Sink Research Natural Area (in the upper Luckiamute) contains the Willamette Valley ecoregion’s designated representative of two “ecological cells” (ORNHP 2003): Douglas fir – grand fir / vine maple – salal, and Slump pond at margin of valley with aquatic beds and marsh shore.

The Luckiamute River itself contains the ORNHP's designated representative of two other ecological cells (ORNHP 2003): Pacific willow shrub swamp, and Oregon ash / Pacific willow woodland.

An ORNHP survey of 172 wetland and riparian sites in the Willamette Valley named the lower Luckiamute as one of the 21 highest-quality areas, chiefly from a botanical perspective (Titus *et al.*, 1996)

More broadly, almost any area of the watershed that has experienced minimal soil disturbance and retains predominantly native vegetation could be important for maintaining the watershed's overall plant diversity. In particular, oak woodlands, prairies, stands of mature madrone, springs, seasonal wetlands, riparian areas, and rock ledges are most likely to support plant species not found elsewhere in the watershed. These are discussed in Section 6 of this report.

Conservation and Restoration Potential: Groups or landowners interested in restoring native plant communities can employ a variety of restoration techniques, depending on the habitat that is being restored. Restoration may consist simply of making a long-term commitment to continually remove invasive species. Or, in the case of prairie and oak woodlands, it may involve controlled burns, mowing, restriction of severe grazing, selective thinning, or other techniques. In the valley bottom and foothills, selective removal of Douglas-fir often benefits oaks and other native plants. For oaks, hot underburns decrease the acorn crop the following year, but in subsequent years acorn production is greater than it was during the pre-burn period. Open-grown trees, and trees with wide crowns, are the best acorn producers. Crowded trees collectively may produce well, but there are fewer acorns per tree. For more information on oaks and oak management in Oregon, see: <http://www.wa.gov/wdfw/hab/oakfinal.pdf> and www.fs.fed.us/pnw/olympia/silv/oak and <http://www.dfw.state.or.us/public/WoodlandArc/WhiteOak.pdf> .

Replanting with native species characteristic of the intended habitat (Table 4.4.2.1c, for example) often speeds the restoration process, but only if conditions are suitable. In many cases, seeds of native species remain in the soil long after land has been cleared and cultivated or overrun by weeds, and may germinate decades later, given the proper conditions. Additional information on prairie restoration strategies and seed sources can be found at:

<http://www.appliedeco.org/RestorationConference2003/ConferenceProceedings03.htm>
http://publicworks.co.marion.or.us/parks/SERConf_Min110901.asp
<http://publicworks.co.marion.or.us/environment/restoration/restorationtypes.asp>
<http://www.nativeseednetwork.org/home/index.php>

However, conservation of relatively intact remnants of native plant communities is ultimately a more successful strategy than restoration. Policies that minimize application of herbicides along roadsides and cropland, as well as minimize soil tillage, compaction, and draining, are most likely to benefit watershed health in the long run.

Table 4.4.2.1c. Native plant species that collectively often characterize *upland prairie* or *oak savannah* in the Willamette Valley.

Note: Upland prairie restoration projects strive to favor these species through seeding, burning, mowing, and other practices. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), and Beazell County Park (B) and may include a few records from outside the watershed in Polk/Benton County. For additional species (including mosses) characteristic of this habitat, see: <http://oregonstate.edu/~wilsomar/Index.htm>

Scientific Name	Common Name	Habit	Luckiamute
<i>Achillea millefolium</i>	western yarrow	Perennial	U, B
<i>Achnatherum lemmonii</i> (<i>Stipa lemmonii</i>)	Lemon's needlegrass	Perennial	B
<i>Allium accuminatum</i>	tapertip onion	Perennial	U
<i>Allium amplexans</i>	slimleaf onion	Perennial	U
<i>Amelanchier alnifolia</i>	western serviceberry	Perennial	U
<i>Apocynum androsaemifolium</i>	spreading dogbane	Perennial.	U, B
<i>Asclepias speciosa</i>	showy milkweed	Perennial	U
<i>Aster curtus</i>	Curtus' aster	Perennial	
<i>Aster hallii/chilensis</i> ssp. <i>chilensis</i>	Hall's aster/ Pacific aster	Perennial	U, B
<i>Berberis aquifolium</i>	tall Oregon grape	Perennial	
<i>Brodiaea (Dichelostemma) congesta</i>	field cluster lily	Perennial	U, B
<i>Brodiaea coronaria</i>	harvest brodiaea	Perennial	
<i>Bromus carinatus</i>	California brome	Perennial	U
<i>Bromus sitchensis</i>	Sitka brome	Perennial	U, B
<i>Calochortus tolmiei</i>	cat's ear	Perennial	B
<i>Camassia leichtlinii</i>	tall camas	Perennial	U
<i>Cardamine oligosperma</i>	little western bittercress	Annual	U, B
<i>Carex tumulicola</i>	foothill sedge	Perennial	U, B
<i>Chamomilla suaveolens</i> (<i>Matricaria</i> sp.)	pineapple weed	Annual	
<i>Clarkia amoena</i> var. <i>caurina</i>	fairwell to spring	Annual	B
<i>Clarkia purpurea</i> ssp. <i>quadravulnera</i>	small-flowered godetia	Annual	U
<i>Collinsia grandiflora</i>	large-flowered blue-eyed Mary	Annual	U, B
<i>Collomia grandiflora</i>	large-flowered collomia	Annual	
<i>Comandra umbellata</i>	bastard toad-flax	Perennial	U
<i>Danthonia californica</i>	California oatgrass	Perennial	B
<i>Delphinium leucophaeum</i>	pale larkspur	Perennial	

Table 4.4.2.1c. Native plant species that collectively often characterize *upland prairie* or *oak savannah* in the Willamette Valley.

Note: Upland prairie restoration projects strive to favor these species through seeding, burning, mowing, and other practices. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), and Beazell County Park (B) and may include a few records from outside the watershed in Polk/Benton County. For additional species (including mosses) characteristic of this habitat, see: <http://oregonstate.edu/~wilsomar/Index.htm>

Scientific Name	Common Name	Habit	Luckiamute
<i>Delphinium menziesii</i>	Menzie's larkspur	Perennial	B
<i>Delphinium oreganum</i>	Oregon larkspur	Perennial	U
<i>Delphinium pavonaceum</i>	peacock larkspur	Perennial	U
<i>Dodocatheon hendersonii</i>	broadleaf shooting star	Perennial	U, B
<i>Elymus glaucus</i>	blue wildrye	Perennial	U, B
<i>Elymus trachycaulus</i>	slender wheatgrass	Perennial	
<i>Erigeron decumbens</i> var. <i>decumbens</i>	Willamette Daisy	Perennial	U
<i>Eriophyllum lanatum</i>	wooly sunflower	Perennial	U, B
<i>Erythronium oreganum</i>	giant fawn lily	Perennial	U, B
<i>Festuca californica</i>	California fescue	Perennial	B
<i>Festuca roemerii</i>	Roemer's fescue	Perennial	B
<i>Fragaria virginiana</i>	blue-leaf strawberry	Perennial	B
<i>Fritillaria affinis</i>	chocolate lily	Perennial	
<i>Geranium oreganum</i>	Oregon geranium	Perennial	B
<i>Heterocodon rariflorum</i>	heterocodon	Annual	
<i>Horkelia congesta</i> ssp. <i>congesta</i>	shaggy horkelia	Perennial	U
<i>Iris tenax</i>	Oregon iris	Perennial	U, B
<i>Juncus tenuis</i>	slender rush	Perennial	U
<i>Koeleria cristata</i>	prairie junegrass	Perennial	B
<i>Lathyrus holochlorus</i>	thin-leaved peavine	Perennial	U, B
<i>Linanthus bicolor</i>	bicolored linanthus	Annual	
<i>Lithofragma parviflora</i>	smallflower woodland star	Perennial	
<i>Lomatium bradshawii</i>	Bradshaw's lomatium	Perennial	
<i>Lomatium dissectum</i>	fern-leaved lomatium	Perennial	U
<i>Lomatium nudicaule</i>	barestem desert-parsley	Perennial	U, B
<i>Lomatium triternatum</i>	nine-leaf lomatium	Perennial	

Table 4.4.2.1c. Native plant species that collectively often characterize *upland prairie* or *oak savannah* in the Willamette Valley.

Note: Upland prairie restoration projects strive to favor these species through seeding, burning, mowing, and other practices. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), and Beazell County Park (B) and may include a few records from outside the watershed in Polk/Benton County. For additional species (including mosses) characteristic of this habitat, see: <http://oregonstate.edu/~wilsomar/Index.htm>

Scientific Name	Common Name	Habit	Luckiamute
<i>Lomatium utriculatum</i>	common lomatium	Perennial	U, B
<i>Lotus micranthus</i>	small-flowered deervetch	Annual	B
<i>Lotus purshianus</i>	Spanish-clover	Annual	
<i>Lupinus micranthus</i>	field lupine	Annual	
<i>Lupinus polyphyllus</i>	bigleaf lupine	Perennial	
<i>Lupinus rivularis</i>	stream lupine	Perennial	
<i>Lupinus suphureus</i> ssp. <i>kincaidii</i>	Kincaid's lupine	Perennial	
<i>Madia elegans</i>	showy tarweed	Annual	U
<i>Madia sativa</i>	coast tarweed	Annual	U, B
<i>Montia howellii</i>	Howell's montia		
<i>Navarretia squarrosa</i>	skunkweed	Annual	U
<i>Nemophila menziesii</i>	baby blue-eyes	Annual	U
<i>Nemophila parviflora</i>	small flower nemophila	Annual	B
<i>Plagiobothrys scouleri</i>	Scouler's popcorn-flower	Annual	U
<i>Plectritis congesta</i>	rosy plectritis	Annual	B
<i>Poa scabrella</i>	pine bluegrass	Perennial	
<i>Polystichum munitum</i>	western swordfern	Perennial	U, B
<i>Prunella vulgaris</i> var. <i>lanceolata</i>	self-heal	Perennial	B
<i>Pteridium aquilinum</i>	western bracken fern	Perennial	B
<i>Quercus garryana</i>	Oregon white oak	Perennial	U, B
<i>Quercus kelloggii</i>	California black oak	Perennial	
<i>Ranunculus occidentalis</i>	western buttercup	Perennial	B
<i>Rosa nutkana</i>	Nootka rose	Perennial	B
<i>Salix scouleriana</i>	Scouler willow	Perennial	B
<i>Sanicula bipinnatifida</i>	purple sanicle	Perennial	U, B
<i>Sanquisorba occidentalis</i>	Annual burnet	Ann./bi	

Table 4.4.2.1c. Native plant species that collectively often characterize *upland prairie* or *oak savannah* in the Willamette Valley.

Note: Upland prairie restoration projects strive to favor these species through seeding, burning, mowing, and other practices. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), and Beazell County Park (B) and may include a few records from outside the watershed in Polk/Benton County. For additional species (including mosses) characteristic of this habitat, see: <http://oregonstate.edu/~wilsomar/Index.htm>

Scientific Name	Common Name	Habit	Luckiamute
<i>Sidalcea campestris</i>	meadow sidalcea	Perennial	U
<i>Sidalcea nelsoniana</i>	Nelson's checkermallow	Perennial	
<i>Sidalcea virgata</i>	rose checker-mallow	Perennial	U, B
<i>Silene hookeri</i>	Indian pink	Perennial	U, B
<i>Solidago canadensis</i>	Canada goldenrod	Perennial	U
<i>Trifolium tridentata</i>	tom-cat clover	Annual	
<i>Trifolium variegatum</i>	white-tip clover	Annl/Per	
<i>Vicia americana</i> v. <i>truncata</i>	American vetch	Perennial	
<i>Viola nuttallii</i> var. <i>praemorsa</i>	canary violet	Perennial	
<i>Wyethia angustifolia</i>	narrow-leaf mule's ears	Perennial	U

4.4.2.2 Wetlands

Overview

Wetlands – which include (for example) areas called marshes, swamps, swales, sloughs, and wet prairies – are an important feature of the Luckiamute watershed. Potentially, wetlands are both a concern and a public asset. They are potentially a concern for two reasons. First, landowners proposing some types of activities in wetlands must first receive permits from federal and state agencies, and this can cause project delays unless planned for in advance. Wetland “removal” and “fill” permits are nearly always granted, but landowners routinely are required to “mitigate” the alteration of wetlands by constructing, or preferably restoring, wetlands elsewhere, sometimes at substantial cost. Second, wetlands are a concern because as a resource, the current area of wetlands in the Willamette Valley is but a tiny fraction of what once existed (Table XX). The loss of so much wetland area is a concern because wetlands in their natural condition typically provide a large number of services and thus are a vital public asset. For example, many wetlands can filter and cleanse surface runoff, help reduce flood peaks in rivers, and provide habitat for species found nowhere else. Infrastructure built in wetlands is susceptible both to frequent flooding and to unstable soils, making wetlands a poor, expensive choice for buildings and roads. Crops planted in many former wetlands are subject to frequent or late damage from frost, floods, and wildlife. While the destruction

of a small portion of a single wetland may seem too trivial to worry about, the repeating of such actions among many wetlands over time eventually adds up and imposes significant costs on taxpayers, e.g., for repair of roads, flood damage payments.

Nonetheless, the conversion of many wetlands to agricultural uses, which mostly occurred in the early 1900's, has helped support the economy of the Luckiamute watershed. Recognizing this, state and federal agencies allow farming to continue in former wetlands where farming has long existed. Grazing also is permitted, and in some instances might benefit wetlands by minimizing incursion of Himalayan blackberry. Existing laws allow small fills in wetlands and other waters (less than 50 cubic yards, or less than 1 cubic yard in waters designated by the state as Essential Indigenous Anadromous Salmonid Habitat). For more information on wetland regulations, functions, and values, see:

<http://www.epa.gov/owow/wetlands/> or <http://statelands.dsl.state.or.us/>.

Seasonally dry wetlands sometimes confuse landowners because such areas aren't obviously distinguishable from non-regulated uplands. To confirm or deny wetland status, an agency wetland specialist (or a consultant trained in use of the standardized federal delineation manual) must visit the site at an appropriate time of year and identify all dominant plant species, carefully examine the soil, and look for indicators of flooding or saturation. To many citizens, it may seem illogical to call areas that appear dry 95% of the year a "wetland." However, often it is the driest wetlands that provide the greatest benefits to society. Like a dry sponge, they are the most effective wetlands for soaking up excess runoff and the pollutants that it bears. When many wetlands in a watershed behave in this manner, downstream flood peaks are lower. Seasonally dry wetlands also host many plants seldom found in uplands or in wetlands that remain inundated year-round (Table 4.4.2.2).

While it is true that not all wetlands are equally effective in supporting functions valued by society, it also is not possible to identify the least effective ("marginal") wetlands by merely applying a one-factor criterion, such as "wetness" or "isolation." Seasonally dry wetlands that appear to be unconnected to other surface waters are very important for some functions, but this can be determined only by considering their watershed, soil type, slope, vegetation structure, and many other interacting factors. Procedures are available that account for such multiple factors in assigning ratings to wetlands and their functions, e.g., Gersib (1997), Adamus and Field (2001). For information on these, see:

http://www.oregonstatelands.us/hgm_guidebook.htm

<http://www.ecy.wa.gov/programs/sea/pubs/97-99/97-99.html>

Wetland maps produced by the National Wetland Inventory may be viewed online at: http://www.nwi.fws.gov/mapper_tool.htm. However, many wetlands that are regulated are not shown, and some of the wetlands that are shown are not legally regulated. As a precaution, landowners planning to fill low spots or excavate in areas mapped as floodplain or having any of the soils listed in Table 4.8.1a, should minimize their legal liabilities by first asking wetland specialists to make and document a "wetland determination" at the project site.

Local Wetland Characteristics

Elevation, slope, soil type, land cover, and other characteristics of local wetlands are summarized geographically in Appendices XXXX.

Conservation and Restoration Potential

Several government programs are available to provide technical support and funds to qualifying landowners who wish to restore wetlands to their property. For example, see:

- <http://www.sherm.com/wild/index.html>
- <http://www.nrcs.usda.gov/programs/>
- <http://www.epa.gov/owow/wetlands/facts/funding.pdf>
- <http://publicworks.co.marion.or.us/parks/nhp/partnerships.asp>
- http://www.oregon-plan.org/archives/steelhead_dec1997/st-12.html

For general information about wetland restoration, consult this report that is downloadable from the internet:

http://www.nmfs.noaa.gov/habitat/habitatprotection/pdf/Wet%20Res%20Guidance_FINAL.pdf

The extent of potential opportunities for wetland restoration, based only on soil conditions, is compiled geographically in Table 4.4.2.2.

Table 4.4.2.2. Native plant species that often characterize <i>seasonal wetlands</i> in the Willamette Valley.			
Note: Wetland restoration projects (except those in floodplains) strive to favor these species through seeding, burning, and/or manipulation of flooding depth, duration, frequency, and seasonality. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), Beazell County Park (B), and may include a few records from outside the watershed in Polk/Benton County			
Scientific Name	Common Name	Type	Luckiamute
<i>Agrostis exarata</i>	spike bentgrass	Perennial	U
<i>Allium amplexans</i>	slimleaf onion	Perennial	U
<i>Aristida oligantha</i>	prairie threeawn	Annual	
<i>Aster curtus</i>	Curtus' aster	Perennial	
<i>Aster hallii</i> ssp. <i>chilensis</i>	Hall's aster/ Pacific aster	Perennial	U, B
<i>Barbarea orthoceras</i>	wintercress	Bi/Peren	U
<i>Beckmannia syzigachne</i>	American sloughgrass	Annual	U
<i>Boisduvalia densiflora</i>	dense spike-primrose	Annual	
<i>Brodiaea (Triteleia) hyacinthina</i>	hyacinth brodiaea	Perennial	
<i>Brodiaea coronaria</i>	harvest brodiaea	Perennial	
<i>Calandrinia ciliata</i>	red maids	Annual	
<i>Callitriche heterophylla</i>	water starwort	Annual	U
<i>Camassia leichtlinii</i>	tall camas	Perennial	U
<i>Camassia quamash</i>	common camas	Perennial	U

Table 4.4.2.2. Native plant species that often characterize seasonal wetlands in the Willamette Valley.

Note: Wetland restoration projects (except those in floodplains) strive to favor these species through seeding, burning, and/or manipulation of flooding depth, duration, frequency, and seasonality. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), Beazell County Park (B), and may include a few records from outside the watershed in Polk/Benton County

Scientific Name	Common Name	Type	Luckiamute
<i>Cardamine penduliflora</i>	Willamette Valley bittercress	Perennial	U
<i>Carex aurea</i>	golden sedge	Perennial	
<i>Carex densa</i>	dense sedge	Perennial	U
<i>Carex echinata</i>	muricate sedge	Perennial	
<i>Carex feta</i>	green-sheath sedge	Perennial	
<i>Carex pachystachya</i>	thick-headed sedge	Perennial	U
<i>Carex unilateralis</i>	one-sided sedge	Perennial	U
<i>Centaurium muhlenbergii</i>	monterey centaury	Annual	
<i>Centunculus minimus</i>	chaffweed	Annual	U
<i>Danthonia californica</i>	California oatgrass	Perennial	B
<i>Deschampsia cespitosa</i>	tufted hairgrass	Perennial	U
<i>Deschampsia danthonioides</i>	Annual hairgrass	Annual	
<i>Deschampsia elongata</i>	slender hairgrass	Perennial	U
<i>Dodocatheon hendersonii</i>	broadleaf shooting star	Perennial	U
<i>Downingia elegans</i>	showy downingia	Annual	U
<i>Downingia yina</i>	Willamette downingia	Annual	
<i>Eleocharis acicularis</i>	needle spike-rush	Annual	
<i>Eleocharis ovata</i>	common spike-rush	Annual	
<i>Epilobium ciliatum</i> var. <i>watsonii</i>	hairy willow-herb	Perennial	
<i>Epilobium paniculatum</i>	autumn willow-herb	Annual	
<i>Erigeron decumbens</i> var. <i>decumbens</i>	Willamette Daisy	Perennial	U
<i>Eriophyllum lanatum</i>	wooly sunflower	Perennial	U, B
<i>Eryngium petiolatum</i>	coyote thistle	Perennial	
<i>Gnaphalium palustre</i>	lowland cudweed	Annual	U
<i>Gnaphalium purpureum</i>	purple cudweed	Ann/bi	
<i>Gratiola ebracteata</i>	bractless hedge-hyssop	Annual	
<i>Grindelia intergrifolia</i>	Willamette Valley gumweed	Perennial	U
<i>Haplopappus racemosus</i>	racemed goldenweed	Perennial	
<i>Hordeum brachyantherum</i>	meadow barley	Perennial	
<i>Horkelia congesta</i>	shaggy horkelia	Perennial	

Table 4.4.2.2. Native plant species that often characterize seasonal wetlands in the Willamette Valley.

Note: Wetland restoration projects (except those in floodplains) strive to favor these species through seeding, burning, and/or manipulation of flooding depth, duration, frequency, and seasonality. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), Beazell County Park (B), and may include a few records from outside the watershed in Polk/Benton County

Scientific Name	Common Name	Type	Luckiamute
<i>Isoetes nutalli</i>	Nuttall's quillwort	Perennial	
<i>Juncus bolanderi</i>	Bolander's rush	Perennial	
<i>Juncus bufonius</i>	toad rush	Annual	U
<i>Juncus nevadensis</i>	Nevada rush	Perennial	
<i>Juncus tenuis</i>	slender rush	Perennial	U
<i>Lasthenia glaberrima</i>	smooth lasthenia	Annual	
<i>Lindernia anagallidea</i>	false-pimpernel	Annual	
<i>Lomatium nudicaule</i>	barestem desert-parsley	Perennial	
<i>Lotus formosissimus</i>	seaside lotus	Perennial	
<i>Lotus pinnatus</i>	meadow deervetch	Perennial	U
<i>Lotus purshianus</i>	Spanish-clover	Annual	
<i>Lupinus polyphyllus</i>	bigleaf lupine	Perennial	
<i>Luzula campestris</i>	field woodrush	Perennial	
<i>Madia glomerata</i>	cluster tarweed	Annual	U
<i>Microseris laciniata</i>	cut-leaved microseris	Perennial	
<i>Microsteris gracilis</i>	pink microsteris	Annual	
<i>Mimulus guttatus</i>	common monkey-flower	Ann/per.	
<i>Montia fontana</i>	water chickweed	Annual	U
<i>Montia linearis</i>	narrow-leaved montia	Annual	U
<i>Myosurus minimus</i>	least mouse-tail	Annual	
<i>Navarretia intertexta</i>	needle-leaved navarretia	Annual	U
<i>Navarretia squarrosa</i>	skunkweed	Annual	U
<i>Orthocarpus bracteosus</i>	rosy owl-clover	Annual	U
<i>Orthocarpus hispidus</i>	hairy owl-clover	Annual	
<i>Panicum capillare</i>	common witchgrass	Annual	U
<i>Panicum occidentale</i>	western witchgrass	Perennial	
<i>Perederidia oregana</i>	Oregon yampah	Perennial	
<i>Perideridia gairdneri</i>	yampah or false-carraway	Perennial	
<i>Plagiobothrys figuratus</i>	fragrant popcorn-flower	Annual	U
<i>Plagiobothrys scouleri</i>	Scouler's popcorn-flower	Annual	U

Table 4.4.2.2. Native plant species that often characterize seasonal wetlands in the Willamette Valley.

Note: Wetland restoration projects (except those in floodplains) strive to favor these species through seeding, burning, and/or manipulation of flooding depth, duration, frequency, and seasonality. List provided by Kathy Pendergrass, US Fish & Wildlife Service. Luckiamute records are from ORNHIC (OH), OSU Herbarium (U), Beazell County Park (B), and may include a few records from outside the watershed in Polk/Benton County

Scientific Name	Common Name	Type	Luckiamute
<i>Poa scabrella</i>	pine bluegrass	Perennial	
<i>Polygonum douglasii</i>	douglas knotweed	Annual	
<i>Potentilla gracilis</i>	slender cinquefoil	Perennial	U
<i>Prunella vulgaris</i> var. <i>lanceolata</i>	self-heal	Perennial	U, B
<i>Psilocarphus elatior</i>	tall wooly-heads	Annual	U
<i>Ranunculus flammula</i>	creeping buttercup	Perennial	
<i>Ranunculus occidentalis</i>	western buttercup	Perennial	
<i>Ranunculus orthorhynchus</i>	straight beaked buttercup	Perennial	U
<i>Rorippa curvisiliqua</i>	western yellowcress	Ann./bi.	U
<i>Rosa nutkana</i>	Nootka rose	Perennial	
<i>Sanquisorba occidentalis</i>	Annual burnet	Ann./bi	
<i>Saxifraga oregana</i>	bog saxifrage	Perennial	B
<i>Sidalcea cusickii</i>	Cusick's checker-mallow	Perennial	
<i>Sisyrinchium idahoense</i> (<i>angustifolium</i>)	blue-eyed grass	Perennial	B
<i>Sisyrinchium hitchcockii</i>	Hitchcock's blue-eyed grass	Perennial	
<i>Spiranthes romanzoffiana</i>	ladies-tresses	Perennial	
<i>Trichostema lanceolatum</i>	vinegar weed	Annual	
<i>Veronica peregrina</i>	purslane speedwell	Annual	
<i>Viola adunca</i>	early blue violet	Perennial	
<i>Wyethia angustifolia</i>	narrow-leaf mule's ears	Perennial	U
<i>Zigadenus venenosus</i>	death camas	Perennial	B

4.4.2.3 Riparian Areas

Riparian zones link terrestrial and aquatic ecosystems. Riparian areas act as sediment filters and areas where organic material and nutrients are transformed before entering the stream; they also provide shade and detritus (dead organic material) directly to the stream ecosystem. Some unconstrained (or floodplain) riparian areas provide important backwater forage areas and refugia from high water flows for juvenile salmonids.

Although riparian areas are frequently disturbed naturally, historic and current land use decisions have dramatically altered the magnitude and frequency of this disturbance.

The historic disturbance regime of the Luckiamute/ Ash Creek watersheds has influenced the current condition of riparian areas. On the valley floor stream channel paths constantly changed, side channels were added to and cut off from the river system, mainly in response to high flow events. In the forested upper reaches of the watershed, riparian vegetation composition was most likely influenced by other disturbances in addition to high water flow, *i.e.*, debris flows, wind throws, fire, etc.

Information on the presettlement vegetation is limited (see Section 4.4.1.1). Again, disturbance probably played an important role in structuring the composition of the plant communities. Plant communities would have changed in response to the type and frequencies of these disturbances. No studies have quantified the relative proportion of hardwoods to conifers along Coast Range streams prior to European settlement. One study of vegetation composition around upland streams concluded that riparian trees were mostly hardwoods, while another study concluded that there were many patches of old growth conifers around streams (Licata *et al.*, 1998).

Most likely, both types of vegetation, conifers and broad-leaved trees, were present along streams in the study area. An interesting consequence of the shift from conifers to alders comes from a recent study. As previously mentioned, the composition of stream side vegetation is known to affect water quality by filtering sediments and coarse organic material from runoff, and by slowing water infiltration so that important chemical transformations can occur (*e.g.*, denitrification and conversion of phosphorus containing compounds; both nitrogen and phosphorus are important nutrients for aquatic life). A recent study in the Salmon River (OR) watershed has linked land cover, in this case alder forest cover, with instream nitrate and dissolved organic N concentrations. The implications of this work indicate that human watershed modifications which increase the proportion of alder in the watershed can directly affect the availability of nitrogen and phosphorus in coastal stream networks (Compton *et al.*, In Press).

In addition to the composition of the riparian plant community, disturbance would most certainly have affected the size of the riparian vegetation. Undoubtedly, riparian areas within the Luckiamute/ Ash Creek study area were a shifting mosaic of different species, ages and sizes of trees. Licata (1998) concluded that there were probably more old trees along the streams historically than there are today, providing more large woody debris to the streams.

4.4.3 LAND USE

An understanding of land use is vital to understanding the function and character of a watershed because land use is linked to many of the enhancements and degradations of a watershed. One land use type, for example agriculture, can produce both watershed enhancement by supporting the economy of the area, producing food, and creating jobs,

and produce degradation through means such as non-point source pollution from fertilizer and pesticides. Although no single land use produces only enhancements or only degradations, it is necessary to thoroughly investigate the land uses within the study area to document what the enhancements and degradations are. Land ownership is also important as it determines some of the land uses in the watershed and influences where monitoring and restoration can take place. Land Cover/ Land Use are summarized for the Luckiamute/ Ash Creek study area in Table 4.4.1.2a.

4.4.4 HISTORIC LAND USE CONDITIONS

Some of the earliest land uses in this region were the construction of homesteads, and the clearing of lands for agricultural and pasture for grazing. The earliest European-American settlers to the valley cleared much of the land for planting and grazing. Pete Frantz, a Luckiamute Valley resident whose grandparents arrived in the Luckiamute Valley in 1866 gives the following account:

“At that time people thought that there would always be timber so they just ignored it or tried to get rid of it. In fact, on our land there were two places where the folks just burnt down the second growth timber...they had to have room to grow grain to feed their stock. There wasn't a demand for second growth timber anyway (Frantz and Brandon, 1976). “

In the early 1900s, timber became a profitable commodity, and logging began to increase in importance in the Luckiamute Valley. Logging was accomplished with horses and steam-powered donkey engines. Small mills were constructed throughout the valley, but most of the logs were sent to Salem for milling. Grains were also historically a major agricultural commodity, and Italian prunes and hops were introduced in 1890. These specialty crops became so valuable that at one point, there were over 4,600 acres of hops planted near Independence (Newton, 1971), however crops of both Italian prunes and hops declined by World War II (Mattson and Gallagher, 2001).

One of the most notable historic land uses within the study area is the army cantonment at Camp Adair during World War II (see Section 4.4.5.5 for more information on Camp Adair). Concerned by the potential contamination of surface water by lead, we were asked by the LWC to report on the use of Camp Adair as a firing range by local area police departments. According to the Benton Co. sheriff's office the range is owned by the National Guard (HQ in Salem) and the range is not currently being used by Benton County.

4.4.5 CURRENT LAND USE CONDITIONS

4.4.5.1 Agriculture

Today, agricultural use within the study area includes production of grass seed, wheat, hay and oats; orchards and vineyards; forest products; and the raising of both cattle and sheep, with the major commodities being grass seed, wheat, hay, and oats and minor commodities including clover, sweet corn, mint, alfalfa, and filberts (Taylor *et al.*, 2003). Other commodities grown in the area include nursery products and Christmas trees (Mattson and Gallagher, 2001). Many orchards exist in the northeastern portion of the watershed just west and north of the city of Monmouth, in the hills above Falls City, and north of the confluence of the Luckiamute and Little Luckiamute Rivers along Elkins Road. There are several vineyards in and around the study area including: Airlie Winery, Chateau Bianca, Eola Hills Wine Cellars, Flynn Vineyards, Schwarzenberg Vineyards, Oak Grove Orchards Winery, Serendipity Cellars, and Van Duzer Vineyards.

Private timber companies own a substantial amount of land in the watershed (about 26%). The major timber companies include Boise and Weyerhouser. Other timber companies include Starker Forests, Inc., Stimson Lumber Co. and B & D Timber Inc (Table 4.4.6a a & b).

Cattle have always been an important part of valley life. Early settlers cleared land so that their livestock could graze. Today, cattle and sheep graze many areas in the Luckiamute Watershed, including Oregon State University's Soap Creek Ranch and Berry Creek Ranch. Approximately 65% of these ranches are open grasslands utilized primarily for forage production. The rest of the ranch land is forested (Oregon State University Animal Sciences Department, 2003). Each ranch has facilities used for research, student projects, classes, and general maintenance of the beef herds. Facilities include scales, covered processing areas, and hay barns. There are currently 225 cows and 4 bulls on the Soap Creek Ranch. One hundred and fifty-five (155) of these are bred to calve spring, and 70 are bred to calve in the fall. The cattle move between the Soap Creek Ranch location and the Berry Creek location, located along Tampico Road, adjacent to Dunn Forest (Sexson, 2003).

The Soap Creek Ranch has expressed interest (by submitting a proposal to OWEB) in constructing fences around the stream (Sexson, 2003). In addition, a team of OSU Engineering students has created a water pumping system on the Soap Creek Ranch site to provide water to cattle. There are currently two pumps located on the ranch site, with a potential to create more if these are successful. Both fences and the watering system would prevent livestock from entering the streams.

4.4.5.2 Mills

The LWC asked team to identify mills associated with some of the early Land Claims. Table 4.4.5.3 lists Donation Land Claims for the Luckiamute/Ash Creek study area.

Table 4.4.5.3. Donation Land Claim settlers and mill sites in the Luckiamute Valley. Listed are the owner, year of claim, township and range, quarter section and notes. Sources of information are listed below.

Owner	Year	Township	Range	Section	Quarter/ Other
Sawmills Identified within the OSU Research Forest (Wisner, 1992)					
Coote/Cornutt	1935-1937	T10S	R5W	9	north half, middle
Coote/Cornutt	1929-1935	T10S	R5W	16	north half, middle
Coote/Weinert	1936-1939	T10S	R5W	15	SE 1/4
Cooper's Mill	1930s	T10S	R5W	8	SW 1/4
Bennett Brothers	1930s	T10S	R5W	16	SW 1/4
Oak Creek Mill	1910-1920	T11S	R5W	7/18	
Oak Creek Mill	1910-1937	T11S	R5W	18	SW 1/4
Oak Creek Mill	1910-1937	T11S	R5W	17	NW 1/4
OSU Mill	1947-1955	T11S	R5W	20	NW 1/4
Govier Mill	1931-1939	T10S	R5W	10	SE 1/4
Soap Creek Mill	1890	T10S	R5W	35	SW 1/4
Valley Mill	1935-1955	T11S	R5W	3	NE 1/4
Calloway Creek Mill	1911-1916	T10S	R5W	36	E middle
Mt. View Mill	1934-1937	T11S	R5W	2	NE 1/4
Zager Mill	1937	T11S	R6W	24	E middle
Zeller Mill	1937	T11S	R6W	12	E middle
Unnamed	1937	T11S	R5W	10	W middle
Sulphur Springs Mill	1890	T11S	R5W	5	NW 1/4
Sharp's Mills		T8W	R4W	20	
Scott's Mill		T8W	R4W	28	
Sawmills identified near Falls City (fall city draft document received from C. Vandenberg)					
John Thorp	1853	T9W	R4W	11	
Shrader-Mowery					on teal creek, near the Falls City water reservoir
Rowell					9 miles west of Dallas,
Palmehn					above Falls City, above Dutch Creek
Sawmills near Kings Valley (Theurer, 2003)					
Barnhart-Kochis					upper end of the Luckiamute
Henry Baumann	1920s				up Luckiamute canyon
Earl Godsey	early 1930s				Luckiamute river, in the "flat field along the river". Powered by steam tractor and did not have a mill pond
Alvin Jones	1916				"in this area"

Moody brothers	1930s			Benton County, below Barnhart's mill, no burner or pond
Frantz family - Big mill at Hoskins	before 1910			on the old Fort Hoskins site. Mill put on Luckiamute and run with water power.
Bayless Moser and Bill Coote	early 1920s			Between Hoskins and Kings Valley, beside railroad
Archie and Dorval Bevins	1923			on the Walter Cosgrove farm, west of Kings Valley
Charles Moser	1931			above Hoskins, on Burgett Creek,
Simpsons	Unknown			Maxfield Creek Road
Christenson	1906			up the "canyon"

4.4.5.3 Mining

Locations of mines, gravel pits and quarries are important watershed features because they can affect water quality and wildlife. We examined available USGS topographical maps for quarries and mines. We found fifty-five quarries or gravel pits located in the study area (Table 5.4.5.4). We also searched the Oregon Department of Geology and Mining Industries (www.oregongeology.com) web page for information on mines in the study area. Unfortunately, we did not find anything for the Luckiamute/ Ash Creek study area.

Table 4.4.5.3. List of Gravel Pits and Quarries by 7th field watershed. Source was 7.5' USGS topographical map.

7th Field HUC	Sub-Basin Name	Feature	Number
17090003060101	Upper Luckiamute	Pit	3
17090003060102	Miller Creek	Pit	1
17090003060201	Wolf Creek	Pit	1
		Quarry	2
17090003060203	Hoskins	Quarry	1
17090003060302	Woods Creek	Quarry	1
17090003060303	Price Creek	Quarry	1
17090003060304	Maxfield Creek	Pit	1
		Quarry	1
17090003060305	Bump Creek	Quarry	1
17090003060401		Quarry	2
17090003060402	Upper Pedee Creek	Pit	2
		Quarry	2
17090003060502	McTimmonds	Quarry	1
17090003060504	Jont Creek	Quarry	1
17090003060601	Upper Little Luckiamute	Pit	4

Table 4.4.5.3. List of Gravel Pits and Quarries by 7th field watershed. Source was 7.5' USGS topographical map.

7th Field HUC	Sub-Basin Name	Feature	Number
17090003060602	Cold Springs	Pit	5
17090003060603	Black Rock Creek	Pit	2
		Quarry	1
17090003060702	Falls City	S	1
17090003060703	Waymire Creek	Quarry	1
17090003060704	Bridgeport	Quarry	4
17090003060801	Upper Teal Creek	Pit	2
17090003060802	Lower Teal Creek	Pit	1
17090003061101	Upper Soap Creek	Quarry	7
17090003061201	Upper Berry Creek	Quarry	2
17090007020503	Harman Slough	Pit	3
17090007020601	Upper North Fork Ash Creek	Quarry	1
		Total	55

4.4.5.4 Landfills

The Coffin Butte Landfill is located near Adair Village just west of Oregon state highway 99. It is situated at the head of an unnamed tributary to the Luckiamute River, between Poison Oak Hill and Coffin Butte. Valley Landfills, Inc., of Corvallis, operates the landfill, and the land now occupied by the landfill was previously used as part of the Camp Adair Army Training Facility. The Coffin Butte Landfill is the second largest landfill in Oregon and is classified by the EPA as a Subtitle D refuse disposal facility. The Valley Landfills, Inc. property covers 700 acres, about 116 acres of which are being used as a landfill at this time. The site has nine cells which will eventually be filled. Cell 1 has been completely filled and cell 2 will be completely filled by the end of 2003. The Coffin Butte landfill receives approximately 1,800 to 2,000 tons of municipal solid waste a day during the spring, and as much as 2,500 to 3,000 tons a day in the summer. Valley Landfills, Inc. plans on increasing tonnage received every year, as demand dictates. The site is equipped with a leachate treatment system, and a methane-base electrical generator (Benson, 2003). There are a water quality monitoring wells located around the facility (see Section 6.6)

4.4.5.5 Military Facilities

Camp Adair, names in honor of Henry Rodney Adair (a West Point graduate and descendant of Oregon pioneers who was killed in 1916) is located 10 miles north of Corvallis. Camp Adair was a military training facility that operated in the southern portion of the Luckiamute watershed between 1941 and 1946. Camp Adair, a WWII army cantonment, occupies 50,000 acres in Benton and Polk Counties. The camp itself occupied only a small portion of that land and covered an area 2 miles wide and 6 miles long, along Oregon state highway 99W near the Benton County line. It was once used as

a training site for army infantry, artillery and engineering units and associated support personnel. The camp had over 1,800 buildings, consisting of barracks, mess halls, offices, churches, five movie theaters, stores, a post office, a bank, and a hospital. Although Camp Adair never reached its full complement of men and women, it quickly became the second largest city in Oregon. Interestingly, full-scale models of European towns were constructed in this area for training troops. During WWII casualties from the Pacific Theater were brought to Camp Adair for treatment and recuperation; the hospital facility could care for 3,600 individuals. Once the troops left Camp Adair, it served as a prisoner-of-war (POW) camp for Italians, then Germans from 1944 through 1946.

Today, the site is owned by state and local governments and a few individuals; only a few buildings and foundations remain of the WWII camp. The former army camp now hosts the E. E. Wilson State Preserve and is home to upland game birds, waterfowl, bald eagles, deer and other species. There are trails, stocked ponds and hunting (during fall and winter) for visitors. Bullets are still being found in trees in the area that is now the McDonald forest (Rogers, 2003). Today, the area that was Camp Adair has many land uses, including Coffin Butte Landfill (see below), the McDonald State Forest, and E.E. Wilson Wildlife area (see below) For more information visit <http://home.teleport.com/~eewilson/campadair.html> or <http://www.ohwy.com/or/e/eewilswa.htm>

4.4.5.6 Wildlife Preserves

The E.E. Wilson Wildlife Area (formerly Camp Adair) is managed by Oregon Fish and Wildlife and Oregon Wetlands. The area supports a diversity of habitats for a variety of sensitive species such as the Sharp-tailed snakes, Red-Legged Frogs, and the Western Pond Turtles. Trails are well established at E.E. Wilson. See Section 4.4.2.1 for information on ORNHP ecological cells.

4.4.6 LAND OWNERSHIP

Land ownership affects the condition of the landscape indirectly because of the various uses for which different owners have used the land. Past land uses have set the stage for the current condition of the natural resources of the Luckiamute/ Ash Creek study area (See earlier Sections on Historical Land Cover/ Land Use). Land ownership also directly affects the current management practices and restoration potential for individual land parcels. Consequently, patterns in land ownership become very important when developing watershed management and restoration plans.

Tables 4.4.6a & b show the current ownership patterns for the study area. In the Luckiamute/ Ash Creek study area, most ownership is private; therefore, most monitoring and restoration is likely to occur on private lands. We recommend that a more detailed ownership assessment be performed (*i.e.*, identify public and willing private land owners) to develop a monitoring and restoration plan. The ownership data set is a combination of the Polk and Benton County tax lot GIS data. After the two data sets were merged in a GIS, the ownership information from the tax lot data was reclassified into public or private ownership.

Ownership	Area (acres)	% Total
Private	170305.6	84.7%
Public, County	1141.0	0.6%
Public, Federal	8339.8	4.1%
Public, Municipal	51.8	0.0%
Public, State	7314.1	3.6%
Unknown	13958.9	6.9%
Total	201111.2	100.0%

Ownership	Area (acres)	% Total
Private	31510.6	93.1%
Public, County	23.5	0.1%
Public, Federal	59.7	0.2%
Public, Municipal	276.3	0.8%
Public, State	255.4	0.8%
Unknown	1714.4	5.1%
Total	33839.9	100.0%

4.5 ZONING AND REGULATIONS

By 1975, the State of Oregon had adopted 19 statewide land use planning goals covering topics from housing to natural resource use, which are achieved through local comprehensive plans. State law requires each city and county to adopt a comprehensive plan that meets the 19 goals, and develop the zoning and land-division ordinances needed to put that plan into effect. These goals are intended to promote consistency in statewide land use and coordination between various local governments. Goal 5, which was amended in 1996, governs natural resources, scenic and historic areas, and open spaces.

**Goal 5: Natural Resources, Scenic and Historic Areas, and Open Spaces
(Oregon Department of Land Conservation and Development, 2003)**

- Plan development to conserve open spaces
- Plan to conserve natural resources: renewable and non-renewable
- Consider efficient consumption of energy when using natural resources
- Protect fish and wildlife areas, in accordance with Oregon Wildlife Commission's fish and wildlife management plans
- Protect stream flow and water levels at an adequate level for fish
- Inventory historically and ecologically significant natural areas

- Investigate building in cluster developments
- State and federal agencies should develop plans for natural resources, historic areas, and open spaces that coordinate with local and regional plans

Goal 5 requires state and local governments coordinate plans for rivers, trails and natural resources, such as wetlands, riparian corridors, wildlife habitat, federal wild and scenic areas. For more information see <http://www.lcd.state.or.us/goalpdfs/goal05.pdf>

The Oregon Legislature passed Senate Bill 1010 to improve agricultural practices near streams. Senate Bill 1010, or the [Agricultural Water Quality Management Act](#) was passed in 1993 at the request of many agricultural interests so that Oregon agriculture could regulate itself as much as possible. Senate Bill 1010 directs the [Oregon Department of Agriculture \(ODA\)](#) to develop an Agricultural Water Quality Management Plan and Rules for watershed in Oregon where there are water quality problems. The ODA along with other agencies identifies priority watersheds for development of Agricultural Water Quality Management Plans. The Luckiamute / Ash Creek study area falls within the Middle Willamette Agricultural Water Quality Management Area. Under this plan local operators will be asked to deal with identified problems such as soil erosion, crop nutrient loss from fields, or degraded streamside areas. Farmers are allowed to choose their own ways of meeting established water quality goals; however, if problems are identified, those who are asked to deal with a problem but continually refuse to do so could be assessed a civil penalty. For more information see <http://www.peak.org/~bentoncd/SB1010.html>.

4.6 ECOREGIONS AND PHYSIOGRAPHIC PROVINCES

There are many ways to group areas into regions. Ecoregions and physiographic provinces are two such methods. Ecoregions are areas that share similar soils, vegetation, and climatic characteristics. The concept of the ecoregion was introduced in 1987 as a water quality management tool that grouped areas based on perceived patterns in climate, soils, potential vegetation, land form and land use (Omernik, 1987). Since the concept was introduced, there have been many different ecoregion classifications schemes produced at many different scales, so it is necessary to know what ecoregion definition is being used to properly understand the designations.

The study area is at the confluence of five ecoregions as defined by the Level IV EPA Ecoregion dataset available from the EPA: Volcanics, Mid-Coast Sedimentary, Valley Foothills, Willamette River Tributaries Gallery Forest, and Prairie Terraces. Descriptions of each of the ecoregions in the study area are shown in Table 4.6.

Table 4.6. Descriptions of the Level IV EPA Ecoregions within the study area (Woods <i>et al.</i> , 2000)	
Level IV EPA Ecoregion	Description

Table 4.6. Descriptions of the Level IV EPA Ecoregions within the study area (Woods *et al.*, 2000)

Level IV EPA Ecoregion	Description
Volcanics	The Volcanics ecoregion varies in elevation from 1,000 to 4,000 feet and is disjunct. Columnar and pillow basalt outcrops occur. Its mountains may have been offshore seamounts engulfed by continental sediments about 200 million years ago. The basaltic substrate preserves relatively stable summer stream flows that still support spring chinook salmon and summer steelhead. Its forests are intensively managed.
Mid-Coast Sedimentary	The Mid-Coast Sedimentary ecoregion is commonly underlain by massive beds of siltstone and sandstone. Its dissected, forested mountains are rugged and are prone to mass movement when the vegetation cover is removed. Stream gradients and fluvial erosion rates can be high.
Valley Foothills	The Valley Foothills ecoregion is a transitional zone between the Willamette Valley, the Cascade Range, and the Coast Range. It has less rainfall than adjacent, more mountainous ecoregions and, consequently, its potential natural vegetation is distinct. Oregon white oak and Douglas-fir were originally dominant but, today, rural residential development, woodland, pastureland, vineyards, tree farms, and orchards are common.
Willamette Valley Tributaries Gallery Forest	In the Willamette River and Tributaries Gallery Forest ecoregion, meandering, low-gradient channels and oxbow lakes are incised into broad floodplains. Deciduous riparian forests that once grew on its fertile, alluvial soils have been largely replaced by agriculture and rural residential, suburban, and urban development.
Prairie Terraces	The undulating Prairie Terraces ecoregion is dissected by low-gradient, meandering streams and rivers. Its fluvial terraces once supported prairie and oak woodlands which were maintained by burning; Oregon ash and fir occurred in wetter areas. Today, grass seed and grain crops are commonly grown.

In addition to ecoregions, physiographic provinces have been used to develop regional descriptions. For example, in their description of the natural vegetation of Oregon and Washington, Franklin and Dyrness (1988) divide the two state area into 15 physiographic provinces. Physiographic provinces are defined natural features of the earth including land formation, climate, and distribution of flora and fauna. The Luckiamute and Ash Creek watersheds fall into the ‘Coast Ranges’ and ‘Willamette Valley’ physiographic provinces [see Chapter II in (Franklin and Dyrness, 1988) for complete descriptions of the

characteristics of each of these physiographic regions]. Two major vegetation zones cover the study area, the *Tsuga heterophylla* (Western Hemlock) and Willamette Valley zones.

4.7 CLIMATE

Climate is one of the factors that determine how watersheds look and behave. For example, climate ultimately determines the type of vegetation (or potential vegetation) that is found in a region. Climate also determines how watersheds function, e.g., the interplay of geology, soils, vegetation, and patterns in rainfall influence sediment and material transport in streams.

4.7.1 TEMPERATURE

Monthly average, and mean high and low temperatures for each month were obtained from observations measured in Dallas, Oregon, from 1971 to 2000 (Oregon Climate Service, 2003). The annual mean temperature of the area is 51.9°F. Mean summer temperatures range from about 55°F in May to 66°F in July and August and mean winter temperatures range from about 39°F in December to 42°F in February (Table 4.7.1). The lowest mean monthly temperature, 33.1 °F, was in January, and the highest mean monthly temperature, 81.5°F, was in August.

Table 4.7.1. Mean temperature (°F) from the years 1971 to 2000 in Dallas, OR. (Oregon Climate Service, 2003)

Month	Mean Maximum	Mean Minimum	Mean
January	45.3	33.1	39.2
February	49.7	34.8	42.3
March	55.2	36.9	46.1
April	60.4	39.4	49.9
May	66.9	43.7	55.3
June	73.0	47.8	60.4
July	80.9	50.4	65.7
August	81.5	49.8	65.7
September	76.7	47.0	61.9
October	64.7	41.2	53.0
November	50.7	37.2	44.0
December	44.2	33.2	38.7
Annual	62.4	41.2	51.9

4.7.2 PRECIPITATION

Precipitation patterns vary from year to year, from season to season and by elevation across the study area. During the period of record (Dallas, OR) from 1935 to 2002 the annual precipitation ranged from 23.0 inches in 2000 to 69.6 inches in 1937 (Oregon Climate Service, 2003). Rainfall in the area is highly seasonal, with 75% of the annual

rainfall occurring between October and March (Table 4.7.2). Most precipitation is delivered in the form of rainfall, such that the rivers are fed primarily by rainfall rather than snowmelt (Clark, 1999). Streamflows are therefore typically highest during the winter months.

Month	Mean
January	7.8
February	6.7
March	5.3
April	3.2
May	2.2
June	1.4
July	0.5
August	0.7
September	1.4
October	3.3
November	7.8
December	8.8
Annual	49.1

Precipitation is also dramatically affected by elevation. Along the northwestern boundary of the watershed in the Coast Range Mountains the annual precipitation is about 141 in. In contrast, in the lowlands of the Willamette Valley the annual precipitation is about 45 inches (Taylor *et al.*, 2003).

4.8 SOILS

The distribution of different soil types can affect land cover/ land use, and hydrology. Hydric soils are formed around lakes and ponds, wetlands, and streams. Erodible soils are a component of the ecological processes that deliver woody debris to the streams that create salmonid habitat. Ultimately, soils affect where people choose to live, as well, as the variety and condition of wildlife within the watershed.

4.8.1 HYDRIC SOILS

Hydric soils, by definition, are soils that formed under conditions of saturation, flooding, or ponding for long enough during the growing season to develop anaerobic conditions. We created a hydric soils layer by merging the SSURGO soils data set with a table of hydric soils (compiled by P. Adamus) listing hydric soils in the Willamette Valley (Table 4.8.1a). Unaltered hydric soils generally support the growth of hydrophytic (wetland) vegetation, and presence of hydric soils is used to assist identification of wetland areas. By acreage, 77% of the watershed's wetlands occur on soils that usually are hydric, and another 18%

occur on soils that sometimes are hydric (*i.e.*, only a portion of the mapped units are hydric). Wetlands that persist on soils mapped as non-hydric (5% of the wetland area) often do so because of artificial impoundments. Conversely, the presence of hydric soils alone does not necessarily indicate the presence of a ‘jurisdictional’ wetland. For example, for the portion of the watershed having both wetland and soils maps available, 24% of the area that is mapped as hydric soil does not currently contain wetlands, probably because of the drainage that has occurred, or because some soils designated as hydric “may have phases that are not hydric depending on water table, flooding, and ponding characteristics” (USDA Natural Resources Conservation Service, 2000).

Many hydric soil areas that currently lack wetlands are prime areas for restoration (Table 5.8.1b). Restoration projects located on Holocene-period alluvial deposits are perhaps the most likely to succeed (D’Amore et al. 2000). Most hydric soils in the Willamette Valley were formed in the Late Pleistocene when glacial floods deposited sediment on older geomorphic soils, resulting in vertically and horizontally abrupt textural differences. Moreover, deposition of lake sediments during the Late Pleistocene (when much of the Willamette Valley was a lake) disrupted gradients along streams well into the Holocene period, as they entered from adjoining foothills, thus causing further deposition of fine alluvial sediments in fans along the valley margin (Balster and Parsons 1968, Reckendorf 1993). The rather impermeable clay layer that was deposited commonly holds rainwater in a “perched” position near the ground surface, giving rise to the chemically-reducing conditions that characterize hydric soils. During dry periods, deeper soils slowly and independently discharge groundwater upward, keeping wet the hydric soils that are closer to the surface (D’Amore et al. 2000).

4.8.1a. Soils types most commonly associated with wetlands in the Luckiamute-Rickreall watershed, based on overlay of NWI map of mid-1980’s wetlands with county soil maps

Soil map unit name	Hydric?	Acres in Wetlands
Waldo silty clay loam	Yes	1012.37
Water	Yes	650.15
Wapato silty clay loam	Yes	319.94
Dayton silt loam	Yes	208.46
Cove silty clay loam	Yes	198.93
Chehalis silty clay loam, occasionally flooded	Some	124.86
Riverwash	Yes	77.54
Xerofluvents, loamy	Some	71.99
Amity silt loam	Yes	68.74
Bashaw silty clay loam	Yes	66.92
Malabon silty clay loam, occasionally flooded	Some	65.95
Cove silty clay loam, thick surface	Yes	62.73
Concord silt loam	Yes	61.52
Mcbee silty clay loam	Some	61.49

4.8.1a. Soils types most commonly associated with wetlands in the Luckiamute-Rickreall watershed, based on overlay of NWI map of mid-1980's wetlands with county soil maps

Soil map unit name	Hydric?	Acres in Wetlands
Coburg silty clay loam, occasionally flooded	Some	60.00
Woodburn silt loam, 0 to 3 percent slopes	Some	50.99
Cloquato silt loam	Some	49.57
Coburg silty clay loam	Some	23.87
Bashaw clay, 0 to 3 percent slopes	Yes	21.67
Dupee silt loam, 3 to 12 percent slopes	No	21.51
Newberg fine sandy loam	Some	21.11
Xerochrepts and haploxerolls, steep	No	13.66
Witham silty clay loam, 2 to 7 percent slopes	No	13.22
Newberg loam	Some	11.30
Pits, quarries	No	10.38
Willamette silt loam, 0 to 3 percent slopes	Some	9.67
Woodburn silt loam, 3 to 12 percent slopes	Some	9.48
Pits	No	8.88
Malabon silty clay loam	Some	8.63
Helvetia silt loam, 0 to 12 percent slopes	No	8.62
Steiwer silt loam, 3 to 12 percent slopes	Some	8.32
Helmick silt loam, 3 to 12 percent slopes	Some	8.06
Hazelair silt loam, 3 to 12 percent slopes	Some	7.78
Mcalpin silty clay loam	No	7.43
Conser silty clay loam	Yes	7.07
Willamette silt loam, 3 to 12 percent slopes	No	7.06
Mcalpin silty clay loam, 0 to 3 percent slopes	No	6.12
Chehalis silty clay loam	Some	5.41
Camas gravelly sandy loam	No	4.67
Mcalpin silty clay loam, 3 to 6 percent slopes	No	4.55
Santiam silt loam, 6 to 15 percent slopes	No	4.08
Salem gravelly loam	No	3.43
Suver silty clay loam, 3 to 12 percent slopes	No	3.20
Pilchuck fine sandy loam	No	3.11
Jory silty clay loam, 12 to 20 percent slopes	No	2.98
Santiam silt loam, 3 to 6 percent slopes	Some	2.95
Holcomb silt loam	Some	2.94
Bellpine silty clay loam, 3 to 12 percent slopes	Some	2.81
Jory silty clay loam, 2 to 12 percent slopes	No	2.74
Chehulpum-steiwer complex, 12 to 40 percent slopes	No	2.70
Willakenzie silty clay loam, 2 to 12 percent slopes	No	2.37
Price-ritner complex, 30 to 60 percent slopes	No	2.33

4.8.1a. Soils types most commonly associated with wetlands in the Luckiamute-Rickreall watershed, based on overlay of NWI map of mid-1980's wetlands with county soil maps

Soil map unit name	Hydric?	Acres in Wetlands
Dupee silt loam, 12 to 20 percent slopes	No	1.76
Suver silty clay loam, 12 to 20 percent slopes	No	1.70
Bellpine silty clay loam, 12 to 20 percent slopes	No	1.44
Steiwer silt loam, 12 to 20 percent slopes	No	1.35
Jory silty clay loam, 20 to 30 percent slopes	No	1.19
Hazelair silt loam, 12 to 20 percent slopes	No	1.11
Jory silty clay loam, 2 to 30 percent slopes	No	0.98
Dixonville silty clay loam, 12 to 20 percent slopes	No	0.93
Helmick silt loam, 12 to 20 percent slopes	No	0.89
Chehulpum silt loam, 3 to 12 percent slopes	No	0.83
Willakenzie silty clay loam, 12 to 20 percent slopes	No	0.80
Dixonville silty clay loam, 3 to 12 percent slopes	No	0.63
Rickreall silty clay loam, 3 to 12 percent slopes	No	0.37
Chehulpum silt loam, 12 to 40 percent slopes	No	0.36
Santiam silt loam, 15 to 20 percent slopes	No	0.34
Nekia silty clay loam, 2 to 12 percent slopes	No	0.33
Bellpine silty clay loam, 30 to 50 percent slopes	No	0.33
Ritner-price complex, 12 to 30 percent slopes	No	0.31
Abiqua silty clay loam, 3 to 5 percent slopes	Some	0.27
Price-ritner complex, 20 to 30 percent slopes	No	0.26
Suver silty clay loam, 20 to 30 percent slopes	No	0.17
Bashaw silty clay loam, 0 to 3 percent slopes	Yes	0.17
Woodburn silt loam, 12 to 20 percent slopes	No	0.14
Willamette silt loam, 12 to 20 percent slopes	No	0.04
Willakenzie silty clay loam, 20 to 30 percent slopes	No	0.04

4.8.1b. Area of hydric soils, by subunit, that have or do not have wetlands at the present time. Blank cells indicate partial or complete lack of wetland maps covering that subunit.

HUC	Hydric and have wetlands (acres)	Hydric but lack wetlands (acres)	Total Hydric Area	% of hydrics lacking wetlands
17090003060101				
17090003060102				
17090003060201				
17090003060202				
17090003060203				

4.8.1 b. Area of hydric soils, by subunit, that have or do not have wetlands at the present time. Blank cells indicate partial or complete lack of wetland maps covering that subunit.

HUC	Hydric and have wetlands (acres)	Hydric but lack wetlands (acres)	Total Hydric Area	% of hydrics lacking wetlands
17090003060204				
17090003060301				
17090003060302	0.98	72.02	73.00	98.65
17090003060303	0.00	11.08	11.08	99.99
17090003060304	66.29	53.06	119.35	44.46
17090003060305	25.79	55.67	81.47	68.34
17090003060401				
17090003060402				
17090003060403				
17090003060404				
17090003060501	20.91	60.65	81.56	74.36
17090003060502	0.04	603.06	603.09	99.99
17090003060503	687.48	756.24	1443.72	52.38
17090003060504	599.94	47.85	647.79	7.39
17090003060601				
17090003060602				
17090003060603				
17090003060701	0.00	2.85	2.85	100.00
17090003060702	0.00	41.74	41.74	100.00
17090003060703	0.00	25.56	25.56	100.00
17090003060704	252.78	88.49	341.27	25.93
17090003060801				
17090003060802	14.51	87.39	101.90	85.76
17090003060803	0.00	45.27	45.27	100.00
17090003060901	797.40	22.90	820.30	2.79
17090003060902	509.69	71.75	581.44	12.34
17090003060903	471.78	18.86	490.64	3.84
17090003061001	542.23	256.29	798.52	32.10
17090003061002	437.33	53.76	491.09	10.95
17090003061003	911.54	55.48	967.02	5.74
17090003061004	578.90	45.96	624.86	7.36
17090003061005	721.82	143.10	864.92	16.54
17090003061101	98.08	0.41	98.49	0.42
17090003061102	623.28	16.36	639.63	2.56
17090003061103	789.60	62.37	851.98	7.32
17090003061201	130.15	23.98	154.12	15.56

4.8.1 b. Area of hydric soils, by subunit, that have or do not have wetlands at the present time. Blank cells indicate partial or complete lack of wetland maps covering that subunit.

HUC	Hydric and have wetlands (acres)	Hydric but lack wetlands (acres)	Total Hydric Area	% of hydrics lacking wetlands
17090003061202	438.79	6.47	445.26	1.45
17090003061203	500.81	285.01	785.82	36.27
17090003061204	405.77	457.65	863.42	53.00
17090003061301	1706.57	240.68	1947.25	12.36
17090003061302	923.77	222.27	1146.03	19.39
17090003061303	544.11	80.80	624.91	12.93
17090007020404				
17090007020501				
17090007020502				
17090007020503				
17090007020601				
17090007020602				
17090007020603				
17090007020604				
17090007020605				
17090007020606				

4.8.2 ERODIBLE SOILS

Under some circumstances, soils can move across the landscape and into the stream network where suspended sediments can dramatically affect the quality of aquatic habitat. Circumstances that foster soil erosion include any actions that remove vegetation (which acts to stabilize soils), or any actions that lead to an increased frequency of mass wasting events. Detailed information on soils can be used to plan actions to minimize the effect on soils prone to erosion. For more information on erosion and landslides in the study area, see Section 8, Sediment Source Analysis.

4.9 LITHOLOGY

Many watershed processes are influenced by bedrock lithology. The geologic formations that underlie each watershed determine how groundwater moves (therefore, stream temperature); how stream channels form; how soils form, weather, and erode; and many other watershed characteristics that directly and indirectly influence salmonid habitat.

There are several types of geologic strata in the study area (Table 4.9a). The Tye Formation dominates the southwestern half of the study area. Five pockets of landslide and debris flow deposits are scattered in the western, higher elevation area of the study area. The southern portion of the study area is dominated by Siletz River Volcanics and

related rocks. The northwestern portion of the study area is classified primarily as Yamhill Formation and related rock and Mafic Intrusions. The eastern portion of the study area is primarily Lacustrine and Fluvial Sedimentary rock, Tuffaceous Siltstone and Sandstone, and Alluvial Deposits.

Table 4.9a. Description of the lithology of the study area (OGDC)

PTYPE	LITHOLOGY	DESCRIPTION
Qal	Alluvial deposits	Sand, gravel, and silt forming floodplains and filling channels of present streams; in places includes talus and slope wash; locally includes soils containing abundant organic material and thin peat beds
Qls	Landslide and debrisflow deposits	Unstratified mixtures of fragments of adjacent bedrock; locally includes slope wash and colluvium
Qs	Lacustrine and fluvial sedimentary rocks	Unconsolidated to semiconsolidated lacustrine clay, silt, sand, and gravel; in places includes mudflow and fluvial deposits and discontinuous layers of peat
Qt	Terrace, pediment, and lag gravels	Unconsolidated deposits of gravel, cobbles, and boulders intermixed and locally interlayered with clay, silt, and sand
Ti	Mafic intrusions	Sheets, sills, and dikes of massive granophyric ferrogabbro; some bodies strongly differentiated and include pegmatitic gabbro, ferrogranophyre, and granophyre
Tsr	Siletz River Volcanics and related rocks	Aphanitic to porphyritic, vesicular pillow flows, tuffbreccias, massive lava flows and sills of tholeiitic and alkalic basalt; upper part of sequence contains numerous interbeds of basaltic siltstone and sandstone, basaltic tuff, and locally derived b
Tss	Tuffaceous siltstone and sandstone	Thick to thinbedded marine tuffaceous mudstone, siltstone, and sandstone; fine to coarse grained; contains calcareous concretions and, in places, is carbonaceous and micaceous
Tt	Tyee Formation	Very thick sequence of rhythmically bedded, medium to finegrained micaceous, feldspathic, lithic, or arkosic marine sandstone and micaceous carbonaceous siltstone; contains minor interbeds of dacite tuff in upper part
Ty	Yamhill Formation and related rock	Massive to thinbedded concretionary marine siltstone and thin interbeds of arkosic, glauconitic, and basaltic sandstone; locally contains interlayered basalt lava flows and lapilli tuff

The Tyee and Lacustrine and Fluvial Sedimentary Formations underlie most of the Luckiamute River watershed (Table 4.9a). Steep areas that occur on Tyee (and similar) geologic formations are prone to slide. Interestingly, the Tyee Formation, such an important formation in the Coast Range, does not extend to the Ash Creek watershed. The Ash Creek watershed is dominated by Lacustrine and Fluvial Sedimentary and Tuffaceous Siltstone and Sandstone (Table 4.9b).

Table 4.9b. Lithology of the Luckiamute Watershed study area (OGDC-citation information is below)

Symbol	Lithology	Area (acres)
--------	-----------	--------------

Qal	Alluvial Deposits	7,992.4
Qs	Lacustrine and Fluvial Sedimentary	35,543.4
Qls	Landslide and Debris Flow Deposits	2,845.8
Ti	Mafic Intrusions	16,912.4
Tsr	Siletz River Volcanics and Related Rocks	36,939.6
Qt	Terrace, Pediment, and Lag Gravels	1,355.3
Tss	Tuffaceous Siltstone and Sandstone	15,574.7
Tt	Tyee Formation	55,622.2
OW	Water Body	33.3
Ty	Yamhill Formation and Related Rock	28,918.9

Table 4.9c. Lithology of the Rickreall Watershed study area (OGDC-citation information is below)

Symbol	Lithology	Area (acres)
Qal	Alluvial Deposits	4,451.6
Qs	Lacustrine and Fluvial Sedimentary	14,329.5
Tsr	Siletz River Volcanics and Related Rocks	207.1
Qt	Terrace, Pediment, and Lag Gravels	333.1
Tss	Tuffaceous Siltstone and Sandstone	11,266.3
OW	Water Body	1,119.4
Ty	Yamhill Formation and Related Rock	2,180.3

Understanding patterns in underlying lithology helps with the interpretation other analyses in this watershed assessment. For example, when analyzing the length of streams with gravel substrate in a watershed, it might be useful to consider the total area of igneous versus sedimentary formations within that watershed. Gravels, cobbles and boulders formed from igneous rock tend to be quite durable, compared to those formed from sedimentary formations, which may break down within periods of tens to hundreds of years (Siuslaw National Forest 1997). The general lithology map can help predict and interpret stream channel morphology data and predict where dramatic changes in stream morphology may occur. For example, igneous intrusions such as dikes and sills can create natural barriers to anadromous migration as headward erosion of streams is impeded (Boateng & Associates Inc. 1999). Finally, groundwater flow in some areas may not follow surface features (Siuslaw National Forest 1997), which may help interpret stream water temperature measurements.

The data set used for this analysis is the 1:500,000 scale data set from the Oregon Geospatial Data Center, taken from the Walker and MacLeod 1991, "Geologic map of Oregon" produced by the USGS. A 1:100,000 scale map of the entire state is being created by the Oregon Department of Geology and Mineral Industries, but this endeavor is still in the process of being completed (<http://pubs.usgs.gov/of/of97-269/staub.html>).

4.10 ELEVATION

Elevation is an important property of a watershed. Temperature, precipitation and rain-on-snow patterns are driven by elevation. Precipitation and rain-on-snow events, in turn, control hydrological patterns in the stream network which influence erosion and sediment transport.

The majority of the Luckiamute / Ash Creek study area lies in lower elevations from 105 to 1,000 feet (72.6%), with a smaller area in mid elevations from 1,000 to 2,000 feet (21.7%), and the remainder in the highest elevations of the watershed from 2,000 to 3,650 feet (5.8%). The Ash Creek study area lies within lower elevations from 105 to 1,000 feet, with the majority of the area (91.1%) occurring from 105 to 500 feet (Tables 4.10a & b).

Elevation (ft)	Area (acres)	% Total
105 - 500	87784.0	43.5%
>500 - 1000	58611.7	29.1%
>1000 - 1500	29902.3	14.8%
>1500 - 2000	13938.6	6.9%
>2000 - 2500	7580.2	3.8%
>2500 - 3000	3618.5	1.8%
>3000 - 3560	302.7	0.2%
Total	201737.9	100.0%

Elevation (ft)	Area (acres)	% Total
105 - 500	30859.3	91.1%
>500 - 1000	3028.1	8.9%
Total	33887.4	100.0%

4.11 AQUATIC RESOURCES

4.11.1 HISTORIC AQUATIC RESOURCE CONDITIONS

The characteristics of aquatic resources in the Willamette Valley were historically much different than they are today. Dynamic river processes, such as frequent flooding events, maintained off channel habitat including side channels, alcoves, sloughs and shallow lakes (Taft and Haig, 2003). Prior to the early 1800s, when the fur trade expanded into this area, beaver contributed to stream complexity by ponding water. Fallen snags and debris jams also created pools of standing water (Taft and Haig, 2003). As farming took hold in the valley and wetlands were drained, much of the side channel habitat was reduced. Large wood was also removed from the streams in order to straighten channels.

Before Himalayan blackberry became omnipresent in bottomlands in the mid-20th century, the dominant riparian shrubs were probably hawthorn (*Crataegus* spp.), hardhack (*Spiraea*), serviceberry (*Amelanchier*), alder (*Alnus*), dogwood (*Cornus*), snowberry (*Symphocarpos*), and willow (*Salix* spp.). Before reed canary-grass blanketed nearly every channel bank and wetland, sedges (*Carex* spp.) and rushes (*Juncus* spp) were probably more widespread than at present. In the overstory, huge cottonwoods (*Populus*), Douglas-fir (*Pseudotsuga*), and grand fir (*Abies*) were more prevalent than today.

Several early logging practices relied on rivers as holding areas and transport system, and greatly affected stream habitat and riparian areas. Small dams known as “splash” dams were built in streams to transport logs down the river (Theurer, 2003). Splash dams got their name from the wave of water preceding the logs as they rushed downstream. Dams were used to create ponds where the logs could float until a log-drive began. To release the logs, some splash dams were dynamited, which resulted in a torrent that carried the logs downstream, other dams were released in a more controlled fashion, and used repeatedly. Log drives involved creating rafts of logs that were then sent down river. The logs, driven by high, fast water, removed riparian vegetation and scoured the stream bottom leading to erosion and loss of instream habitat (Theurer, 2003).

The advantage of splash dams was that it did not require sophisticated tools or machinery to move logs from the forests to the mills. The disadvantage was that the use of splash dams was damaging to stream ecosystems in many ways; many streams in Oregon still bear the signs of splashing damming. Unintended results of splash damming included: the down-cutting of stream channels, scouring the creek bottoms, sometimes to bedrock; loss of natural logjams; loss of deep pool habitat; stream channel channelization; loss of stream side vegetation and eroded stream banks; and barriers to migrating salmon. The damage must have been quite apparent because there are reports of farmers in the valley complaining that the log drives were causing erosion along the banks of their land on the river (Frantz and Brandon, 1976). The number of complaints eventually led to legal battles to stop log drives beginning in 1914; however, log drives continued until 1925 (Theurer, 2003).

Historic records indicate that there were from 80-100 splash dams on the Luckiamute River (Licata *et al.*, 1998), with some major ones documented on Pedee Creek and Ritner Creek (Theurer, 2003). A larger dam was located near the mouths of the north and south fork of the Luckiamute River, where “the water backed up nearly to Camp Walker” (Theurer, 2003).

Splash dams were not the only log transport tool associated with stream degradation. Another tool in early logging, the donkey engine, also affected instream habitat. These steam-driven engines were set up in a canyon or stream bottom. Logs were attached to the engine and dragged along the ground and along streambeds. This probably resulted in loss of structure from stream bottom. The use of donkey engines may have increased landslide frequency and contributed to sediment delivery in streams (Licata *et al.*, 1998).

Other early settlement practices such as allowing livestock to trample streamside vegetation and enter waterways, and clearing of vegetation for homesteads and agriculture also affected stream habitat. The clearing of the land eliminated most large riparian conifers which had a potential to fall into streams to serve as large woody debris (Licata *et al.*, 1998).

4.11.2 CURRENT AQUATIC RESOURCE CONDITIONS

4.11.2.1 Streams

Streams are referred to by stream order. Headwater streams are known as first order streams. Where two first order streams join, they form a second order stream, and so on. In general, higher order streams are large and lower order streams are small. Because the streams form a network, factors that influence the headwaters of a stream also affect higher order streams. Thus, higher order streams are said to “express” the cumulative effects of the entire watershed. Tables 4.11.2.1a & b summarizes the total length of streams in the study area based on stream order.

Most streams in the Luckiamute/ Ash Creek study area are first order streams. Woody debris, gravel, and sediments enter the stream network from first order streams. In addition, shade along first order stream will help to keep water temperatures cool. Although salmon do not ordinarily have access to first order streams, these streams are an important, and frequently overlooked, component of salmonid habitat. We recommend that first order streams be evaluated for shade and for the potential to deliver large wood to the stream networks.

Table 4.11.2.1a. Stream length by order for the Luckiamute Watershed study area (Miller *et al.*, 2001)

Stream Order	Length (ft)	Length (mi)	% Total
1	2,565,780.0	485.9	49.5%
2	1,175,520.6	222.6	22.7%
3	620,055.0	117.4	12.0%
4	311,985.3	59.1	6.0%
5	297,699.3	56.4	5.7%
6	197,966.1	37.5	3.8%
7	13,260.4	2.5	0.3%
Total	5,182,266.7	981.5	100.0%

Increasing stream size ↓

Table 4.11.2.b. Stream length by order for the Ash Creek Watershed study area (Miller *et al.*, 2001)

Stream Order	Length (ft)	Length (mi)	% Total
1	244,007.0	46.2	45.7%
2	125,478.6	23.8	23.5%
3	73,818.6	14.0	13.8%
4	75,862.4	14.4	14.2%

Increasing stream size ↓

5	15,094.0	2.9	2.8%
Total	534,260.6	101.2	100.0%

USGS operates a streamflow gauging station on the mainstem of the Luckiamute River near Suver at an elevation 171.9 ft. This station, number 14190500, has been in operation since 1874 and measures the discharge for an area of 240 mi². USGS has recorded “acceptable” data only from the period 1940 to 1988. Data on daily flow, monthly means and yearly means are available (see Section 6.4).

4.11.2.2 Lakes and Reservoirs

There are a few major reservoirs in the Luckiamute/ Ash Creek study area. These include Buchanan Reservoir and Hamilton Reservoir northeast of Airlie, McCrae Reservoir and Bauman Reservoir west of Monmouth, Fall City Reservoir south of Falls City, Lake of the Winds behind the dam on Burgett Creek, and Emry Pond/Moore behind the dam on the tributary of Maxfield Creek (see Section 6.7.2).

4.11.2.3 Springs

Springs provide cold water to streams and can lower stream temperatures. We examined USGS 7.5' topographical maps and found that there were twenty-six springs identified (Table 4.11.2.3). Only a few of the springs were named including: Cauthorn, Cold, Crystal, Fort Hoskins, Maple, Nelson, Rattling, Sulfur, Thistledew, and Vitae Springs. Spring water can be used, in some cases, without obtaining a water use permit (see Section 6.5 on Water Use).

7th Field Watershed	Sub-Basin Name	Number	Feature
17090003060203	Hoskins	2	Spring
17090003060204	Vincent Creek	1	Spring
17090003060301	Plunkett Creek	2	Spring
17090003060401	Unnamed	2	Spring
17090003060404	Ritner Creek	2	Spring
17090003060502	McTimmonds	3	Spring
17090003060503	Middle Luckiamute	1	Spring
17090003060602	Cold Springs	3	Spring
17090003060702	Falls City	1	Spring
17090003060802	Lower Teal Creek	1	Spring
17090003060803	Grant Creek	2	Spring
17090003060901	Fern Creek	1	Spring
17090003060902	Lower Little Luckiamute	3	Spring
17090003061101	Upper Soap Creek	1	Spring
17090003061103	Rifle Range	1	Spring
	<i>Total</i>	<i>26</i>	<i>Total</i>

We recommend that these sites be verified and that the condition of the springs be recorded, if possible.

4.12 FISH AND WILDLIFE

Fish and wildlife are of interest to the Luckiamute Watershed Council because of their aesthetic, recreational, and economic importance. In addition, particular groups of species, such as PNW salmon, are also of interest because they act as indicators of environmental quality. PNW salmon have complex life history strategies which depend on the presence of suitable environmental conditions in freshwater, estuarine, and marine environments in order to successfully complete their life cycles. If the ecological requirements in any one of their life history stages (e.g., spawning, rearing, or growth) are not met, their populations will decline. Thus, salmon are sensitive to conditions in streams where they spawn and rear in the upper watershed, tidal marshes in the estuary, and the ocean environment.

Fish and wildlife populations can be characterized in several different ways. Abundance and distribution can be directly observed either anecdotally (casually) or through scientific study. In many cases, biologists qualitatively assess environmental conditions and use their best professional judgment to determine where populations of organisms are likely to be (see Section 7.11). Alternatively, populations can be indirectly assessed by understanding the relationship that exists between an organism and its habitat through the use of a habitat suitability model. Numerous Habitat Suitability Index models have been developed. Some models have been rigorously developed using quantitative data and empirical species-habitat relationships (see (Johnson and O'Neil, 2001) and others tend to be more descriptive and of unknown accuracy. Finally, historic population levels of fish and wildlife are known through archeological studies. For this report, we have acquired and evaluated information developed using all of these approaches.

4.12.1 FISH

An understanding of both natural and anthropogenic (human influenced) factors is necessary to explain the current distribution of fish species in the Luckiamute / Ash Creek study area. For example, Willamette Falls (at RM 48) has selectively blocked runs of anadromous fish from reaching upriver areas including the Luckiamute study area. While anadromous runs of winter steelhead and spring chinook are able to pass Willamette Falls during seasonal high water flows, other runs are not (Mattson and Gallagher, 2001). Archeological evidence suggests that this has been the case since pre-historic times. Archeological sites in the Willamette Valley show evidence of prehistoric fishing, including bone points that represent parts of composite harpoons or fish spears, and grooved pebbles that may have served as sinkers (Aikens, 1986); however, salmon did not seem to be as important a food source to the early peoples in Willamette Valley as they were to the peoples living along the Columbia River and coastal Oregon. Moreover, early accounts of European-American travelers in the Willamette Valley rarely mention fish

(Aikens, 1986) suggesting that salmon were not as abundant in the Willamette Valley as they were in other parts of the state.

Fish species found in the Luckiamute / Ash Creek study area today either occur there naturally or they were stocked in the watershed (or nearby areas). In all cases, environmental conditions within the watershed must fulfill the ecological requirements of each species for populations to persist. For example, fall chinook salmon, not naturally occurring in the basin, were released into the Luckiamute and Little Luckiamute Rivers in May 1974 and 1976; however, these populations failed to become established, perhaps due to the low flows and high water temperatures characteristic of the subbasin (Wevers *et al.*, 1992).

The abundance and distribution of the fish species occurring in the study area is not well known. A recent watershed assessment by the BLM states that, "There are no known data pertaining to populations of these resident fish. However, it is believed that many second-order, and all third-order streams (*i.e.*, those having gradients < 8 percent) and below have fish present." (Licata *et al.*, 1998). To further complicate matters, uncertainty pertaining to the origin stocks arises when past stocking practices supplemented naturally occurring species with stocked varieties of the same species. For example, juvenile coho were found in the Luckiamute River prior to 1955 before any releases were made and the origin of these fish remains unknown: they were most likely strays from other subbasins (Wevers *et al.*, 1992).

The distribution of salmonids in the watershed is not well known. The Oregon Department of Fish & Wildlife (ODFW) publishes several fish distribution maps showing their best professional judgment on the current distribution of coho and winter steelhead salmon (Maps 9-11). In general, the distribution of fish would be limited by unfavorable environmental conditions in the upper stream reaches or by a barrier (see Section 6.7.3: Barriers). In the Luckiamute River basin, winter steelhead are distributed along the Little Luckiamute and Luckiamute Rivers, and along Soap Creek. Winter steelhead also occur along the lower reaches of Ash Creek. Winter steelhead are believed to be distributed along 25.5% and 0.7% length of the streams (1:100K) in the Luckiamute and Ash Creek basins, respectively (Map 9). The distribution of coho are also mapped by ODFW although they are not native to the Luckiamute and Ash Creek study. Coho distribution in the Luckiamute is similar to that of winter steelhead, occupying slightly less (20.1% of the length of the 1:100K streams) of the available stream length (Map 10). Coho do not occur in Ash Creek. Spring chinook are distributed along the lower reaches of the Luckiamute River, and Soap and Ash Creeks. Spring chinook are distributed along approximately 2.5% of the streams (1:100K) in the Luckiamute and 0.6% of the streams in the Ash Creek watersheds (Map 11). Interestingly, cutthroat trout are widely distributed in the study area, even above some of the barriers that block other species. In fact, isolated populations of cutthroat exist in the Little Luckiamute drainage. Population densities of cutthroat above the falls on the Little Luckiamute are much greater than other streams in the study area (Wevers *et al.*, 1992).

The Luckiamute Watershed Council Technical Advisory Team asked for information on the fish species shown in Table 4.12.1. This table is not meant to include all the fish species known from the watershed.

Table 4.12.1: Important fish in the study area. Shown are common name, scientific name, whether the fish is native to the study area, where it occurs in the watershed and other notes.

Species	Scientific Name	Native to Luckiamute	Stocked	Watershed Use	Notes
Salamonids					
Winter Steelhead	<i>Oncorhynchus mykiss</i>	Y	1964-1982	Occur in upper reaches	Occur in the upper reaches of the watershed.
Cutthroat Trout	<i>Oncorhynchus clarki</i>	Y	1920's-	Occur in most perennial steams and some intermittent streams.	Not anadromous above Willamette Falls. Isolated populations occur above barriers in Little Luckiamute, Teal, Burgett and Rock Pit Creeks.
Coho	<i>Oncorhynchus kisutsh</i>	N	1920-1980's	Occur in most streams including small first and second order streams	Coho may compete with native cutthroat and winter steelhead in the basin.
Spring Chinook	<i>Oncorhynchus tshawytscha</i>				
Fall Chinook	<i>Oncorhynchus tshawytscha</i>	N	1974; 1976		Not believed to be in basin today. Failed to become established due to low flows and high temps.
Rainbow Trout	<i>Oncorhynchus mykiss</i>	N	1920's-1990's		Stocked in Luckiamute and Little Luckiamute Rivers.
Other Species					
Pacific Lamprey	<i>Lampetra tridentata</i>				
Western Brook Lamprey	<i>Lampetra richardsoni</i>				
Whitefish		Y			Member of the trout family and have been observed in the Little Luckiamute River.
Oregon Chub	<i>Oregonichthys crameri</i>	Y	No		Found in sand and gravel pools and backwaters of creeks and small rivers, often in vegetation. Currently there are no known populations of the Oregon Chub in the study

Table 4.12.1: Important fish in the study area. Shown are common name, scientific name, whether the fish is native to the study area, where it occurs in the watershed and other notes.

Species	Scientific Name	Native to Luckiamute	Stocked	Watershed Use	Notes
					area.
Sandroller	<i>Percopsis transmontana</i>		No		Found in low gradient streams, active at night.
Speckled Dace	<i>Rhinichthys osculus</i>				
Sculpin (several species)					
Warm Water Fish					
Largemouth Bass	<i>Micropterus salmoides</i>	N		Shallow weedy lakes and backwater areas of large rivers.	Prefers water temps 60°F
Smallmouth Bass	<i>Micropterus dolomieu</i>	N	Introduced to US West in late 1800's	Streams with alternating pools and riffles, lakes and reservoirs.	Prefers water temps 55-65°F
Black Crappie	<i>Pomoxis nigromaculatus</i>	N			
White Crappie	<i>Pomoxis annularis</i>	N			
Bluegill	<i>Lepomis macrochirus</i>	N			
Pumpkinseed	<i>Lepomis gibbosus</i>	N			
Warmouth	<i>Lepomis gulosus</i>	N			
Yellow Perch	<i>Perca flavescens</i>	N			
Brown Bullhead	<i>Ameiurus nebullosus</i>	N			

^A=ODFW. 2000. Listed Species of Fish in Oregon. ^B=Wevers et al. 1992

4.12.1.1 General Fish Habitat Summary

For an individual salmonid to hatch, grow and return to a stream to spawn specific ecological conditions must be met during each of its life history stages. Typically, the

following life history stages are recognized for salmonids: spawning, incubation, emergence, summer rearing, and overwintering. Each species of salmonid has different ecological requirements and may be more or less vulnerable in each of these stages than another species in a particular watershed. The interplay between a salmonid's life history requirements and its environment (*i.e.*, watershed, estuary, and ocean) lead to a diversity in life history strategies among different populations of salmon. These differences permit one species of salmonid to do better than another within the same basin, or one basin to support higher populations of a particular species than an adjacent basin. The relationship between salmonid populations and their environments lead to genetically distinct (locally adapted) populations.

Fisheries biologists generally determine the habitat requirements of salmonids by matching abundance and distribution data with the physical and biological characteristics of where the fish are observed. Observed habitat-species relationships guide much of the salmonid restoration plan in the PNW. In addition to observational data, laboratory studies can also be used to determine the physiological response to fish to environment factors. For example, lethal and sub-lethal water temperature guidelines were probably determined from laboratory studies. In all cases, biologists are particularly interested in determining what environmental factors limit the abundance and distribution of salmonid species. In-stream complexity (used by salmonids to escape periods of high water flows, to forage, to escape predation, *etc.*) is generally believed to be the factor that most frequently limits salmonid populations in the PNW. For this reason, most salmonid habitat restoration efforts center on enhancing or creating in-stream structure. In-stream structure, in turn, can lead to the development of pools and can trap and sort substrates (gravel beds used for spawning: see Sections 4.4.1.1 and 7.1.2).

One source of information describing the condition of salmonid habitat is the ODFW Aquatic Habitat Inventory (AHI) data sets (see Section 6.1.1.2). AHI data are gathered in the field by county, state and federal agencies and by private industrial groups using similar protocols. Field teams make measurements and observations on stream gradient, substrates, channel form, stream-side vegetation, *etc.* Data are organized on USGS topographic maps at 1:24K. ODFW then converts field data to GIS data sets. There are two things to be aware of when using AHI GIS data. First, there are two different data sets, habitat unit-level and reach-level GIS spatial data sets. Habitat units, the fundamental surveyed units, are defined by the surveyor based on breaks in geomorphology, flow characteristics, *etc.* (Moore *et al.*, 2002). Reach-level summaries, produced by ODFW, are based on summaries of habitat unit data. Second, all field data are 'calibrated': calibration consists of converting field measured distances to map distances. Therefore, AHI GIS layers should not be viewed as spatially accurate descriptions of instream habitat, but rather a generalized description of the patterns of instream habitat.

The area of the Luckiamute and Ash Creek basins covered by AHI data is quite limited (Maps 9-11): approximately 12.2% of the streams (1:100K) have been surveyed by AHI field crews. Aquatic habitat conditions can reflect watershed processes that occur over large areas. However, when only a limited proportion of the stream network is surveyed,

it is possible that the conditions found may not be typical of the entire watershed. In addition, some of the surveys from the Luckiamute / Ash Creek study area are more than nine years old. While these data will remain useful until they are replaced by more recent surveys, it is important to note that the Coast Range of Oregon is a dynamic environment where the conditions reported in some of the older surveys may no longer be accurate.

The primary advantage of using AHL data is that the condition of streams across the state can be evaluated as salmonid habitat by matching the ecological requirements of salmonids with in-stream conditions. Scientists at ODFW and NMFS are currently working on a model that ranks data collected during AHL into 'good', 'fair' or 'poor' habitat for steelhead, chinook and sockeye salmon. Although this model is being developed for the Middle Deschutes River Basin it is "intended for general application to the Pacific Northwest Basins" (Burke *et al.*, in prep). An advantage of this model is that it can be tied to existing GIS coverages created from AHL data to give a quick overview of habitat quality in a particular area for each of three life stages: spawning (spawning+ incubation+ emergence), summer rearing and overwintering.

Jennifer Burke *et al.* (in prep) have reviewed the literature and developed a set of optimal habitat requirements for various life stages of steelhead, chinook, and sockeye salmon. For steelhead, Burke *et al.* summarized ecological requirements for substrate, pool area and depth, temperature, flows, large woody debris, and cover and developed a Habitat Quality Rating Model (HabRate). This model was developed as a decision making tool and is "intended to provide a qualitative assessment of the habitat potential of stream reaches" (Burke *et al.*, in prep). This spreadsheet model is ideally suited for interpreting AHL data. We developed a reach-level ranking of AHL data from the Luckiamute and Ash Creek study area using criteria established in this model for steelhead (see Section 7).

Unfortunately, a similar model was not available for cutthroat trout, chinook or coho, the other salmonids found in the Luckiamute / Ash Creek study area. Cutthroat trout are widely distributed in the Luckiamute and Ash Creek basins. They occur in most perennial streams and occasionally in intermittent streams. Spawning occurs in November in the Luckiamute River although they are known to spawn earlier in other areas (Table 4.12.1.1). Following spawning, cutthroat drop back to the Willamette River in late March (in (Wevers *et al.*, 1992). Cutthroat have been observed to spawn in Soap Creek from January through May.

Fall chinook salmon require higher water flows and lower water temperatures than those occurring in the study area. Releases of fall chinook failed to establish viable populations in the Luckiamute and Little Luckiamute rivers in the mid 1970's (Wevers *et al.*, 1992).

Coho have similar ecological requirements to cutthroat and winter steelhead in the Coast Range Basin. Competition between coho and native salmon is believed to have limited the successful establishment of coho populations in the study area (Wevers *et al.*, 1992).

Table 4.12.1.1 shows generalized spawning times for many of the fish found in the study area.

Table 4.12.1.1: Spawning times for selected fish found in the Luckiamute and Ash Creek watersheds.												
Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Winter Steelhead	Spawning											
Cutthroat Trout	Spawning									Spawning		
Coho										Spawning		
Spring Chinook	Spawning											
Fall Chinook	Spawning							Spawning				
Rainbow Trout	Stocked											
Oregon Chub			Spawning									
Sandroller					Spawning							
Warm Water Fish												
Largemouth Bass				Spawning								
Smallmouth Bass				Spawning								

4.12.1.2 Historic Catch Records (steelhead, cutthroat)

There is information on the status of salmonid populations in the Pacific Northwest; however, much of this information is anecdotal. Fish populations are frequently assessed using a variety of different survey methods, including catch data, dam counts, and more formalized juvenile counts and spawning surveys. Unfortunately, many of these survey methods, like most sample methods, include some sort of sampling bias. For example, catch data may be influenced by conditions other than the abundance of fish. Catch records are frequently used to assess the status of game fish populations. Catch data also are difficult to standardize; however, catch data are often expressed as the number of fish caught per level of effort, usually per angler hour. However, many factors can influence the number of fish that are actually caught. This makes catch records an unreliable tool to assess populations.

Many other survey techniques also have sampling bias. For example, some spawning surveys are frequently conducted along subjectively selected stream segments and therefore are not suitable for use in developing accurate, basin-wide estimates of fish populations. In addition, reported results from many surveys may incorporate of some sort of “correction factor” intended to account for sample bias. Examples of correction factors commonly used include mortality estimates, exploitation rates, and/or bias correction (Botkin *et al.*, 1993). While correction factors are not entirely bad, these factors are often employed without being defined or their assumptions being documented. This makes it very difficult to determine and interpret what was actually measured. Most importantly, one has to accept population size estimates at face value without know how accurate or representative they are (see Section 3.2.3).

Winter steelhead harvest, determined from catch records, in the Coast Range subbasin ranged from 24 to 262 tags during 1977 to 1989. Catch records for winter steelhead for the Luckiamute River are shown in Fig. 4.12.1.2.




Figure 4.12.1.2. Corrected Winter Steelhead Harvest in the Luckiamute River from 1976-1989.

4.12.1.3 Historic/Recent Juvenile and Spawner Surveys

Surveys of both juvenile and spawning adults are also used to assess fish populations. Many of these techniques also have bias, and knowledge of protocols and data handling methodologies are necessary before results can be interpreted and compared. This criticism was recognized by ODFW personnel in 1980 when they made several recommendations to improve accuracy and precision of coho surveys. These improvements included the expansion of the number of index streams, to replace peak counts with estimates derived from Area-Under-the-Curve (AUC) techniques, and to separate indices from streams influenced by hatchery fish from others (Wemple, 1994). Therefore, care must be taken when interpreting and comparing earlier records.

Unfortunately, for the Luckiamute / Ash Creek study area not much is known about recruitment of winter steelhead and cutthroat. Moreover, the release of winter steelhead into the basin from 1964 to 1982 and cutthroat stocking in the early 1920s would make trend analysis of juvenile counts problematic. One way to estimate potential recruitment is from the number of redds. Wevers *et al.* (1992) summarized the number of winter steelhead redds per mile for the Luckiamute River between 1985 and 1991. The

average number of redds per mile was 10.6 and the number redds per mile peaked in 1988 but dropped to the lowest number during the period of record in 1990 (Fig 4.12.1.3). The ODFW basin plan did not include information for cutthroat.




Figure 4.12.1.3. Number of winter steelhead redds per mile for the Luckiamute River (Wevers *et al.* 1992). The average number of redds per mile was 10.6.

4.12.1.4 Hatcheries

No hatcheries are located in the Coast Range Subbasin (Wevers *et al.*, 1992), the area in which the Luckiamute and Ash Creek Study area is located. No hatchery cutthroat trout or whitefish have been released in the study area. However, several nearby hatcheries supplied smolts in the recent past which were released in or near the study area. Winter Steelhead were released from the Big Creek and Klaskanine (sic?) hatcheries or the Roaring River Hatchery. Coho released into the subbasin were supplied from the Bonneville, Oxbow, Eagle Creek, Cascade, and Sandy hatcheries. Coho eggs used in the STEP program were supplied by the Sandy Hatchery or from the Cowlitz Hatchery (WA). Fall Chinook were introduced to the Luckiamute in the mid 1970's from the Cowlitz Hatchery. Rainbow trout currently released into the subbasin are supplied by the Roaring River Hatchery. Non-native warmwater game fish are obtained from hatcheries and released into the subbasin (Wevers *et al.*, 1992).

4.12.1.5 Endangered/Threatened Fish Species and Species of Concern

Much of what is being done to manage and restore watersheds in Oregon is the result of the Oregon Plan. The Oregon Plan can be traced back to 1996 when then Governor John Kitzhaber initiated Oregon's Coastal Salmon Restoration Initiative (OCSRI). The goal of OCSRI was to develop a plan to restore the vitality of wild salmon, steelhead, and cutthroat trout in coastal watersheds. The OCSRI fostered active partnerships between state and federal agencies, local governments, conservation organizations, industry representatives, watershed councils, and private landowners. The goal of OCSRI was to develop a plan to restore the vitality of wild salmon, steelhead, and cutthroat trout in coastal watersheds.

Elements of the OCSRI Plan included:

- Specific actions to conserve "core" populations of salmon.
- Procedures to provide continuing leadership and improve interagency cooperation.
- Adjustments in harvest management and hatchery programs.
- Goals for riparian management in land-use planning.
- Measures to improve the condition of streams and riparian habitats.
- Proposals for funding and economic incentive programs.
- Opportunities to improve compliance with existing environmental laws.
- Public education programs.
- A proposal describing a comprehensive monitoring program.
- Descriptions of watershed council restoration projects.

The multifaceted OCSRI was later renamed the Oregon Plan. The Oregon Plan now forms the basis for salmon recovery strategies in Oregon.

Table 4.12.1.5 shows the current status, state and federal, as well as sensitive species (designated by ODFW).

Table 4.12.1.5: Current status of selected fish species.			
Species	Scientific Name	ODFW Status	Federal Status ^A
Winter Steelhead	<i>Oncorhynchus mykiss</i>	Not Listed	Threatened
Cutthroat Trout	<i>Oncorhynchus clarki</i>	ODFW species of concern ^B	
Spring Chinook	<i>Oncorhynchus tshawytscha</i>	Not Listed	Threatened
Oregon Chub	<i>Oregonichthys crameri</i>	Not Listed	Endangered
Pacific Lamprey	<i>Lampetra tridentata</i>		
Sandroller	<i>Percopsis transmontana</i>	ODFW stock of concern ^C	

Species	Scientific Name	ODFW Status	Federal Status ^A
---------	-----------------	-------------	-----------------------------

^A=ODFW. 2000. Listed Species of Fish in Oregon. ^B=BLM Watershed Analysis ^C=Wevers et al. 1992

4.12.2 WILDLIFE

4.12.2.1 Insects

Although not generally regarded as “wildlife,” terrestrial and aquatic invertebrates (including insects) are numerically the largest component of the Luckiamute’s biodiversity. Moreover, many insect species help control weeds and pest insects, as well as pollinate plants, aerate soil, support fish and other wildlife, serve as indicators of ecosystem health, and provide aesthetic enjoyment (e.g., butterflies).

Data Availability

No comprehensive surveys have been undertaken of invertebrate or insect diversity in the Luckiamute watershed, or of the distribution and prevalence of particular insect pests. Lists of aquatic invertebrates may be available based on samples collected in a few Luckiamute tributaries and wetlands by DEQ, EPA, OSU, and WOU investigators.

Species Status and Geographic Patterns

Of particular concern are invertebrate species that are rare or believed to have declined severely or disappeared entirely from the region in recent years. The ORNHIC database includes six such species that have been reported specifically from the Luckiamute Watershed (Table 4.12.2). It is virtually certain that other rare or declining invertebrate species are present in the watershed but are not included in this table because of the lack of a survey covering all watershed lands.

Table 4.12.2. Notable invertebrate species or subspecies from the Luckiamute Watershed reported in the ORNHIC database.

Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Driloleirus (Megascolides) macelfreshi</i>	Oregon giant earthworm	G1 S1; 1 Fed=SOC	Last reported before 1985
<i>Icaricia icarioides fenderi</i>	Fender’s blue butterfly	G5T1 S1; 1 Fed=LE	Foothill meadows (upland prairies) with Kincaid’s lupine. See: http://www.epa.gov/fedrgstr/EPA-SPECIES/1998/January/Day-27/e1851.htm
<i>Speyeria zerene bremneri</i>	Valley silverspot butterfly	G5T3T4; 2 SH	No recent records; presumed extirpated. Wet prairies and marshes.
<i>Speyeria callippe</i> ssp.	Willamette callipe fritillary butterfly	G5TH SX; 1	No recent records; presumed extirpated
<i>Euphydryas editha taylori</i>	Taylor’s checkerspot butterfly	G5T1 S1; 1 Fed=SOC	Dry prairies, open oak stands

Table 4.12.2. Notable invertebrate species or subspecies from the Luckiamute Watershed reported in the ORNHIC database.			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Rhyacophila fenderi</i>	Fender's caddisfly**	unlisted	Rapid streams
<p>* LE= legally listed as Endangered; LT= legally listed as Threatened; SOC= Species of Concern; SU= status undetermined; SV= vulnerable; SC= critical. Critical habitat can be protected legally only for LE and LT species.</p> <p>G= global status (ONHP), S= state status (ONHP). 1= critically imperiled; 2= imperiled and vulnerable to extinction; 3= rare but not immediately imperiled, 4= not rare but of long-term concern, 5= widespread and secure</p> <p>Number following the semicolon: 1= extinction threatened or presumed, 2= extirpation from Oregon threatened or presumed, 3= insufficient information, 4= of concern but not immediately imperiled</p>			

Important Habitats and Communities

Almost any area of the watershed that has experienced minimal soil disturbance, is distant from areas of pesticide application, has unpolluted water, and/or retains predominantly native vegetation could be important for maintaining the watershed's overall diversity of native invertebrates. Remnant native prairie – a particularly rare habitat -- is especially important to several species. Just north of the Luckiamute Watershed, the conversion to wheat field of part of one such prairie near Buell in the mid-1990's is believed responsible for the disappearance of a rare species in that area (US Fish & Wildlife Service: <http://www.epa.gov/fedrgstr/EPA-SPECIES/1998/January/Day-27/e1851.htm>).

Conservation and Restoration Potential

Projects to restore native prairies, wetlands, and oak woodlands are especially beneficial to rare invertebrate species. Improvements in stream water quality will benefit a wide variety of aquatic invertebrates. Efforts to encourage retention of downed dead wood and planting of a diversity of native vegetation in yards and other horticultural settings will benefit many species.

4.12.2.2 Amphibians and Reptiles

This section covers turtles, lizards, snakes, salamanders, frogs, and toads. Species in these groups not only contribute to the Luckiamute Watershed's biodiversity, but also help reduce pest insects and maintain healthy aquatic and terrestrial ecosystems.

Data Availability

No comprehensive surveys have been undertaken of amphibian and reptile diversity in the Luckiamute watershed. Surveys of amphibians and reptiles have been conducted at specific locations in the watershed, such as the Camp Adair Military Training Area west of Coffin Butte (Henny *et al.* 1999) and [McDonald-Dunn Forest?]. Also, reports covering larger regions that include the Luckiamute have been published (e.g., St. John 1987, Vesely *et al.* 1999, Adamus 2003), and anecdotal observations of amphibians and reptiles have been recorded at the E.E. Wilson Wildlife Area, portions of Boise timber land, and a few other locations.

Species Status and Geographic Patterns

A total of 14 amphibian and 13 reptile species are known to occur within the Luckiamute watershed (Appendix E: Amphibian and Reptile). Although no field data are available to depict their overall distribution within the watershed, we used a modeling approach to tentatively identify patterns of “weighted species richness” (WSR) (Figure xx). Areas with higher WSR scores (colored orange or red on the accompanying maps) can be assumed to individually be providing better habitat to a larger number of amphibian and reptile species. In general, western parts of the watershed have greater WSR, as do riparian areas, and the WSR of the Luckiamute watershed is mostly greater than that of the Rickreall (Ash Creek) watershed.

To compute WSR, the author (with subsequent review by a team of herpetologists) rated the land cover type predominating in each 30 x 30 m square (pixel). The land cover type was rated on a 1-10 scale with regard to its suitability for each of the 14 amphibians and 13 reptiles, and then those scores of the 27 species were (a) summed, and (b) averaged. Only the summed scores are depicted in Figure xx. This spatial modeling approach allowed us to systematically consider the adjacency of each pixel to other pixels and features (e.g., streams) of synergistic habitat types when scoring each species in each pixel. Image analysts at the Forest Sciences Laboratory, Oregon State University, had initially assigned each pixel to one of about 30 land cover types based on its condition as detected by satellite imaging in the spring and summer of 1992. Note that many features important to reptiles and amphibians cannot be detected directly by satellite imagery. Also note that some cover types, while not supporting large numbers of species, may provide the only quality habitat for a few species and so may be quite significant; this is not reflected by the modeling approach. Maps for individual species could be generated if desired.

Of particular concern are species that are rare or believed to have declined severely or disappeared entirely from the region in recent years. The ORNHIC database includes four such species that have been reported specifically from the Luckiamute Watershed (Table 4.12.2.2a). It is likely that other rare or declining amphibians and reptiles are present in the watershed but are not included in this table because of the lack of a survey covering all watershed lands.

Table 4.12.2.2a. ORNHIC database records of notable amphibian and reptile species reported from the Luckiamute Watershed			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Emmys marmorata marmorata</i>	Northwestern pond turtle	G3T3 S2; 1 ODFW= SC Fed= SOC	Ponds or wetlands near permanent water, especially where only partially surrounded by woodland.
<i>Chrysemys picta</i>	Painted turtle	G5 S2; 2 ODFW= SC	Ponds or wetlands near permanent water, generally in wooded areas
<i>Contia tenuis</i>	Sharptail snake	G5 S3; 4 ODFW= SV	Meadows, prairies, woodland edges

Table 4.12.2.2a. ORNHIC database records of notable amphibian and reptile species reported from the Luckiamute Watershed			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Ascaphus truei</i>	Coastal tailed frog	G4 S3; 2 ODFW= SV Fed= SOC	Large, wooded, moderately steep streams
<p>* SOC= Species of Concern; SU= status undetermined; SV= vulnerable; SC= critical. These state (ODFW) and federal (Fed) designations are advisory only and confer no additional legal protection to the species. G= global status (ONHP), S= state status (ONHP). 1= critically imperiled; 2= imperiled and vulnerable to extinction; 3= rare but not immediately imperiled, 4= not rare but of long-term concern, 5= widespread and secure</p> <p>Number following the semicolon: 1= extinction threatened or presumed, 2= extirpation from Oregon threatened or presumed, 3= insufficient information, 4= of concern but not immediately imperiled</p>			

Pest Species

The watershed hosts no reptile or amphibian species that cause significant economic harm. Bullfrogs are not native to the Willamette Valley, but because of their status as predators are believed to be having major effects on some native amphibians and reptiles, e.g., red-legged frog, sub-adult pond turtles. Also, a variety of non-native turtles kept as pets have been illegally released into the wild, and are believed to sometimes harbor diseases that harm native species.

Important Habitats and Communities

Almost any area that has experienced minimal soil disturbance and retains predominantly native vegetation could be important for maintaining the watershed’s overall reptile and amphibian diversity. In particular, mature conifer forests, oak woodlands, prairies, springs, wetlands, ponds, riparian areas, unpolluted streams, and rock ledges and quarries are particularly likely to support species not found broadly in the watershed.

Conservation and Restoration Potential

Projects that restore native prairies, wetlands, and oak woodlands will benefit many native reptiles and amphibians. Efforts to encourage landowners to retain and pile (rather than burn) brush and downed dead wood on their land also will be a help. Root wads and large limbs from logging operations could be donated to pond owners willing to provide essential basking sites for rare western pond turtles (Adamus 2003). Additional practices beneficial to turtles are described at:

http://www.dfw.state.or.us/ODFWhtml/springfield/W_Pond_Turtle.htm

Table 4.12.2.2b. Amphibians and reptiles documented as occurring within the study area		
Common Name	Scientific Name	Documentation of local occurrence
Amphibians		
Northwestern Salamander	<i>Ambystoma gracile</i>	1;2;4
Long-Toed Salamander	<i>Ambystoma macrodactylum</i>	1;2;3a;4
Clouded Salamander	<i>Aneides ferreus</i>	1;2;4
Ensatina	<i>Ensatina eschscholtzii</i>	1;2;4; 6

Common Name	Scientific Name	Documentation of local occurrence
Amphibians		
Dunn's Salamander	<i>Plethodon dunni</i>	1;2
Western Red-Backed Salamander	<i>Plethodon vehiculum</i>	1;2;4
Roughskin Newt	<i>Taricha granulosa</i>	1;2;3c; 4; 6
Pacific Giant Salamander	<i>Dicamptodon tenebrosus</i>	2; 4
Southern Torrent Salamander	<i>Rhyacotriton variegatus</i>	1;2
Tailed Frog	<i>Ascaphus truei</i>	5
Pacific Treefrog (Chorus Frog)	<i>Pseudacris regilla</i>	1;2;3a; 4; 6
Red-Legged Frog	<i>Rana aurora</i>	1;2;3u; 4; 6
Bullfrog	<i>Rana catesbeiana</i>	1; 3a; 4; 6
Reptiles		
Painted Turtle	<i>Chrysemys picta</i>	1; 4; 5
Western Pond Turtle	<i>Clemmys marmorata</i>	1;2;3u; 4; 5
Northern Alligator Lizard	<i>Elgaria coerulea</i>	1;2; 6
Southern Alligator Lizard	<i>Elgaria multicarinata</i>	1;2;3r; 4
Western Fence Lizard	<i>Sceloporus occidentalis</i>	1;2;3r; 4; 6
Western Skink	<i>Eumeces skiltonianus</i>	1; 2;4; 6
Rubber Boa	<i>Charina bottae</i>	1; 3c;4
Racer	<i>Coluber constrictor</i>	1;2;3c; 4
Sharptail Snake	<i>Contia tenuis</i>	1;2;3u; 4; 5
Ringneck Snake	<i>Diadophis punctatus</i>	1;2;3a; 4
Gopher Snake	<i>Pituophis catenifer</i>	1;2;3a; 4; 6
Northwestern Garter Snake	<i>Thamnophis ordinoides</i>	1;2;3c; 4
Common Garter Snake	<i>Thamnophis sirtalis</i>	1;2;3a; 4; 6
<p>NOTE 1: Western Rattlesnake (<i>Crotalus viridis</i>) was formerly reported at Coffin Butte and west of Dallas but no verified records exist for recent years.</p> <p>NOTE 2: Western Terrestrial Garter Snake (<i>Thamnophis elegans</i>) has been reported rarely from the adjoining Yamhill watershed in habitat similar to that existing in the Luckiamute.</p> <p>*1= documented in Polk Co. by Nussbaum et al. 1983; 0= not documented in Nussbaum et al. 1983</p> <p>2= documented by St. John (1987) in Luckiamute watershed</p> <p>3= on E.E. Wilson Wildlife Area list, based on reports by Russell Oates, John Crawford, Richard Hoyer, and/or David Budeau (A= abundant, C= common, U= uncommon, R= rare)</p> <p>4= on Willamette Valley National Wildlife Refuges list (not necessarily in Luckiamute)</p> <p>5= Oregon Natural Heritage Program database</p> <p>6= Henny et al. 1999 (Camp Adair Military Training Area)</p>		

4.12.2.3 Birds

Birds not only contribute to the Luckiamute Watershed’s biodiversity, but also help reduce pest insects and maintain healthy aquatic and terrestrial ecosystems. Several species are legally hunted, and a few damage crops. All provide recreation and aesthetic enjoyment to birders and the public generally.

Data Availability

Although no watershed-wide surveys have been undertaken of birds, more data are available for birds than for other wildlife groups. e.g.

Breeding Bird Atlas

Breeding Bird Survey route
OBOL and BirdNotes archives
Winter Waterfowl Surveys
Christmas Bird Counts: Dallas, Airlie-Albany
Baskett Slough & E.E. Wilson & Camp Adair
Hagar & Stern 2001, theses
MacForest
Camp Adair
Boise

Species Status, Trends, and Geographic Patterns

A total of 232 bird species have been documented within or near the Luckiamute watershed (Appendix F: Birds). Of these, 131 (56%) breed somewhat regularly, 87 (38%) are year-round residents (although numbers may change greatly seasonally), 20 (9%) have been recorded only as migrants, and 25 (11%) have been recorded only in winter. Based on percent of watershed comprised of “optimal” habitat for the species, the Luckiamute and/or Rickreall watersheds rank in the top 10% of Willamette watersheds (among 72 fifth-field HUC watersheds) for the following species, according to a regional analysis conducted by Adamus *et al.* (*in press*): Western pond turtle, American beaver, mink, California ground squirrel, Camas pocket gopher, coast mole, Townsend’s mole, White-tailed Kite, Wild Turkey, N. Rough-winged Swallow, Grasshopper Sparrow, Savannah Sparrow, Vesper Sparrow, and Western Meadowlark. In addition, based on percent of watershed comprised of “good” habitat for the species, the two watersheds rank in the top 10% of Willamette watersheds for the following species: Pacific treefrog, ring-necked snake, Spotted Sandpiper, Horned Lark, Common Yellowthroat, and Red-winged Blackbird.

Bird species that likely were common in the vicinity of the watershed around the time of pioneer settlement, but now absent or nearly so, include Snow Goose, Greater White-fronted Goose, California Condor, Sandhill Crane, Long-billed Curlew, and Yellow-billed Cuckoo (Taft and Haig, 2003). In addition, actual numbers of herons, egrets, geese, ducks, swans, and shorebirds were much greater than presently. Declines in many other species (and increases in others) from their pre-settlement abundance levels can be presumed based on documented changes in land cover types and patterns, but documentation is lacking.

As we did with amphibians and reptiles, we used a modeling approach to tentatively identify patterns of “weighted species richness” (WSR) for birds (Figure -xx). Areas with higher WSR scores (colored orange or red on the maps) can be assumed to individually be providing better habitat to a larger number of bird species. In general, western parts of the watershed have greater WSR, as do riparian areas, and the WSR of the Luckiamute watershed is mostly greater than that of the Rickreall (Ash Creek) watershed. Modeling procedures and limitations were described in the Amphibians and Reptiles section.

Of particular concern are species that are rare or believed to have declined severely or disappeared entirely from the region in recent years. The ORNHIC database includes seven such species that have been reported specifically from the Luckiamute Watershed (Table xx). It is possible that other rare or declining birds are present in the watershed but are not included in this table because of the lack of a survey covering all watershed lands.

Table 4.12.2.3a. ORNHIC database records of notable bird species reported from the Luckiamute Watershed			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Branta canadensis leucopareia</i>	Aleutian Canada Goose	G5T3 S2N; 1 Fed= LE ODFW= LT	open land
<i>Oreortyx pictus</i>	Mountain Quail	G5 S4; 4 Fed= SOC ODFW= SU	foothill forests, clearcuts
<i>Brachyramphus marmoratus</i>	Marbled Murrelet	G3G4 S2; 2 Fed= LE ODFW= LT	old-growth forest and stands
<i>Strix occidentalis caurina</i>	Northern Spotted Owl	G3T3 S3; 1 Fed= LE ODFW= LT	multi-layered mature and old-growth forest
<i>Eremophila alpestris strigata</i>	Streaked Horned Lark	G5T2 S2; 2 Fed= SOC ODFW= SC	sparsely-vegetated fields, prairie, overgrazed pastures
<i>Pooecetes gramineus affinis</i>	Oregon Vesper Sparrow	G5T3 S3B; 2 Fed= SOC ODFW= SC	young tree plantations
<i>Progne subis</i>	Purple Martin	G5 S3B; 2 Fed= SOC ODFW= SC	large snags near water
<p>* LE= legally listed as Endangered; LT= legally listed as Threatened; SOC= Species of Concern; SU= status undetermined; SV= vulnerable; SC= critical. Critical habitat can be protected legally only for LE and LT species. G= global status (ONHP), S= state status (ONHP). 1= critically imperiled; 2= imperiled and vulnerable to extinction; 3= rare but not immediately imperiled, 4= not rare but of long-term concern, 5= widespread and secure Number following the semicolon: 1= extinction threatened or presumed, 2= extirpation from Oregon threatened or presumed, 3= insufficient information, 4= of concern but not immediately imperiled</p>			

A significant number of other rare or declining bird species with special status (according to federal or state agencies) are known to occur in the watershed, at least sporadically, but have not been registered yet in the ORNHIC database:

Table 4.12.2.3b.			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Ammodramus savannarum</i>	Grasshopper Sparrow	G5 S2B; 2 ODFW= P	see Altman (1997)
<i>Chlidonias niger</i>	Black Tern	G4 S3B; 4 Fed= SOC	large permanently-flooded marshes
<i>Chordeiles minor</i>	Common Nighthawk	G5 S5; 4	gravel bars, large clearcuts, young tree

Table 4.1 2.2.3b.			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
		ODFW= C	plantations
<i>Columba fasciata</i>	Band-tailed Pigeon	G5 S4; 4 Fed= SOC	Conifer forests, large cottonwoods. Declining statewide.
<i>Contopus cooperi</i>	Olive-sided Flycatcher	G5 S4; 4 Fed= SOC ODFW=V	Tall open conifer forest. Declining statewide.
<i>Cypseloides niger</i>	Black Swift	G4 S1B ODFW= P	waterfalls
<i>Dryocopus pileatus</i>	Pileated Woodpecker	G5 S4; 4 ODFW=V	woods with large-diameter trees
<i>Empidonax traillii brewsteri</i>	Willow Flycatcher	G5TU SUB; 4 ODFW=V	Riparian shrubs, clearcuts. Declining statewide.
<i>Falco peregrinus anatum</i>	American Peregrine Falcon	G4T3 S1B; 2 Fed= LE	open land, wetlands, cliff faces
<i>Icteria virens</i>	Yellow-breasted Chat	G5 S4; 4 Fed= SOC ODFW=C	brushy fields; see Altman (1997)
<i>Melanerpes formicivorus</i>	Acorn Woodpecker	G5 S3; 4 Fed= SOC	oak woodlands
<i>Melanerpes lewis</i>	Lewis's Woodpecker	G5 S3; 4 Fed= SOC ODFW= C	oak woodlands, riparian areas
<i>Sialia mexicana</i>	Western Bluebird	G5 S4; 4 ODFW=V	uncultivated fields, clearcuts, and pasture near oak woodlands
<i>Sturnella neglecta</i>	Western Meadowlark	G5 S5; 4 ODFW= C	uncultivated fields and pasture with widely spaced shrubs; see Altman (1997). Declining statewide.

Some bird species occurring in the watershed are recognized as “priority species” by the U.S. Bureau of Land Management (BLM 2003). They include (in addition to some of those listed above) Trumpeter Swan, Blue Grouse, Western Screech-Owl, Vaux’s Swift, Rufous Hummingbird, Pacific-slope Flycatcher, Cassin’s Vireo, Hutton’s Vireo, Chestnut-backed Chickadee, Golden-crowned Kinglet, Swainson’s Thrush, Wrenit, Black-throated Gray Warbler, Hermit Warbler, MacGillivray’s Warbler, and Black-headed Grosbeak.

Also, some Luckiamute species may be of particular concern because of statistically-significant declines in their regional or local breeding populations, as suggested by annual Breeding Bird Survey (BBS) data, 1966-2002 (Sauer et al. 2002, see Appendix F). The following declining species have not already been mentioned and meet this criterion: Cinnamon Teal, American Kestrel, Ring-necked Pheasant, Killdeer, Mourning Dove, Western Wood-Pewee, Barn Swallow, Red-breasted Nuthatch, Ruby-crowned Kinglet, Hermit Thrush, Orange-crowned Warbler, Chipping Sparrow, Fox Sparrow, Song Sparrow, White-crowned Sparrow, Brewer’s Blackbird, and both American and Lesser Goldfinches. A comparison of nine oak stands throughout the Willamette Valley, 1967-68 vs. 1994-96, suggested a decrease has occurred in species commonly associated with open-stand oaks, such as Bushtit and Chipping Sparrow (Hagar and Stern 2001). The widely-noted and nearly complete disappearance of Lewis’s Woodpecker from the

Willamette Valley (as documented in part by the Dallas Christmas Bird Count) may be attributed partly to the loss of mature large-diameter oak stands which the species depends on for acorn storage. A preliminary analysis of data from the Dallas Christmas Bird Count (Adamus, unpublished, see Appendix F) also suggests the possibility that wintering populations of the following year-round resident species might be declining significantly in or near parts of the Luckiamute watershed: Ring-necked Pheasant, Northern Flicker, Black-capped Chickadee, White-breasted Nuthatch, American Dipper, Dark-eyed Junco, Brewer's Blackbird, and Evening Grosbeak. In addition, the following visiting species from other regions are now being found in fewer numbers than 30 years ago, in or near the Luckiamute watershed: Northern Pintail, Rough-legged Hawk, Northern Shrike, and Ring-billed Gull.

Past and future decline in the extent and quality of oak stands is causing local declines in many bird species. When oak stands are invaded by Douglas-fir, their avifauna is replaced mostly by species that already occur widely in coniferous forests of the adjoining Oregon coast range. Many oak woodland inhabitants do not regularly use coniferous stands. These include Mourning Dove, Acorn Woodpecker, Downy Woodpecker, White-breasted Nuthatch, Western Wood-Pewee, Bewick's Wren, House Wren, Western Scrub-Jay, Cassin's Vireo, Lazuli Bunting, and American Goldfinch. More species of Neotropical migrants – a bird group of particularly high conservation concern – use oak stands than coniferous forests (Hagar and Stern 2001).

In the Luckiamute watershed, surveys specifically targeting particular legally-listed threatened, endangered, and sensitive species have been published only for the Camp Adair Military Training Area (Schreiber 2002) and Beazell County Park (Schreiber 2001). Using recognized survey techniques in habitat of at least minimal suitability, the surveys at Camp Adair had no breeding-season detections of Spotted Owl, Common Nighthawk, Western Meadowlark, or Horned Lark, but did detect Northern Pygmy-Owl, Pileated Woodpecker, Yellow-breasted Chat, Willow Flycatcher, Olive-sided Flycatcher, and Western Bluebird (Schreiber 2002). The briefer survey at Beazell County Park found Pileated Woodpecker but no Spotted Owl or other sensitive species, although habitat conditions for many seemed promising and further surveys were recommended (Schreiber 2001).

Pest Species

There are no bird species that cause major, widespread economic damage to crops or property in the watershed. Waterfowl and blackbirds cause sporadic damage to crops in some areas. Jays, crows, woodpeckers, and a few other species occasionally damage filbert and fruit orchards. Starlings and robins can inflict damage at vineyards and other places where berries are grown. Minor annoyances are caused by Vaux's swifts that sometimes fly down (and occasionally nest in) chimneys, as well as swallows, starlings, barn owls, and house sparrows that nest in or on barns and other structures. However, all of these species are very useful as controllers of insect or rodent pests. Information on dealing with wildlife damage can be found at:

<http://www.dfw.state.or.us/springfield/wildlifeandpeople.html>

Game Birds

Birds subject to legal hunting as game in the Luckiamute watershed include waterfowl (29 species), quail (2 species), grouse (2 species), wild turkey, pheasant, pigeons (2 species), and mourning dove. Harvest data specific to the watershed are not available. For Polk County as a whole, wintering waterfowl numbers based on aerial surveys are given in Table 4.12.2.3c.

	January 2000	January 2001	January 2002	January 2003
Ducks	36,662	34,297	24,404	24,896
Geese	32,946	30,975	8371	22,701
Swans	105	206	218	205
Total*	69,713	65,478	32,993	47,802

* at Baskett Slough NWR alone, roosting geese numbered 14,204 (in 2001), 16,400 (in 2002), and 26,240 (in 2003)

Important Habitats and Communities

In the Luckiamute watershed, some of the rarest birds are associated with mature conifer forests, oak woodlands, prairies (or uncultivated fields), wetlands, and riparian (especially cottonwood) areas. Within these habitats, large snags are particularly important. Some types of built structures also are important. Barn owls depend on open abandoned buildings and barns and silos near pasture or wetlands. Beneficial swallows and bats use these structures as well as larger buildings and bridges for nesting or roosting. Tall, weathered, unused chimneys provide essential roosting habitat for large congregations of Vaux’s Swift during late summer. Just prior to migrating south, thousands of swallows sometimes depend on large, lowland, unharvested corn fields for roosting and feeding.

Conservation and Restoration Potential

Activities to conserve and restore native plant communities will generally benefit native bird communities. However, when mowing or burning is used to restore native prairies, it should be done after July 15 or before April 15 in order to minimize harm to nesting species. Planting willows and other native woody species along waterways, as an alternative to Himalayan blackberry, will benefit most bird species. Also especially important to birds is the retention of standing and downed dead trees (snags). In areas where large snag densities are fewer than 1 per acre, creation of snags by intentionally killing standing trees, at least those of lower commercial value, is warranted where they do not pose a hazard to humans. This provides longer-term benefits than the construction and deployment of bird boxes, but boxes nonetheless may be worthwhile in areas with few trees, e.g., edges of fields. Where abandoned buildings do not pose a safety hazard, they should be allowed to remain standing for the shelter they sometimes provide to swallows, swifts, and barn owls. Maintaining water sources (small seeps, ponds, natural drainages) is important to all wildlife species, so plans to extract groundwater for

irrigation or other uses should be carefully considered for their potential impacts to near-surface water tables.

4.12.2.4 Mammals

This section covers rodents, canids, ungulates, bats, and other wild mammals. Species in these groups not only contribute to the Luckiamute Watershed’s biodiversity, but also help reduce pest insects and maintain healthy aquatic and terrestrial ecosystems. Several species are legally hunted, and some cause damage to crops and livestock.

Data Availability

No comprehensive surveys have been undertaken of mammal diversity in the Luckiamute watershed. Limited surveys of mammals have been conducted at specific locations in the watershed, such as the Camp Adair Military Training Area west of Coffin Butte (Henny et al. 1999) and in McDonald-Dunn Forest. Also, reports covering larger regions that include the Luckiamute have been published (e.g., Verts & Carroway 1998), and anecdotal observations of mammals have been recorded at the E.E. Wilson Wildlife Area and a few other locations.

Species Status and Geographic Patterns

A total of 69 mammal species are known or likely to occur within the Luckiamute Watershed (Appendix G: Mammals).

As we did with birds, amphibians, and reptiles, we used a modeling approach to tentatively identify patterns of “weighted species richness” (WSR) for mammals (Figure xx). Areas with higher WSR scores (colored orange or red on the accompanying maps) can be assumed to individually be providing better habitat to a larger number of mammal species. In general, western parts of the watershed have greater WSR, as do riparian areas, and the WSR of the Luckiamute watershed is mostly greater than that of the Rickreall (Ash Creek) watershed. Modeling procedures and limitations were described in the Amphibians and Reptiles section.

Of particular concern are species that are rare or believed to have declined severely or disappeared entirely from the region in recent years. The ORNHIC database does not include any records of such species specifically from the Luckiamute Watershed. However, the following mammal species that could potentially occur (or are known to occur) in the watershed would be of particular note:

Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Sciurus griseus</i>	western gray squirrel	G5 S4; 3 ODFW= SU	oak woodlands; residential areas
<i>Thomomys bulbivorus</i>	Camas pocket gopher	G3G4 S3S4; 3 Fed= SOC	open land
<i>Antrozous pallidus pacificus</i>	Pacific pallid bat	G5T3T4 S3; 2 Fed= SOC ODFW= SV	ranges widely over areas with large sustained outputs of flying insects

Table 4.12.2.4.			
Scientific Name	Common Name	Listing Status*	Habitat/ Comments
<i>Arborimus (Phenacomys) albipes</i>	white-footed vole	G3G4 S3; 4 Fed= SOC ODFW= SU	riparian deciduous shrubs in conifer forest
<i>Corynorhinus (Plecotus) townsendii townsendii</i>	Pacific western big-eared bat	G3T3T4 S2; 2 Fed= SOC ODFW= SC	ranges widely over areas with large sustained outputs of flying insects, and caves or abandoned buildings for roosting
<i>Lasionycteris noctivagans</i>	silver-haired bat	G5 S4; 4 Fed= SOC ODFW= SU	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Lasiurus cinereus</i>	hoary bat	G5 S4; 4	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Myotis evotis</i>	long-eared myotis (bat)	G5 S3; 2 Fed= SOC ODFW= SU	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Myotis thysanodes</i>	fringed myotis (bat)	G4G5 S2; 2 Fed= SOC ODFW= SV	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Myotis volans</i>	long-legged myotis (bat)	G5 S3; 4 Fed= SOC ODFW= SU	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Myotis yumanensis</i>	Yuma myotis (bat)	G5 S3; 4 Fed= SOC	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<i>Tadarida brasiliensis</i>	Brazilian free-tailed bat	G5 S2; 2	ranges widely over areas with large sustained outputs of flying insects and caves or abandoned buildings for roosting
<p>* SOC= Species of Concern; SU= status undetermined; SV= vulnerable; SC= critical. These state (ODFW) and federal (Fed) designations are advisory only and confer no additional legal protection to the species. G= global status (ONHP), S= state status (ONHP). 1= critically imperiled; 2= imperiled and vulnerable to extinction; 3= rare but not immediately imperiled, 4= not rare but of long-term concern, 5= widespread and secure Number following the semicolon: 1= extinction threatened or presumed, 2= extirpation from Oregon threatened or presumed, 3= insufficient information, 4= of concern but not immediately imperiled</p>			

Game and Furbearer Species Species in the Luckiamute watershed that can be trapped commercially for their fur are muskrat, nutria, mink, river otter, beaver, bobcat, fox (2 species), coyote, raccoon, and opossum. Mammals harvested as game include black bear, elk, deer, cougar, squirrels (3 species), and rabbits (4 species).

Pest Species

Several mammals cause (or have the potential to cause) widespread economic damage or damage to habitat of other species. Deer, elk, bear, opossum and rodents frequently cause damage to crops, tree plantations, landscaping shrubs, and/or gardens. Beaver plug culverts and in doing so, flood property. Nutria alter wetland vegetation structure and thus potentially affect habitat quality for many species. Feral cats prey on many native mammals and birds.

Important Habitats and Communities

In the Luckiamute watershed, some of the rarest mammals are associated with mature conifer or oak woodland, and riparian areas. Within these habitats, areas with extensive downed wood and large standing stags are particularly important.

Conservation and Restoration Potential

Activities to conserve and restore native plant communities will generally benefit native mammals. Planting willows and other native woody species along waterways, as an alternative to Himalayan blackberry, will benefit many mammal species. However, in some instances new plantings may be removed by beaver. Just as important to mammals as to birds are adequate densities of snags (for roosting bats) and downed wood (used by numerous rodents). In areas where large snag densities are fewer than 1 per acre, creation of snags by intentionally killing standing trees, at least those of lower commercial value, is warranted where they do not pose a hazard to humans. Where natural snags are few (due to early successional stage of the forest), installation of boxes designed for bats is warranted. Bats using buildings and other structures should be accommodated whenever possible and not harrassed. Maintaining water sources (small seeps, ponds, natural drainages) is important to all wildlife species, so plans to extract groundwater for irrigation or other uses should be carefully considered for their potential impacts to near-surface water tables.

Table xx. Mammals documented as occurring (or potentially occurring) within the study area		
<i>Common Name</i>	<i>Scientific Name</i>	<i>Documentation of local occurrence</i>
American Beaver	<i>Castor canadensis</i>	3c;4;6
Virginia Opossum	<i>Didelphis virginiana</i>	1;2;3c;4;6
Vagrant Shrew	<i>Sorex vagrans</i>	1;2;3a;4;6
Pacific Shrew	<i>Sorex pacificus</i>	2;6
Pacific Water Shrew	<i>Sorex bendirii</i>	1
Trowbridge's Shrew	<i>Sorex trowbridgii</i>	1;2;3u;4;6
Baird's Shrew	<i>Sorex bairdi</i>	2
Fog Shrew	<i>Sorex sonomae</i>	1;2
Shrew-Mole	<i>Neurotrichus gibbsii</i>	1;2;4
Townsend's Mole	<i>Scapanus townsendii</i>	1;3c;4;6
Coast Mole	<i>Scapanus orarius</i>	1;2;4
Little Brown Myotis	<i>Myotis lucifugus</i>	4;6
Yuma Myotis	<i>Myotis yumanensis</i>	1;6
Long-Eared Myotis	<i>Myotis evotis</i>	1;4;6
Fringed Myotis	<i>Myotis thysanodes</i>	0
Long-Legged Myotis	<i>Myotis volans</i>	6
California Myotis	<i>Myotis californicus</i>	4;6

<i>Common Name</i>	<i>Scientific Name</i>	<i>Documentation of local occurrence</i>
Silver-Haired Bat	<i>Lasionycteris noctivagans</i>	1;4
Big Brown Bat	<i>Eptesicus fuscus</i>	1;4;6
Hoary Bat	<i>Lasiurus cinereus</i>	0;4
Townsend's Big-Eared Bat	<i>Corynorhinus townsendii</i>	0
Pallid Bat	<i>Antrozous pallidus</i>	0
Brazilian Free-Tailed Bat	<i>Tadarida brasiliensis</i>	0
Brush Rabbit	<i>Sylvilagus bachmani</i>	1;2;3c;4;6
Eastern Cottontail	<i>Sylvilagus floridanus</i>	1;3a;4;6
Snowshoe Hare	<i>Lepus americanus</i>	0
Black-Tailed Jackrabbit	<i>Lepus californicus</i>	1;3r;4;6
Mountain Beaver	<i>Aplodontia rufa</i>	0
Townsend's Chipmunk	<i>Tamias townsendii</i>	1;2;3u;4;6
California Ground Squirrel	<i>Spermophilus beecheyi</i>	1;2;3a;4;6
Western Gray Squirrel	<i>Sciurus griseus</i>	1;3u;4;6;7
Douglas' Squirrel	<i>Tamiasciurus douglasii</i>	1;2;4;6
Northern Flying Squirrel	<i>Glaucomys sabrinus</i>	1;2;4;6
Western Pocket Gopher	<i>Thomomys mazama</i>	1
Camas Pocket Gopher	<i>Thomomys bulbivorus</i>	3a;4;5;6
Deer Mouse	<i>Peromyscus maniculatus</i>	1;2;3a;4;6
Dusky-Footed Woodrat	<i>Neotoma fuscipes</i>	1;2;3c;4;6
Bushy-Tailed Woodrat	<i>Neotoma cinerea</i>	1;2;4
Western Red-Backed Vole	<i>Clethrionomys californicus</i>	1;2;4
White-Footed Vole	<i>Phenacomys albipes</i>	0
Red Tree Vole	<i>Phenacomys longicaudus</i>	1;4
California Vole	<i>Microtus californicus</i>	0
Townsend's Vole	<i>Microtus townsendii</i>	1;2;3a;4;6
Long-Tailed Vole	<i>Microtus longicaudus</i>	1
Creeping (Oregon) Vole	<i>Microtus oregoni</i>	1;2;3c;4;6
Gray-Tailed Vole	<i>Microtus canicaudus</i>	1;3a;4;6
Muskrat	<i>Ondatra zibethicus</i>	3c;4
Black Rat	<i>Rattus rattus</i>	4
Norway Rat	<i>Rattus norvegicus</i>	1;3c
House Mouse	<i>Mus musculus</i>	1;3c;4
Pacific Jumping Mouse	<i>Zapus trinotatus</i>	1;2;3u;4;6
Common Porcupine	<i>Erethizon dorsatum</i>	3r;4
Nutria	<i>Myocastor coypus</i>	3c;4
Coyote	<i>Canis latrans</i>	1;3c;4;6
Red Fox	<i>Vulpes vulpes</i>	3u;4
Gray Fox	<i>Urocyon cinereoargenteus</i>	1;3r;4
Black Bear	<i>Ursus americanus</i>	1;3r;4
Raccoon	<i>Procyon lotor</i>	1;3c;4;6;7
Ermine	<i>Mustela erminea</i>	1;2;3r;4
Long-Tailed Weasel	<i>Mustela frenata</i>	1;4;7
Mink	<i>Mustela vison</i>	3u;4
Western Spotted Skunk	<i>Spilogale gracilis</i>	4
Striped Skunk	<i>Mephitis mephitis</i>	3c;4;6
Northern River Otter	<i>Lutra canadensis</i>	3u;4
Mountain Lion	<i>Felis concolor</i>	3r;4;7
Feral House Cat	<i>Felis catus</i>	4;6

Table xx. Mammals documented as occurring (or potentially occurring) within the study area

<i>Common Name</i>	Scientific Name	Documentation of local occurrence
Bobcat	<i>Lynx rufus</i>	3u;4;6
Elk	<i>Cervus elaphus</i>	3r;4;6;7
Black-Tailed Deer	<i>Odocoileus hemionus</i>	1;3c;4;6;7

*0= is potentially present but no documented records
 1= documented in Polk Co. by Verts & Carraway 1998
 2= documented in McDonald-Dunn State Forest by Dave Waldien (pers. comm.)
 3= on E.E. Wilson Wildlife Area list, based on reports by Russell Oates, John Crawford, Richard Hoyer, and/or David Budeau (A= abundant, C= common, U= uncommon, R= rare)
 4= on Willamette Valley National Wildlife Refuges list (not necessarily in Luckiamute)
 6= Henny et al. 1999 (Camp Adair Military Training Area)

4.13 EXISTING WATERSHED ANALYSES

U.S. Bureau of Land Management

In 1998 the BLM conducted a watershed assessment of the Rowell Creek/Mill Creek/Rickreall Creek/Luckiamute River Watershed (Licata *et al.*, 1998).

Western Oregon University

The biology and earth and physical sciences departments of Western Oregon University created a manual for environmental educators to use in studying the Luckiamute River watershed (Taylor *et al.*, 2003).

Rickreall Watershed Assessment

The Rickreall Watershed Council completed a watershed assessment for the Rickreall Creek watershed in January 2001. A small portion of the land examined in the Rickreall Watershed Assessment is included in the northeast corner of the present Luckiamute Watershed Assessment area (Mattson and Gallagher, 2001).

Midcoast Watershed Assessment

This assessment, completed by Earth Design Consultants, Inc. and Green point Consulting, covers a portion of Polk and Benton Counties (Garono and Brophy, 2001). This report is available online at www.midcoastwatershedcouncil.org.

Yamhill Watershed Assessment

The Yamhill River Basin contains eight sub-watersheds: Willamina, North Yamhill River, Chehalem Creek and South Yamhill River. All of these sub-basins are contained primarily in Yamhill and Polk Counties. For more information visit

<http://www.co.yamhill.or.us/ybc/assessments.htm>

4.14 EXISTING PRESERVES AND WATERSHED RESTORATION PROJECTS

E.E. Wilson Wetlands, ODFW <http://wetlands.dfw.state.or.us/projects/willamette.html>

These wetlands are on land managed by the Oregon Department of Fish and Wildlife (ODFW). ODFW began restoring wetlands here in 1992. By 1995 several dozen small ponds and wetlands were constructed or enhanced, totaling 170 acres of improvements. The area is still managed to control non-native vegetation, especially reed canary grass.

Fort Hoskins, Benton County http://www.co.benton.or.us/parks/fh_final_plan.htm

The land around the historic Fort Hoskins site (1856 – 1865) is now a park owned by Benton County. The County’s goals for the property include 1) active restoration of pre-settlement native oak savanna, 2) sustainable forest management using environmentally sensitive techniques, and 3) creating reserve areas to protect unique resources. Promoting recreation activities and public education are also major goals for projects in the park.

Luckiamute Landing State Park

http://wetlands.dfw.state.or.us/pdfs/willamette_valley_draft.pdf

This restoration project is in an area just north of Albany, at the confluence of the Luckiamute, Santiam and Willamette Rivers. The Western Rivers Conservancy and State Parks have proposed to acquire and protect a large block of floodplain habitats around Luckiamute Landing.

In addition to these preserved lands, there are also numerous small preservation and restoration projects going on within the basin. We checked the REO web site for information on existing watershed restoration projects. We found one restoration site from this database: It was listed as a culvert upgrade in the headwaters of the Luckiamute (IRDS_ID: 444729612331517). The Greenbelt Land Trust, a volunteer non-profit organization in the mid-Willamette Valley, involved in the protection of open space, has organized and compiled a list of regional restoration projects. Table 4.14 lists the restoration sites compiled by the Greenbelt Land Trust. It is not clear whether some of these sites are being routinely monitored or not. We highly recommend incorporating these sites into the LWC monitoring plan.

Table 4.14. List of restoration sites compiled by the Greenbelt Land Trust. Shown are watershed ID code, type of restoration, habitat affected, a brief description, partners, whether the site is being monitored or not and the number of acres.

HUC	TYPE	HABITAT	DESCRIPTION	PARTNERS	MON	ACRES
3060704				ODFW		41.0
3060704				ODFW		5.0
3060901		wetlands		ODFW		5.0
3060901				ODFW, S*, owner		6.0
3060901		woodlands		ODFW, B*, owner		12.0
3060704		woodlands		ODFW, K*, owner		9.0
3060704		oak woodlands		ODFW, D*, owner		20.0
3060802				ODFW, R*, owner		1.0

Table 4.14. List of restoration sites compiled by the Greenbelt Land Trust. Shown are watershed ID code, type of restoration, habitat affected, a brief description, partners, whether the site is being monitored or not and the number of acres.

HUC	TYPE	HABITAT	DESCRIPTION	PARTNERS	MON	ACRES
3061004				ODFW, P*, owner		9.0
3060503				ODFW, W*, owner		22.0
3061002	wetland restoration	wetlands		USFWS, R*, owner	y	15.0
3061002	oak restoration	oak woodlands		USFWS, R*, owner	y	0.0
3061002	wetland restoration	wetland, oak		USFWS, J*, owner		20.0
3061004	oak restoration	oak woodlands		USFWS, J*, owner		0.0
3061004	wetland restoration	wetland		ODFW, R*, owner, USFWS	y	25.0
3061005	wetland restoration	wetland		ODFW, S*, owner, NRCS		40.0
3061005	easement	riparian, upland		ODFW, B*, owner		60.0
3061005	wetland restoration	wetland		ODFW, K*, owner	y	40.0
3061303	wetland restoration	riparian, wetland		NAWCA state park, Vanderpool, USFWS	y	230.0
3061302	wetland restoration	wetland		ODFW, USFWS, G*, owner	y	15.0
3061303	wetland restoration	wetland		ODFW, USFWS, U*, owner	y	20.0
3061204	wetland restoration	wetland		ODFW, USFWS, O*, owner	y	76.0
3061302	wetland restoration	wetlands		ODFW, OWEB, Winter Creek	y	52.0
3061302	riparian	riparian		ODFW, OWEB, Winter Creek	y	0.0
3060204				Benton County		0.0
3060301	wetland restoration	wetlands		ODFW, USFWS, E*, owner, OWEB	y	120.0
3060301	oak restoration	oak savanna		ODFW, USFWS, E*, owner, OWEB	y	0.0
3061101	easement			Greenbelt Land Trust	yes	0.0
3061005	riparian		riparian planting	OPRD, ODFW, farmer		0.0
3061103	easement	wetlands		NRCS, owner		58.0
3060703	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.0
3060701	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.6
3060701	riparian	forest	riparian tree	Willamette	no	4.1

Table 4.14. List of restoration sites compiled by the Greenbelt Land Trust. Shown are watershed ID code, type of restoration, habitat affected, a brief description, partners, whether the site is being monitored or not and the number of acres.

HUC	TYPE	HABITAT	DESCRIPTION	PARTNERS	MON	ACRES
			retention	Industries, Inc.		
3060701	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	6.1
3060701	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	4.6
3060701	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	34.0
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.0
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.0
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.0
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	4.5
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.9
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.7
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	4.1
3060603	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.2
3060601	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.9
3060601	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.2
3060601	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.0
3060601	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.3
3060601	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	4.0
3060801	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.7
3060802	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.8
3060803	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.7
3060802	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.3
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	4.6
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.5
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.5

Table 4.14. List of restoration sites compiled by the Greenbelt Land Trust. Shown are watershed ID code, type of restoration, habitat affected, a brief description, partners, whether the site is being monitored or not and the number of acres.

HUC	TYPE	HABITAT	DESCRIPTION	PARTNERS	MON	ACRES
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.2
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.2
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	6.3
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.1
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.5
3060902	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	3.9
3060503	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.2
3060503	riparian	forest	riparian tree retention	Starker Forests, Inc.	no	1.0
3060503	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.5
3060503	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.0
3060501	riparian	forest	riparian tree retention	Boise Cascade	no	1.4
3060501	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	8.4
3060501	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.1
3060501	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	3.2
3060501	riparian	forest	riparian tree retention	Starker Forests, Inc.	no	2.0
3060501	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.9
3060402	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	0.7
3060401	riparian	forest	riparian tree retention	boise	no	4.2
3060404	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	1.2
3060204	riparian	forest	riparian tree retention	Starker Forests, Inc.	no	1.1
3060204	riparian	forest	riparian tree retention	Starker Forests, Inc.	no	0.6
3061201	riparian	forest	riparian tree retention	Boise Cascade	no	4.0
3061102	riparian	forest	riparian tree retention	Willamette Industries, Inc.	no	2.0
3061101	riparian	forest	riparian tree	Starker Forests,	no	0.5

Table 4.14. List of restoration sites compiled by the Greenbelt Land Trust. Shown are watershed ID code, type of restoration, habitat affected, a brief description, partners, whether the site is being monitored or not and the number of acres.

HUC	TYPE	HABITAT	DESCRIPTION	PARTNERS	MON	ACRES
			retention	Inc.		
3061101	riparian	forest	riparian tree retention	Starker Forests, Inc.	no	2.0

5 LAND COVER ANALYSIS

5.1 TERRESTRIAL ANALYSES

5.1.1 FORESTED AND NON-FORESTED AREAS

Using GIS, we combined cover classes from the pre-settlement vegetation and current vegetation data layers (see Section 4.4.1) and calculated the change that has occurred in the spatial extent of these major cover classes (TABLE 5.1.1).

Table 5.1.1: Change in major vegetation cover classes from pre-Settlement to present day. Shown are acres and percent change.

	Pre-Settlement Cover	Current Cover	Change (Acres)	Percent Change
Forest (Unspecified or Mixed)	17,306.1	42,550.2	25,244.1	10.6%
Herbaceous	53,674.9	70,632.7	16,957.8	7.1%
Urban	0.0	8,210.5	8,210.5	3.4%
Forest (Deciduous)	13,821.1	19,606.7	5,785.6	2.4%
Forest (Conifer)	76,002.5	76,153.3	150.8	0.1%
Water	1,123.4	1,107.9	-15.5	0.0%
Shrub	76,114.6	18,185.4	-57,929.2	-24.3%

The proportion of area covered by many of the cover types is surprisingly similar to pre-settlement areas. The largest changes are in the shrub category (a large amount of wetland shrub communities existed at the time of settlement: see Section 4.4) and in the mixed forested classes. Judging from the results presented in Table 5.1.1, shrubby areas have been replaced by all of the other cover classes (which experienced net increases). Of course, the distribution of these major cover classes have changed on the landscape even though the total areas remained similar (see Maps 4 and 5). The fragmented modern landscape has profound implications for wildlife.

Not only has the spatial extent of forest changed, but so has its composition and structure. Logging in the past 150 years has manipulated the vegetation towards the younger seral stages (less than 80 years-old: see Section 4.4.4). Late-seral and old growth forests once

occupied 60 to 80% of the Coast Range landscape [Agee 1993, in (Licata *et al.*, 1998)], In contrast, 96% of the Luckiamute and Rickreall watersheds are currently in early and mid-seral stages (Licata *et al.*, 1998).

5.1.2 OAK SAVANNAH

According to our summary of historic landcover, oak savannah (as part of a deciduous forest), covered about 7% of the Luckiamute/ Ash Creek study area (see Section 4.4.1.1). The Yamhill Basin Council (2001) recognized that there is a difference between Oak Savannah and Oak forests (see Section 4.4.1.1); using the existing data, it is impossible to determine how much of the 7% of the deciduous forest was actually oak savannah. Nevertheless, we do know that modern oak stands have become increasingly dominated by Douglas-fir in many parts of the watershed. This may be a consequence of fire suppression. Also, in the absence of fire many oak stands have become crowded, leading to slower growth rates, stunting, smaller diameters, and fewer tree cavities essential to several nesting birds and mammals (see Sections 4.1.2.23 and 4.1.2.2.4). A survey of nine Willamette Valley oak stands (not just in the Luckiamute watershed) found two-thirds had measurable Douglas-fir in the canopy and subcanopy (Hagar and Stern 2001).

The LWC members recognized the importance of oak savannahs and expressed an interest in mapping the current extent of Oak savannah within the study area. We were unable to find a data source describing the extent of this type of habitat. After reviewing a number of data sets, we presented the council with a series of maps that should help them to develop their own data set. We recommend that the locations on the maps be visited and observations recorded and that oak forests be separated from 'true' oak savannahs. These observations can be use to refine the GIS data sets which describe the extent of oak savannahs.

5.2 WETLAND ANALYSES

As previously mentioned, there are no good data sets describing the spatial extent of wetlands for the entire study area. National Wetland Inventory maps exist for portions of the study area and we used land cover and hydric soils to describe the current distribution of 'potential' wetlands.

Wetland conservation and restoration will most likely be an important component of the LWC action planning process. Since it was not the intention of this analysis to select individual sites for consideration (see Section 2), we recommend that when it comes time to prioritize sites, the riparian, soils, and wildlife WSR grids be used to prioritize sites.

5.3 RIPARIAN ANALYSES

We were unable to find a data set describing the current riparian conditions with in the Luckiamute/ Ash Creek study area. Information on riparian condition can be used to for planning monitoring and restoration actions. Therefore, at the request of LWC we created

a GIS data layer describing 11 cover classes within the riparian zone. We defined the riparian zone as extending 100ft on either side of the streams. In this case, we used the 1:24,000 streams layer. In GIS, we use the digital orthoquads (black and white photographs) to identify and outline areas belonging to the following cover classes: bare areas having little or no vegetation; developed areas with obvious buildings, structures or parking lots; coniferous forests both dense and sparse categories; deciduous forests both dense and sparse categories; mixed forests (not readily identifiable as coniferous or deciduous) both dense and sparse categories; treed-strips areas with a narrow strip of large trees along the river (may provide shade) but predominantly non-forested; and water.

All cover classes were subjectively interpreted from the photographs (Fig. 5.2.1). Due to time constraints, no assessment of classification accuracy was made. We recommend that LWC ground truth the riparian vegetation layer created in this assessment.

In general, vegetation in the riparian areas mirrors patterns in land cover in the rest of the watersheds (see Section 4.4.1). The Luckiamute watershed is dominated by dense conifer forests (43.5% of the riparian zone) and dense deciduous forests (19.4% of the riparian zone). About 18.4% of the riparian zone is herbaceous land cover indicating that there may be ample opportunity for stream side planting wherever appropriate. In contrast, the Ash Creek watershed is dominated by herbaceous cover (45.3% of the riparian zone) and only 13.5% and 12.9% of the riparian zone is covered with dense coniferous and deciduous forests, respectively.

We then examined the condition of the riparian zone within each 7th field watershed by tallying the area covered by each of the cover classes. Cover data are expressed as a percent of the total riparian zone area for each 7th field watershed (Table 5.3b).




Figure 5.2.1. Example of how riparian areas were interpreted from black and white DOQs. Top shows 100ft buffer strip. Bottom shows areas of different cover classes: dark green is dense deciduous forest, yellow is herbaceous, blue is sparse coniferous forest, and pink is sparse mixed forest (see text for details).

Table 5.3b. Land cover within the riparian zone for each of the 7th field watersheds in the study area. Shown are the % of the total riparian zone by cover class for each 7th field watershed.

7th Field HUC	Name	Bare	Developed	Forest						herbaceous	treed-strip	water
				Conifer		Deciduous		Mixed				
				Dense	Sparse	Dense	Sparse	Dense	Sparse			
3060101	Upper Luckiamute	0.0%	0.0%	69.0%	6.2%	3.1%	0.0%	16.3%	0.0%	5.4%	0.0%	0.0%
3060102	Miller Creek	0.0%	0.0%	72.8%	12.0%	9.6%	0.0%	0.0%	0.0%	5.6%	0.0%	0.0%
3060201	Wolf Creek	0.0%	0.0%	74.2%	2.3%	12.0%	0.6%	0.0%	0.0%	11.0%	0.0%	0.0%
3060202	Cougar Creek	0.0%	7.2%	70.7%	0.4%	11.6%	0.2%	0.0%	0.0%	9.8%	0.0%	0.0%
3060203	Hoskins	0.0%	5.4%	43.5%	3.1%	21.4%	4.4%	2.5%	0.8%	17.7%	0.8%	0.3%
3060204	Vincent Creek	0.0%	0.0%	61.2%	15.3%	10.2%	1.5%	0.0%	0.0%	11.4%	0.4%	0.0%
3060301	Plunkett Creek	0.0%	0.6%	25.8%	5.2%	27.2%	4.6%	15.3%	0.4%	12.5%	8.5%	0.0%
3060302	Woods Creek	0.0%	0.0%	16.6%	5.7%	33.3%	2.3%	15.2%	0.0%	20.5%	6.0%	0.4%
3060303	Price Creek	0.0%	0.0%	18.9%	40.7%	19.6%	1.1%	0.0%	1.9%	17.2%	0.6%	0.0%
3060304	Maxfield Creek	0.0%	0.0%	25.1%	25.4%	22.2%	4.6%	1.4%	1.0%	12.1%	7.5%	0.6%
3060305	Bump Creek	0.0%	0.2%	62.8%	8.0%	15.2%	0.4%	7.2%	0.0%	3.5%	2.6%	0.1%
3060401	Lower Pedee Creek	0.0%	0.0%	52.3%	6.9%	21.2%	3.1%	1.8%	1.3%	12.4%	1.0%	0.0%
3060402	Upper Pedee Creek	0.6%	0.0%	87.5%	0.1%	3.8%	0.0%	2.7%	0.0%	4.7%	0.6%	0.1%
3060403	Clayton Creek	0.0%	0.0%	90.5%	1.1%	4.2%	0.7%	0.0%	0.0%	3.4%	0.0%	0.0%
3060404	Ritner Creek	0.0%	0.2%	71.9%	8.0%	10.3%	0.8%	0.0%	1.6%	6.8%	0.5%	0.0%
3060501	Ira Hooker	0.0%	0.0%	41.3%	2.8%	31.4%	8.4%	2.3%	0.1%	13.9%	0.0%	0.0%
3060502	McTimmonds	0.0%	0.3%	41.4%	16.5%	11.4%	0.6%	2.1%	1.8%	22.4%	3.5%	0.0%
3060503	Middle Luckiamute	0.0%	0.1%	17.4%	7.6%	38.0%	2.3%	0.0%	0.0%	27.5%	6.9%	0.1%
3060504	Jont Creek	0.0%	0.0%	33.7%	7.0%	16.6%	6.3%	6.7%	1.9%	25.9%	1.9%	0.1%
3060601	Upper Little Luckiamute	0.1%	0.0%	80.3%	9.9%	0.0%	0.1%	1.3%	0.0%	8.3%	0.0%	0.0%
3060602	Cold Springs	0.0%	0.0%	68.3%	13.4%	8.9%	4.3%	0.4%	0.0%	4.8%	0.0%	0.0%
3060603	Black Rock Creek	0.0%	0.0%	56.1%	10.1%	7.3%	1.1%	0.4%	0.5%	24.4%	0.0%	0.0%

Table 5.3b. Land cover within the riparian zone for each of the 7th field watersheds in the study area. Shown are the % of the total riparian zone by cover class for each 7th field watershed.

7th Field HUC	Name	Bare	Developed	Forest						herbaceous	treed-strip	water
				Conifer		Deciduous		Mixed				
				Dense	Sparse	Dense	Sparse	Dense	Sparse			
3060701	Socialist Valley	0.0%	0.1%	65.0%	3.1%	10.9%	0.0%	0.0%	0.0%	18.5%	2.5%	0.0%
3060702	Falls City	0.0%	0.0%	35.5%	2.4%	35.5%	13.8%	1.7%	0.0%	5.8%	5.2%	0.0%
3060703	Waymire Creek	0.0%	0.0%	57.5%	2.7%	8.5%	15.1%	0.0%	0.0%	9.7%	6.6%	0.0%
3060704	Bridgeport	0.5%	0.1%	28.5%	9.2%	13.8%	11.0%	6.3%	3.3%	15.8%	11.0%	0.4%
3060801	Upper Teal Creek	0.2%	0.0%	62.1%	6.4%	8.4%	3.3%	0.0%	4.0%	15.6%	0.0%	0.0%
3060802	Lower Teal Creek	0.0%	0.5%	41.1%	2.8%	29.1%	3.7%	5.0%	0.0%	16.7%	0.9%	0.1%
3060803	Grant Creek	0.0%	0.0%	47.1%	11.0%	23.5%	0.0%	9.8%	0.0%	8.7%	0.0%	0.0%
3060901	Fern Creek	0.0%	0.2%	21.3%	10.5%	5.9%	8.5%	0.0%	1.2%	40.3%	11.9%	0.2%
3060902	Lower Little Luckiamute	0.0%	0.7%	12.0%	5.3%	41.7%	3.1%	0.6%	0.8%	27.4%	7.3%	1.0%
3060903	Cooper Creek	0.0%	0.0%	11.2%	2.5%	24.8%	6.1%	0.0%	0.3%	37.7%	17.4%	0.0%
3061001	Simpson	0.0%	0.6%	10.5%	1.9%	51.0%	7.2%	0.0%	0.7%	16.7%	11.4%	0.0%
3061002	Zumwalt	0.0%	0.0%	13.7%	1.2%	17.6%	3.9%	0.0%	0.0%	35.6%	28.0%	0.0%
3061003	Helmick	0.0%	1.8%	0.0%	0.0%	73.1%	1.2%	0.0%	0.0%	23.9%	0.0%	0.0%
3061004	Parker	0.0%	1.4%	9.9%	0.0%	9.3%	14.2%	0.0%	0.0%	55.3%	9.9%	0.0%
3061005	Luckiamute Landing	0.0%	0.0%	0.6%	1.8%	71.0%	0.0%	0.0%	0.0%	21.5%	0.0%	5.1%
3061101	Upper Soap Creek	0.0%	0.1%	41.7%	11.4%	17.3%	1.6%	17.3%	0.0%	9.3%	1.3%	0.0%
3061102	Middle Soap Creek	0.0%	0.0%	31.9%	3.5%	33.6%	7.2%	0.0%	0.0%	14.4%	9.5%	0.0%
3061103	Rifle Range	0.0%	0.6%	30.3%	1.3%	30.8%	2.4%	0.0%	0.0%	21.7%	12.9%	0.0%
3061201	Upper Berry Creek	0.0%	0.0%	72.2%	1.7%	8.5%	2.2%	2.9%	0.0%	11.4%	1.1%	0.0%
3061202	Staats Creek	0.0%	0.0%	26.5%	7.0%	8.9%	3.4%	0.0%	0.0%	48.1%	5.9%	0.2%
3061203	Peterson Creek	0.0%	0.0%	11.3%	0.6%	2.1%	4.4%	0.0%	0.0%	78.0%	3.6%	0.0%
3061204	Lower Berry Creek	0.0%	0.1%	9.0%	1.9%	12.3%	5.6%	0.0%	0.0%	66.9%	2.1%	2.2%
3061301	E.E. Wilson	0.0%	0.2%	0.0%	0.0%	83.1%	1.6%	0.0%	0.0%	10.9%	4.3%	0.0%

Table 5.3b. Land cover within the riparian zone for each of the 7th field watersheds in the study area. Shown are the % of the total riparian zone by cover class for each 7th field watershed.

7th Field HUC	Name	Bare	Developed	Forest						herbaceous	treed-strip	water
				Conifer		Deciduous		Mixed				
				Dense	Sparse	Dense	Sparse	Dense	Sparse			
3061302	Palestine	0.0%	1.4%	0.0%	0.0%	24.3%	21.0%	0.0%	0.0%	48.8%	4.5%	0.0%
3061303	Springhill	0.0%	1.7%	1.7%	3.5%	29.6%	12.2%	0.0%	0.0%	42.9%	8.6%	0.0%
7020404	Buena Vista	0.0%	1.3%	0.0%	0.0%	16.8%	0.0%	0.0%	0.0%	78.6%	3.2%	0.2%
7020501	American Bottom	0.0%	0.0%	0.0%	0.0%	50.2%	9.3%	0.0%	0.0%	24.8%	0.0%	15.7%
7020502	Duck Slough	0.0%	0.2%	3.2%	4.2%	9.6%	2.6%	3.2%	0.0%	61.9%	15.2%	0.0%
7020503	Harman Slough	0.0%	0.0%	5.5%	0.0%	11.0%	8.2%	0.0%	0.0%	61.6%	12.8%	0.8%
7020601	Upper North Fork Ash Creek	0.0%	1.4%	16.7%	20.4%	1.2%	8.4%	0.0%	3.2%	25.2%	23.2%	0.4%
7020602	Middle North Fork Ash Creek	0.0%	4.7%	5.7%	3.9%	4.3%	17.9%	0.0%	0.0%	49.5%	13.9%	0.0%
7020603	Lower North Fork Ash Creek	0.0%	4.1%	0.6%	5.0%	30.6%	6.7%	0.0%	0.0%	53.0%	0.0%	0.0%
7020604	Middle Fork Ash Creek	0.0%	0.4%	24.7%	11.1%	14.9%	5.1%	0.0%	0.0%	32.0%	11.7%	0.0%
7020605	Upper South Fork Ash Creek	0.0%	0.1%	24.0%	3.1%	9.3%	3.3%	0.6%	0.0%	51.5%	7.7%	0.4%
7020606	Lower South Fork Ash Creek	0.0%	0.0%	7.5%	0.3%	40.7%	9.2%	3.6%	4.6%	26.8%	4.8%	2.4%

5.3.1 STREAM CHANNEL PERCENT SHADE

Vegetation and landforms provide shade to streams thereby keeping water temperatures cool. We used the riparian GIS layer described above to rank 7th field watersheds by the amount of dense forests (either coniferous or deciduous) in the riparian zones. Table 5.3.1 shows the proportion of dense forest, either coniferous, deciduous, or mixed, in the riparian buffer zone. We calculated the proportion of dense forest by summing the total percentages for each of the dense forest classes from Table 5.2.1 and dividing by 3. Lower ranked watersheds would be those watersheds that may benefit from riparian plantings.

Table 5.3.1. List of 7 th field watersheds ranked according to the amount of dense forest occurring within the riparian zone.		
7th Field HUC	Name	Proportion of Dense Forest
3060403	Clayton Creek	31.6%
3060402	Upper Pedee Creek	31.3%
3060101	Upper Luckiamute	29.4%
3060201	Wolf Creek	28.7%
3060305	Bump Creek	28.4%
3061201	Upper Berry Creek	27.9%
3061301	E.E. Wilson	27.7%
3060102	Miller Creek	27.5%
3060202	Cougar Creek	27.5%
3060404	Ritner Creek	27.4%
3060601	Upper Little Luckiamute	27.2%
3060803	Grant Creek	26.8%
3060602	Cold Springs	25.9%
3061101	Upper Soap Creek	25.4%
3060701	Socialist Valley	25.3%
3060401	Lower Pedee Creek	25.1%
3060802	Lower Teal Creek	25.1%
3060501	Ira Hooker	25.0%
3061003	Helmick	24.4%
3060702	Falls City	24.2%
3061005	Luckiamute Landing	23.9%
3060204	Vincent Creek	23.8%
3060801	Upper Teal Creek	23.5%
3060301	Plunkett Creek	22.8%
3060203	Hoskins	22.5%
3060703	Waymire Creek	22.0%
3061102	Middle Soap Creek	21.8%
3060302	Woods Creek	21.7%
3060603	Black Rock Creek	21.3%

Table 5.3.1. List of 7th field watersheds ranked according to the amount of dense forest occurring within the riparian zone.

7th Field HUC	Name	Proportion of Dense Forest
3061001	Simpson	20.5%
3061103	Rifle Range	20.4%
3060504	Jont Creek	19.0%
3060503	Middle Luckiamute	18.5%
3060502	McTimmonds	18.3%
3060902	Lower Little Luckiamute	18.1%
7020606	Lower South Fork Ash Creek	17.3%
7020501	American Bottom	16.7%
3060304	Maxfield Creek	16.2%
3060704	Bridgeport	16.2%
7020604	Middle Fork Ash Creek	13.2%
3060303	Price Creek	12.8%
3060903	Cooper Creek	12.0%
3061202	Staats Creek	11.8%
7020605	Upper South Fork Ash Creek	11.3%
3061002	Zumwalt	10.4%
3061303	Springhill	10.4%
7020603	Lower North Fork Ash Creek	10.4%
3060901	Fern Creek	9.1%
3061302	Palestine	8.1%
3061204	Lower Berry Creek	7.1%
3061004	Parker	6.4%
7020601	Upper North Fork Ash Creek	6.0%
7020404	Buena Vista	5.6%
7020503	Harman Slough	5.5%
7020502	Duck Slough	5.3%
3061203	Peterson Creek	4.4%
7020602	Middle North Fork Ash Creek	3.4%

5.4 INTERPRETATION AND RECOMMENDATIONS

Land Cover/ Land Use

- Develop a GIS data set that describes Oak Savannah. Develop protocols that separate oak forests from 'true' oak savannahs.

- Develop or obtain up to date land cover information that reflects current conditions in the watershed.
- Ground-truth and update riparian vegetation information, especially in areas known to have spawning salmonids. Coordinate with DEQ's efforts for riparian vegetation monitoring.
- Locate and map exotic plant species.
- Determine the condition of fences along riparian corridors.

- Incorporate tax lot and building information into the GIS when it becomes available.
- Update and map changing land use information, e.g. timber harvest plans, pesticide application areas, construction projects.
- Update or map floodways along rivers.
- Educate landowners to make long-term commitments to continually remove invasive plant species.
- Whenever possible use native vegetation to landscape.

Wildlife

- Monitor existing restoration sites.
- Consider controlled burns or mowing on remnant prairies or oak woodlands. Avoid burning or mowing before 15 April or after 15 July to avoid disturbing breeding birds.
- Gather information on roadless areas and consider performing a roadless areas analysis.
- Identify areas with minimal soil disturbance having native vegetation: these areas are important refugia for insects and wildlife.
- Educate land owners not to burn brush piles; leave dead standing wood (if it will not cause damage or be a safety issue) for wildlife.

5.4.1 RECOMMENDED ANALYSES

5.4.1.1 Locate Riparian Wetlands

We recommend using the GIS to map locate potential wetland especially in the riparian areas. We have prepared a GIS layer showing hydric soils and soils with hydric inclusions. We recommend mapping the hydric soils upon the riparian land cover GIS layer to locate potential off channel wetlands. These areas can then be field verified and the data layer updated.

5.4.1.2 Locate Riparian Planting Areas

We recommend using the GIS to locate bare areas within the riparian buffer zone. These areas can then be evaluated for suitability for riparian planting. Riparian planting will provide shade to streams and slow sediment entry to the stream network.

6 AQUATIC ANALYSIS

6.1 STREAM CHANNEL MORPHOLOGY

6.1.1 DATA AVAILABLE AND DATA USED

6.1.1.1 DEM data

Many key data sets were not available for the entire study area. Therefore, we used GIS to develop surrogate or stand-in data sets for our analyses. The advantage to using these data sets is that they are comparable across the entire study area and are of a uniform spatial scale. The disadvantage is that without field verification it is impossible to know how well these data represent the actual watershed condition. We highly recommend that these data sets be field checked or that the analyses be re-run if better data sets become available.

Like any other data set, DEM (Digital Elevation Models) are appropriate for answering some questions not so good for answering others. DEM files are computer representations of topography. In a DEM grid, a 30ft X 30ft square is assigned a single elevation. These are the same data that are used to generate the familiar paper USGS 7.5' topographic quadrangles. These data are frequently used at the scale of the stream reach or watershed because general trends in topography are apparent. These data are, perhaps, not appropriate to use if one is interested in mapping subtle elevation changes along narrow first order streams.

We used DEMs to derive slopes, stream gradients, stream confinement (described below) and describe elevations within the study area (see Section 4.10)

6.1.1.2 AHI data

We were asked to summarize salmonid habitat using existing AHI data. As previously mentioned, only 12.2% (1:100K) of the streams in the study area have been surveyed by AHI crews (Maps 9-11). In addition, the dates of some surveys are more than nine years old. Because only a limited proportion of the stream network is surveyed and much of the existing information is quite old, we urge caution when interpreting summaries of these data.

6.1.1.3 GIS data

6.1.2 STREAM CONFINEMENT

Stream confinement refers to the extent to which streams are confined by hills, cliffs, terraces or other landscape features. Confinement is not the same as stream entrenchment. Stream confinement is an important factor in many watersheds because it is directly related to watershed characteristics and functions, such as presence and formation of wetlands, floodplain connectivity, availability of off-channel habitat, and flooding and peak flows. Since stream confinement information was not available for the study area, we used ARCGIS to generate a stream confinement layer from the 30ft DEMs. We

visually classified the 1:24,000 scale streams layer with a slope map that was classed into flat ($\leq 4\%$ slope) and non-flat areas ($>4\%$ slope). Table 6.1.2 shows the parameters that were used to generate unconfined and confined stream reaches. This information was also used to determine channel type.

The following table outlines the distances used to determine the channel type:

Table 6.1.2. Parameters that were used to generate stream confinement. Shown are stream order and distances from the stream that were examined (see text for explanation).		
Order	Unconfined	Confined
1-2	> 30 feet (1 pixel)	≤ 30 feet (1 pixel)
3-4	> 60 feet (2 pixels)	≤ 60 feet (2 pixels)
5-7	>120 feet (4 pixels)	≤ 120 feet (4 pixels)

6.1.3 STREAM GRADIENT

Stream gradient, the slope of the streambed, is an important watershed attribute and is an important component of salmonid habitat. Typically, steeper stream gradients have faster water velocities than flat streambeds. As water velocity increases, so does the water’s capacity to transport sediment and other materials, including large gravel, suspended sediments and large woody debris.

There are several ways to measure stream gradient. In the field, stream gradient can be measured directly with a clinometer. More commonly, stream gradient is measured from USGS 7.5 min topographic maps by measuring the change in vertical elevation (rise) over the stream segment length (run). One method is to count the number of contour intervals within a given map distance on a topographic map (Watershed Professionals Network, 1999). Stream gradient can be expressed two ways, as a percent slope (length of rise over length of run) or as the number of degrees of slope (ranges from 0° or horizontal, to 90° or vertical).

Existing stream gradient information was not available for the entire study area. We used GIS and the DEMs to calculate stream gradients.

6.1.4 CHANNEL TYPES

The relationship between a stream channel and its gradient, floodplain, and the shape of its valley are the main physical factors that structure in-stream fish habitat. Steep sided, constrained valleys and high stream gradients can result in rapidly moving water which is capable of transporting gravel and large woody debris within the stream network. Low gradient streams with unconstrained, broad floodplains are areas where transported sediment, gravel and organic debris are deposited. The relatively low gradient streams area areas where salmon spawn and juveniles rear. It is important to consider that the gravel and woody debris in these spawning and rearing stream reaches come from lower

order, head water streams. The interplay between erosion, transport and deposition is largely a function of the landforms through which a stream network flows.

Recognizing the different roles that various types of stream channels play in salmonid habitat, the OWEB manual recommends that a channel type classification be performed as part of the assessment. Channel type information was not available for the entire study area. We used GIS to classify stream channel types based on the stream order, DEM-derived stream gradient and stream confinement.

For this assessment, the derivation of stream channel types was dependent on DEM-derived stream gradients and DEM-derived stream confinement. The digital elevation model (DEM) cells are 10m X 10m in size and are, therefore, too large to detect small but biologically important topographic detail such as 1 or 2 m slope breaks that form confining stream terraces. As a result, the GIS methods used to derive confinement and gradient probably obscure important topographic information necessary to better describe these two stream attributes. Despite these limitations, we proceeded with derivation of important characteristics (gradient, confinement and stream location) from the DEM data, because at the time of this study, no other complete data were available for our analysis. The first approximation of these important stream characteristics generated from the DEM will serve as a stand-in until comprehensive data are available and have been field-checked for accuracy.

We assigned stream channel types to each stream reach using information on stream order, stream gradient and stream confinement. Due to limitations with existing data, we were not able to distinguish between moderately confined and confined streams; therefore, we did not classify moderately stream channel types (*i.e.*, LM and MM). Moreover, we did not attempt to classify Bedrock Canyons (BC) or Alluvial Fan (AF) channel types using existing information. Finally, to assign all stream reaches to a channel type, we found that it was necessary to create several new categories that are not described in the OWEB manual (Table 6.1.4). These categories represent areas where the stream channel is not confined in the upper watershed; these unconfined upper watershed stream reaches accounted for about 26.2% of the study area stream length. We highly recommend that these data be field checked.

Since stream channel types blend into one another and some channel types may fit into more than one category; therefore, the order in which we assigned channel types was important. We focused on the two ends of the stream channel continuum first, the Floodplain and Headwater stream channel types. We felt that these stream channel classes were most clear cut. Stream channel types were assigned to stream reaches in the following order: (1) Floodplain; (2) Steep Narrow Valley and Very Steep Headwater; (3) Low Gradient Confined; (4) Moderate Gradient Headwater; (5) Moderate Steep Narrow Valley; and (6) Moderate Gradient Confined (Table 6.1.4).

Table 6.1.4. Types of stream channels recognized in the OWEB manual. Shown are also the length and proportion of streams in the study area by stream channel type. Note that the estuarine channel types (ES & EL) were not present in the study area, and it was not possible to classify AF, LM, MM, BC using the data available.

Stream Channel Type	Description	Slope (%)	Stream Confinement	Stream Order	% of Study area
FP1	Low gradient large floodplain	< 1	Unconfined	6-7	3.7
FP2	Low gradient medium floodplain	< 2	Unconfined	5	5.2
FP3	Low gradient small floodplain	< 2	Unconfined	3-4	15.9
LC	Low gradient confined	< 2	Confined	variable	9.7
LU*	Low gradient unconfined	< 2	Unconfined	variable	22.2
MC	Medium gradient confined	2-4	Confined	variable	0.8
MH	Medium gradient headwater	1-6	Confined	1-2	28.5
MV	Medium gradient steep narrow valley	3-10	Confined	1-3	6.6
HU*	High gradient unconfined	> 2	Unconfined	variable	4.0
SV	Steep narrow valley	8-16	Confined	1-2	3.4
VH	Very steep headwater	>16	Confined	1-2	0.0

* Category added to accommodate unclassified stream segments

The study area is dominated by streams with Moderate Gradient Headwater (MH) channel types. This is not surprising since this stream channel category has one of the broadest definitions. It is characterized by stream gradients ranging from 1-6 degree slopes and is variable in stream order. The MH, MC, and MV channel type classes were among the most difficult to separate because of these broad definitions. The Floodplain channel types, characterized by low gradient, unconfined streams, were also well represented in the study area (24.8%). Of particular interest are the low stream order, unconfined streams (both HU and LU) that were identified in our analysis. We recommend ground truthing these reaches to see if they can be classified using more convention stream channel types.

6.1.5 SIDE CHANNEL/SECONDARY CHANNEL HABITAT

Side channels and secondary channels are important to salmonids because they provide refuge from rapid stream velocities during high flow events. The ODFW stream survey protocols define each habitat unit as either a primary (mainstem) or secondary (side) channel.

6.1.6 RIFFLES

The occurrence of riffle areas in streams is useful in evaluating fish habitat, particularly for steelhead. Under ODFW protocols, riffles are defined as areas of fast, shallow flow. The ODFW protocol divides riffles into two types, "Riffle" and "Riffle with pockets."

6.1.7 PERCENT POOLS

Pools are important to salmonids because they provide a diversity of habitats in the stream system. The variety of channel bed form and flow characteristics provided by pools give salmonids many different environments for foraging, shelter from predators

and high stream velocities, and resting. Water temperatures in pools are often layered in summer, providing deeper, cooler water for escape from high surface temperatures. ODFW stream survey protocols define pools as areas of little or no water surface gradient, having a hydraulic control such as a log, impinging streambank, boulder, bedrock wall, or other obstruction.

6.1.8 CHANNEL WIDTHS PER POOL

Pool frequency expresses how many pools are found per unit of stream length or stream area. "Channel widths per pool" is an inverse measure of pool frequency. A higher value of channel widths per pool represents a lower pool frequency, i.e., a less desirable condition (fewer pools).

6.1.9 CHANNEL WIDTH-TO-DEPTH RATIO

Channel width-to-depth ratio is of interest in watershed assessment because it is one way of describing stream channel morphology. A high width-to-depth ratio is considered undesirable because shallow water can be warmed rapidly by sunlight and surrounding warm soil and air in summer, creating temperatures too high for salmonids.

6.1.10 CHANNEL MODIFICATIONS

Channel modifications include areas where the natural stream channel has been altered or where the stream banks have been modified. Modifications include rip-rap, bridge structures, channelization, etc. Existing information on channel modifications come from the AHI data.

6.2 IN-STREAM STRUCTURE

6.2.1 POOL COMPLEXITY

Pool complexity is an estimated area of pool that has cover from wood, large substrate, and undercut banks. Average pool complexity.

6.2.2 LARGE WOODY DEBRIS

Large wood in streams provides shelter for salmon and contributes organic material (see Section 6.2.2.2). Large woody debris (LWD) and Key LWD (defined by ODFW as pieces of woody debris over 60 cm in diameter) are important components of stream structure. Large wood provides cover that can shelter salmonids from predators, and contributes organic material to the aquatic food chain. Logs provide stream structure to help reduce stream velocities, create pools, and generally diversity in the stream environment

6.2.2.1 LWD Frequency

Large woody is measured during AHI surveys and the frequency that large wood is encountered is recorded. During stream surveys, the quantity of large woody debris in a stream is expressed as "wood frequency", or pieces of wood per 100m of stream length.

6.2.2.2 LWD Source Areas

The importance of large woody debris is recognized in Pacific Northwest forests. Woody debris can directly affect the organisms that inhabit our forests by serving as shelter or as a food source. In addition, woody debris and other organic material can affect the physical environment of the forest (thus, indirectly affecting organisms) by slowing down water moving over the forest floor and into streams. Reduction of water velocity can lead to reductions in sediment delivery to forest streams. Therefore, large wood can play a role in establishing the complex terrestrial and in-stream environments favorable for many organisms, such as salmon. Of all the structural components in the terrestrial ecosystem, woody debris is one of the slowest components of the forest ecosystem to recover after disturbance (Spies *et al.* 1988). A watershed management strategy should strive to (1) identify and preserve areas that serve as large wood sources and (2) regenerate areas where large wood may no longer be present.

Many salmon habitat restoration actions involve the short-term measure of placing large wood directly in streams to enhance salmonid habitat. Longer-term strategies can also be used to manage watersheds. For example, watershed managers can plan for large wood recruitment by allowing trees to reach larger sizes in areas that may be prone to mass wasting events.

Charlie Dewberry (Simenstad *et al.*, 1997) describes a process where small "hollows" were identified in Knowles Creek. Dewberry recognized that these hollows accumulate sediments over thousands of years. During winter storms, debris torrents can originate in these hollows leading to the delivery of sediments and large wood to the stream network. Therefore, the restoration of Knowles Creek not only called for the placement of large woody debris in streams, but also planted and planned for the maturation of trees in and down slope from these hollows. These actions focused on both watershed structure and watershed function (the ecological process of sediment and large wood delivery to streams).

An analysis, such as that described by Dewberry is beyond the scope of this assessment. However, the data exist to perform this analysis and should the placement of LWD become a dominant management tool by the LWC, we recommend conducting a LWD source area analysis.

6.3 SUBSTRATES

AH1 survey crews estimate the percent of the streambed covered by each substrate particle size (silt and fine organic matter, sand, gravel, cobble, boulder, and bedrock). These data are available in the ODFW habitat-unit-level GIS layer.

6.4 STREAM FLOW

6.4.1 GAGE DATA

Stream flow is an important component of salmonid habitat. Salmon may not be able to move into, spawn or forage into some stream reaches during low flows. Humans compete with fish and wildlife for use of the water (see Section 6.5). Low flows are also related to warm stream temperature because lower volumes of water heat up more quickly than deeper water. Finally, low flows can further isolate the stream from its floodplain. Patterns in steam flows are measured by stream gauges.

We searched USGS and the Oregon Water Resource Department web sites (<http://nwis.waterdata.usgs.gov/> or nwis/discharge/ and http://www.wrd.state.or.us/surface_water/index.shtml, respectively) for information on stream gages. We found records from ten stream gauges located in the Luckiamute / Ash Creek study area (Table 6.4.1). The dates between USGS and WRD data sets closely correspond but vary in some of the stream gauge stations.

Table 6.4.1. Names and periods of record for stream gauges located in the Luckiamute/ Ash Creek study area. We included dates from both USGS and OWRD where there were discrepancies in dates. Sources were USGS and OWRD webpages.

Stream Gauge Name	Start Date	End Date	Source
Hoskins	1/5/1934	9/30/1978	USGS
	5/1934	10/1978	WRD
Little Luckiamute	1/8/1965	9/30/1971	USGS
	8/1965	9/2000	WRD
Pedee	1/10/1940	9/30/1970	USGS
	10/1940	9/1971	WRD
Rickreall Creek-Dallas	6/1926	11/1978	WRD
Rickreall Creek-Dallas Dam	10/1970	9/1979	WRD
Rickreall Creek-Mercer Dam	4/1979	9/1979	WRD
Rickreall Creek-Rickreall	4/1964	9/1985	WRD
Rickreall Creek-Salem	4/1964	9/1965	WRD
Suver	1/8/1905	9/30/2002	USGS
	8/1905	9/1901	WRD
Teal Creek	6/1968	9/1973	WRD

We gathered data from the Suver gage because it was one of the gauges with the longest periods of record. Graphs in Figure 6.4.1 show patterns in the flows for the past 99 years. The inset on the top figure shows the variability in daily mean stream flow for the early 1900s and from the late 1930's on. Notice the variability between years. The black box shows the area of the graph that is detailed in the top panel. Seasonal pattern are visible at this scale. Finally, the lower graph shows the annual mean stream

flow for the period of record. This shows multi-year drought patterns especially in the early 1990s.





Figure 7.4.1. Hydrograph the Suver gauge. Top inset shows variability in daily mean flow (CFS) for period of record, top shows season patterns in daily mean flow (CFS), and bottom shows annual mean flow (see text for details).

6.4.2 PEAK FLOWS

6.4.2.1 Land Use

Land use changes have undoubtedly had an impact on hydrologic patterns. Any action that results in less watershed water storage or quickly routes water into the stream network will increase peak flows. These land use changes and management practices include, but are not limited to, soil compaction, increases in impervious cover, tiling of agricultural fields and installation of culverts in forest road systems, draining of wetlands, channelization of stream beds and loss of flood plain connectivity.

If hydrologic changes due to land use alterations are of major concern to the LCW, we recommend the development of a GIS-based hydrologic model. Developing a hydrologic model was beyond the scope of this assessment; however, a useful model was constructed for the Tillamook Bay watershed (Melancon, 1999) and was used to evaluate hydrologic patterns (and bacterial and sediment transport patterns) as they are affected by land use changes. The advantage of creating a calibrated computer model is that multiple scenarios can be evaluated using the computer before committing to the expense of making actual changes on the ground. In this way, the cost to benefit of management actions can be evaluated.

6.4.2.2 Roads

The Watershed Assessment Manual details two methods for evaluating the impact of roads on peak flows:

1. Use of urban road density (expressed as miles of road per mi² of watershed) as a surrogate for Total Impervious Area;
2. Rural road density expressed as the percentage of total watershed area occupied by road surfaces.

We calculated road density (mi road/ mi²) for each 7th field watershed. We found that a few of the 7th field watersheds (Table 6.4.2.2a) had road densities in the high risk category (>5.5 mi roads/mi²) using the urban road density method (Watershed Professionals Network 1999). As the name implies, this screening tool is most appropriate for urban watersheds; therefore, non-urban watersheds that appear to be at risk using this approach should also be evaluated using the rural road density method (below).

<p>Table 6.4.2.2a. Seventh field watershed that are in the moderate (4.2-5.5) and high (>5.5) risk category for the potential impact of roads on peak flows. Shown are watershed HUC and name, length of roads in watershed (mi), area of watershed (mi²) and road density (mi road/ mi² of watershed).</p>
--

HUC	Name	Length (mi)	Area of HUC (mi ²)	Road Density (mi road/ mi ² watershed)
3060101	Upper Luckiamute	52.0	9.3	5.6
3060102	Miller Creek	49.5	8.9	5.6
3060201	Wolf Creek	28.9	5.0	5.7
3060404	Ritner Creek	50.3	10.0	5
3060602	Cold Springs	23.0	2.9	7.8
3060603	Black Rock Creek	33.9	5.3	6.4
3060701	Socialist Valley	46.7	7.8	6
3060702	Falls City	36.9	5.2	7.1
3060801	Upper Teal Creek	18.5	3.4	5.4
3060802	Lower Teal Creek	40.9	7.7	5.3
3060803	Grant Creek	17.0	3.2	5.3
3061101	Upper Soap Creek	55.2	9.2	6
3061302	Palestine	38.0	6.5	5.9
7020601	Upper North Fork Ash Creek	25.0	4.3	5.8
3060402	Upper Pedee Creek	33.9	7.2	4.7
7020606	Lower South Fork Ash Creek	24.1	5.2	4.7
7020603	Lower North Fork Ash Creek	24.9	5.4	4.6
3060601	Upper North Fork Ash Creek	42.4	9.4	4.5
3060303	Price Creek	23.4	5.2	4.5
3061102	Middle Soap Creek	30.0	6.8	4.4
3060304	Maxfield Creek	40.3	9.1	4.4
3060401	Lower Pedee Creek	40.7	9.4	4.3
3060305	Bump Creek	28.1	6.6	4.3
3060501	Ira Hooker	17.5	4.1	4.2

We also calculated the percent of watershed area in roads using the rural road density method (Watershed Professionals Network 1999). We assumed that the average width of a road is 35 ft (Watershed Professionals Network, 1999). This resulted in the area of road (mi²) per mi² of watershed area. The number of square miles of impervious surface per mi² of watershed ranged from 0.8 mi² roads/mi² to 5.2 mi²roads/ mi². As with the previous analysis, we found that several 7th field watersheds that were at risk for peak flow increases using the rural road density method (Table 6.4.2.2b).

We recommend that watersheds identified in Tables 6.4.2.2a & b be evaluated for mechanisms to keep water from entering the stream networks from roads. Road densities are just a screening tool to alert the watershed council to potential problems. If problems are found to exist, then various management tools/ approaches that would increase

watershed water storage could be evaluated including, detention ponds or ditches, vegetated buffer strips, etc.

Table 6.4.2.2b. Seventh field watershed that are in the moderate (4.0-8.0) risk category for the potential impact of roads on peak flows. Shown are watershed HUC and name, length of roads in watershed (mi), area of watershed (mi²) and road density (mi road/ mi² of watershed).

Watershed ID	Name	Length (mi)	Watershed area (mi ²)	Area of roads (mi ²)	Area of roads per 7 th field watershed (mi ² /mi ²)
3060602	Cold Springs	23.0	2.9	0.2	5.2%
3060702	Falls City	36.9	5.2	0.2	4.7%
3060603	Black Rock Creek	33.9	5.3	0.2	4.2%
3060701	Socialist Valley	46.7	7.8	0.3	4.0%

6.4.2.3 Rain-on-Snow

When rain falls on snow, water does not infiltrate the soil, as it normally does. Instead, water runs over the surface of the ground into the receiving stream network. This can result in high water levels in streams (high peaks on the hydrograph). Therefore, rain on snow (ROS) events can dramatically impact the pattern of water delivery to streams. As more water enters the stream network, water velocities increase, so does the capacity of the water to erode banks and down cut streambeds.

The OWEB watershed assessment manual (Watershed Professionals Network, 1999) describes watersheds as having potential impact from ROS events if two conditions are met in 20% or more of the watershed area: (1) less than 30% crown closure and (2) elevations suitable for ROS events (not defined in OWEB manual). However, the manual does not have specific guidelines for mapping these areas in the Coast Range.

We used GIS to locate areas of **potential** ROS impact, that is, areas where conditions exist that could *potentially* lead to ROS events. This is not to say that ROS events always occur in these zones. ROS events have a greater probability of occurring under certain conditions. In the Oregon Coast Range, ROS events can have return intervals of several years to tens of years.

We used land cover (see Section 4.4) and elevation (see Section 4.10) in our analysis to locate higher elevations areas that were non-forested. Non-forested areas have the potential to form a layer of snow from which rain could run off quickly into streams. The Luckiamute/ Ash Creek study area is a relatively low elevation study area compared to other areas in the Coast Range. We used GIS to locate all non-forested areas about 2,500 ft. We found one 7th field watershed, the Upper Little Luckiamute (17090003060601), in the study area which had 0.22 acres about 2,500ft. We

conclude that ROS events probably occur infrequently in the Luckiamute / Ash Creek watersheds.

6.4.3 FLOODING

Historically, there have been several large floods in or near the Luckiamute / Ash Creek study area. See Appendix B for a timeline showing important flood events including the flood of 1861, 1864, 1898, 1948, and 1996. Changes in the watershed have probably reduced the frequency of flooding along the streams in the study area. According to a BLM report (Licata *et al.*, 1998),

“Nearly all of the observed response channels in the analysis area are incised and moderately to highly unstable. Channels are “disconnected” from their floodplains (over-bank flooding occurs only during extreme storm events, if at all) which now primarily function as terraces. Water storage in floodplains has been reduced, contributing to the reduction in summer baseflows, and water quality has been degraded.”

While the down cutting of streams and lack of a connection between streams and their floodplains may have reduced the frequency of flooding, it has not improved habitat for instream organisms. Licata (1998) reports that stream down cutting has “lead to an overall reduction in the quantity and quality of aquatic life relative to reference conditions throughout the analysis area.” A certain amount of over bank flooding is necessary and floodplain connection is the natural (desirable) condition for stream ecosystems.

6.4.4 STREAM DYNAMICS

The morphology or physical form of a stream channel at any point is a dynamic expression of the climate (as it affects stream flows) and the geology (as it affects sedimentation) of a stream basin. Other variables, such as resistance to flow (friction) and bed particle size, also influence important channel variables, such as width, depth, velocity, slope, and pattern. Both human and natural disturbances to a fluvial (river) system can result in site-specific channel changes (e.g., changes in cross-sectional geometry at the point of disturbance) and/or channel morphology adjustments longitudinally over an area of stream downstream or upstream from the point of disturbance (after M. Reiter, 1995).

Human modifications of channels can cause an array of effects depending on the inherent characteristics of a system. In larger river systems these modifications rarely occur in isolation, but interact with other upstream and downstream alterations to channel morphology.

Channel erosion is the detachment and transport of material from a gully or stream channel. The material may be derived from the channel itself or material that has been deposited within the channel by surface or mass erosion. The size, complexity (sinuosity) and transport capability of channels is determined by the energy of the water, which

flows through the channel. High gradients, low friction and unimpeded water flow characterize high-energy channel systems.

Channels are unique; therefore their responses to natural factors and human-induced modifications are also unique. Changes in channel form and process occur longitudinally along a stream. In the downstream direction, the gradient decreases, sinuosity ("curviness") increases, the ratio of **bedload to total sediment load** decreases, the grain size of material which can be transported decreases, and the total **discharge** or **streamflow** increases. Large-scale determinants of channel morphology include the following factors; climate, geology/topography, vegetation and soils, land use practices, and in-channel modifications.

Fluvial processes are structured by hydrology, sediment load and movement, and the resistance of the channel to flow and sediment movement. Components of hydrology include the type of flow (baseflow, **bankfull flow**, and highflow), **stream power**, and the hydrological disturbance regime. Sediments can differ in their source, type (suspended load, bedload, turbidity), and size. Changes in sediment load that occur through land use practices can result in sediment accumulation (**aggradation**) or loss (**degradation**) in portions of the stream. Channel resistance is determined by the bank and bed material, vegetation (large wood, riparian vegetation, and roots), and physical form of the channel. Adjustments of channels include a number of factors. Channels have four degrees of freedom or ways in which the form can change: 1) **the longitudinal profile**, 2) **channel sinuosity**, 3) roughness of bed or bank, and 4) **the hydraulic radius**.

The LWC instructed the assessment team to search for information on factors that affected stream dynamics particularly log jams. We found several potential sources of this information but it was beyond the scope of this project to prepare and analyze these data sets. We recommend that the digital orthoquads be reviewed and that log jams be located and mapped. In addition, land owners can be contacted and stream walked to record log jams and other stream channel elements (i.e., riprap, boat launches, bridge supports, etc.). Finally, comment fields from Aquatic Habitat Inventory data can be queried for observations of field crews.

6.5 WATER USE

In the State of Oregon all water is publicly owned. This includes the surface water and groundwater that flows past or beneath privately owned property. A property owner does not automatically have a right to the water that flows on or near their property and, with a few exceptions, must obtain appropriate permits (water rights) to use water whether it is underground or on the surface (streams and lakes). The Oregon Water Resources Department (OWRD) issues water use permits.

Oregon's water laws, like many western US states, are based on the principle of prior appropriation. In Oregon, water appropriation doctrine has been law since February 24,

1909. Prior appropriation means that the first person to acquire a water right on a stream is the last person to be prevented from withdrawing water during low flow conditions. For this reason, water rights are ordered by date (generally date of application) and the oldest water right can demand use of the water resource to the exclusion of junior water right holders. Thus, water demand is filled by sequentially meeting demands from each diversion point from oldest to youngest.

Water use permits are acquired for particular uses. These uses include irrigation, municipal, industrial, commercial, domestic, agricultural, and other. Some uses, however, do not require permits. These water uses are called “exempt uses.” Exempt uses of **surface** water include:

1. Natural springs: if a spring does not form a natural channel or flow off of the property at any time of the year;
2. Stock watering: if livestock drink directly from surface water and there is no diversion or other modification to the water source; this includes watering livestock from a permitted reservoir or water piped (under certain conditions) to a watering tank or trough;
3. Salmon: egg incubation projects under the Salmon and Trout Enhancement Program (STEP), and as water used for fish screens, fishways and bypass structures;
4. Fire control: water withdrawn for use in emergency fire fighting;
5. Forest management: water withdrawn (under some circumstances) for certain activities such as slash burning and mixing pesticides.
6. Land management practices: water withdrawn for certain land management practices where water use is not the primary intended activity;
7. Rainwater: collection and use of rainwater does not require a permit if it is collected from an impervious surface.

Like surface water, there are some uses of ground water that do not require a permit. Generally, ground water use is allowed for a beneficial purpose without waste. These uses include:

1. Stock watering;
2. Lawn or non-commercial garden watering: of not more than one-half acre in area;
3. Single or group domestic purposes: for no more than 15,000 gallons per day;
4. Single industrial or commercial purposes: not exceeding 5,000 gallons per day;
5. Down-hole heat exchange uses;

6. Watering: the grounds, ten acres or less, of schools located within a critical ground water area.

In all cases of ground water use, Oregon's minimum well construction standards must be followed for the construction, maintenance, and abandonment of exempt wells. **For any water use, please check with the regional water master.**

If there is a conflict over use of water, the date of permit application (date of priority) determines who has first right to the water without regard to the water's use. If the dates of priority are the same in the conflict, then the law indicates that domestic use and livestock watering have preference over all other uses. However, under drought conditions, preference can be given to stock watering and domestic/ household use. Finally, ground water rights for geothermal uses are always junior in priority to other uses of water unless the water is also used for another purpose, such as irrigation, or injected back into the ground water reservoir.

The Oregon Water Trust has identified five basins in the state for water right acquisition. These priority basins are areas where there is the potential for instream water rights to benefit fish and water quality. The five basins include the Rogue, Umpqua, Deschutes, John Day and Umatilla. For more information on the Oregon Water Trust see www.owt.org.

6.5.1 GROUND WATER

In 1992 the Oregon Water Resources Department (OWRD) identified four major ground water issues in the Willamette Valley: (1) a sound, quantitative understanding of the ground-water hydrology was necessary to better manage ground- and surface-water; (2) long-term ground-water declines needed to be controlled; (3) low-yield aquifers needed to be better developed and managed; and (4) areas prone to natural ground-water quality problems needed to be identified. These issues were addressed in a two phase study initiated by the USGS and OWRD. The study examined the water resources in the Willamette Valley and focused on ground water (Oregon Water Resources Department and Oregon Department of Land Conservation and Development, 2002).

The goals of the USGS/ OWRD study were to: (1) provide a quantitative understanding of the regional ground-water flow system of the Willamette Valley sufficient to effectively evaluate the hydrologic effects of land- and water-use policies and climate changes; (2) develop the understanding and tools necessary to quantitatively evaluate the timing, location and magnitude of streamflow depletion caused by ground-water pumping; (3) characterize the unique hydrology of basalt aquifers within the Willamette Valley, particularly as related to water availability and management of multiple water-bearing zones; (4) develop a better understanding of the relations between well-yield and factors such as geology, well construction, and siting in areas underlain by low-yield aquifers; and (5) develop a better understanding of the origins and distribution of selected types of naturally occurring poor-quality ground water. The first phase of the study examined

regional patterns in water-budgets, ground-water flow directions, and relations between the streams and aquifers in the basin. The second phase of the study focused data collection at specific sites within the basin. A two phased approach was necessary because the size of the Willamette River Basin study area did not permit detailed study of every area. Studies carried out during the second phase were used to develop an understanding of specific hydrologic and geologic settings.

Western Oregon is known for its abundant precipitation (see Section 4.7); however, most precipitation falls from October through May and summer precipitation may not be sufficient to meet water demands. Therefore, many streams in the Willamette River Basin are closed to additional out-of-stream appropriations in the summer and ground water is used satisfy the growing water demand (Lee and Risley, 2002).

The OWRD has identified Critical Ground Water Areas (CGWA) where ground water resources are of special concern. In these areas, OWRD can address interference between wells, excessive water level declines, and water quality degradation. In CGWA, the OWRD can create preferences of use without regard to water right seniority, or deference to the order in which water rights were granted; however, because CGWA can require use reductions, they may be difficult to establish. There are no CGWA in the Luckiamute/Ash Creek study area (Oregon Water Resources Department and Oregon Department of Land Conservation and Development, 2002). A word of caution: where OWRD has not restricted development of the ground water resources, one cannot assume that the resource is capable of sustaining additional water use. Land use decisions and water use decisions may exacerbate water supply problems. We recommend that local planning groups work together with OWRD to prevent rural water supply problems.

The LWC requested that this assessment summarize existing ground water withdrawals. We obtained municipal well water use data from the OWRD online Water Use Reporting System. Well information includes point of diversion (POD) ID number, permit number, certification number, priority, township and range, section number, use, rate, stream name and source. We found records for 14 permitted wells in Monmouth, nine in Independence and six in Falls City. Not all wells, however, had reported withdrawals (some had 0 withdrawals listed). In addition, for those records that did have reported use, the units varied from year to year. Therefore, we converted all units to cubic feet per second (CFS) for the following comparisons. In reviewing the data, we found that following our conversions, the patterns seemed to indicate that the units were incorrectly entered into the permitting system. We followed up with the Watermaster, Bill Ferber (Dec 2003), to make sure that we interpreted the unit codes correctly. Unfortunately, there is still reason to believe that either some of the data were entered incorrectly or that the units were incorrectly recorded. Before the well withdrawal data are used for planning purposes, we strongly recommend that the Watermaster be consulted and that the data be carefully reviewed.

Peak well water usage in Monmouth was during the period of 1993-1997 (Table 6.5.1 a). According to information obtained from WARS, the well water usage is decreasing at each of Monmouth's three points of diversion (POD). The City of Independence shows

similar trends at one POD; however, the usage data were only available from 1997 to 2000. The remaining Independence PODs show an increase in water usage. Finally, PODs from Falls City recorded usage in 1989, 1999, and 2000. Recorded usages from two PODs decreased from 1989 to 2000 while the remaining two PODs each had only one record.

Tables 6.5.1 a-c show the total groundwater withdrawals for each municipality.

Table 6.5.1a. Total reported groundwater withdrawal from four Falls City wells. Units are cubit feet per second (CFS). Data were originally reported in CFS, gallons per minute (GPM), and Million Gallons per Day according to watermaster Bill Ferber. Data were converted from GPM to CFS by multiplying GPM by 0.0022280 and from MGD to GPM by multiplying by 694.44. Data are from OWRD online Water Use Reporting database.

Year	POD 11474	POD 11475	POD 11479	POD 11480
1989	126,261	NA	138,264	138,153
1999	NA	45	65	NA
2000	42	NA	117,217	NA

Table 6.5.1b. Total reported groundwater withdrawal from three Monmouth wells. Units are cubit feet per second (CFS). Data were originally reported in both CFS and gallons per minute (GPM) according to watermaster Bill Ferber. Data were converted from GPM to CFS by multiplying GPM by 0.0022280. Data are from OWRD online Water Use Reporting database.

Water	POD 11794	POD 11795	POD 30217
Year			
1989	405,000	0	NA
1990	NA	NA	NA
1991	0	0	NA
1992	0	0	NA
1993	2,404,000	2,149,100	NA
1994	3,123,500	3,877,778	34,909,100
1995	827,200	1,851,097	35,834,692
1996	667,781	358,703	43,142,327
1997	1,651,470	692,740	43,355,549
1998	8,827	4,690	790,571
1999	NA	NA	48,860,696
2000	49,515	67,091	640,593
2001	13,208	86,296	639,299

Table 6.5.1c. Total water reported water withdrawal from three Independence wells. Units are cubic feet per second (CFS). Data were originally reported in both CFS and gallons per minute (GPM) according to watermaster Bill Ferber. Data were converted from GPM to CFS by multiplying GPM by 0.0022280. Data are from OWRD online Water Use Reporting database.

Year	POD 19923	POD 24740	POD 26997	POD 30064	POD 33137	POD 33138	POD 33139
1997	NA	NA	NA	NA	118,358	NA	NA
1998	93,342	82,299	12	109,092	65,183	74,906	NA
1999	96,843	79,110	0	119,962	102,683	70,667	67,309
2000	NA	NA	45	121,739	181,783	71,200	73,939

More information on the Willamette Basin Ground-Water Study is online at http://oregon.usgs.gov/projs_dir/willgw/index.html and from <http://www.wrd.state.or.us.gov>.

6.5.2 SURFACE WATER

Water availability is determined for basins in the state through the use of water availability models. Briefly, water availability models are computer programs that are designed to predict the naturalized stream flow based on a previous period of record. For Oregon, the water availability model uses the period of record from 1958 to 1987. Once naturalized stream flows are modeled, consumptive and instream uses are subtracted according to the use permit. A detailed document describing how water allocations are made in Oregon is available online at http://www.wrd.state.or.us/programs/sw_studies/OpenFileSW02-002.pdf.

Water availability models are generally run to show water availability at the 50% and 80% exceedance levels. Fifty percent exceedance flow is the modeled stream flow that occurs, on average, one of every two years, or 50% of the time. This modeled water availability is used to determine if water can be captured for water storage. The 80% exceedance level, the amount of streamflow that occurs on average four out of five years, is used to determine if water is available for issuing a new surface water permit. Modeled streamflows are generally calculated for every month at particular positions along the stream network.

Generally, before issuing a new water right, available water is compared against permitted withdrawals. If a stream is fully appropriated (all the water is permitted) no new withdrawals can be permitted.

6.5.2.1 Points of Diversion & Types of Water Use

In order to determine the status of the water resources in the Luckiamute basin, we first queried the Oregon Water Resources GIS database containing the point of diversion and places of use. This GIS layer was built in March of 2001. Permitted and certificated water rights were digitized (excluding municipal rights) and have a planned update cycle of 6 weeks. In this GIS theme, each point represents a location where water is diverted for

use according to the terms in the water right. We found that there was a total of 933 unique permit numbers for the study area, 711 in the Luckiamute watershed and 222 in the Rickreall watershed. Each permit may have multiple uses designated. We found a total of 1,458 designated water uses in 26 use categories listed on these permits (Table 6.5.2.1). Irrigation was the most commonly designated water use accounting for 53.6% of the designated water uses. Other common water uses included: livestock (16.8%), storage (7.3%), domestic (5.6%), supplemental irrigation (4.5%), and municipal (2.6%).

Keep in mind that not all water rights are in use at any one time. And, during times of water shortages, junior permit holders may be denied use. Finally, we did find at least one permit that was incorrectly entered into the WARS database; Permit S 41181 was identified as being on the Mary's River although according to the GIS data set it was located in the Luckiamute watershed. Although we carefully examined the data in Table 6.5.2.1, there may be other errors. We recommend that the Watermaster be contacted before POD data are used for detailed planning purposes.

Table 6.5.2.1. Summary of the water withdrawal permits in the study area by watershed (Luckiamute and Rickreall). Shown are the source streams and permitted usage for all types of permits. AG= agriculture, CM=commercial, DI= domestic including lawn and garden, DN=domestic expanded including non-commercial garden , DO=domestic , DS=domestic & livestock , FI=fish culture, FP=fire protection , FW=fish & wildlife , GD=group domestic , I*=irrigation, livestock & domestic, ID=irrigation and domestic , IL=irrigation & livestock , IM=manufacturing , IR=irrigation, IS=supplemental irrigation, LV=livestock, MM=unspecified, MU=municipal, PW=power development, QM=quasi-municipal, RC=recreation, ST=storage , TC=temperature control, WI=wildlife, and BL=unspecified. For details on permitted uses see www.wrd.state.or.us.

Watershed	STREAM2_	A	C	D	D	D	D	F	F	F	G	I	I	I	I	I	L	M	M	P	Q	R	ST	T	W	B	Tot	
	NA	G	M	I	N	O	S	I	P	W	D	*	D	L	M	IR	S	LV	M	M	P	Q	R	ST	T	W	B	al
Luckiamute	AIRLIE CR														1								1				2	
	BEAVER CR							1							1													2
	BERRY CR					2										3		4					2					11
	BONNER CR					1																						1
	BOUGHEY CR			1																								1
	BUMP CR			1		1												4										6
	COLUMBIA R															1												1
	COOPER CR															1		6					4					11
	CRABTREE CR								1									1										2
	DUTCH CR			1														4										5
	EVERZ CR					2												1					1					4
	FARLEY CR																	3										3
	FERN CR					1			1						1			1					3		1			8
	FULLER CR																						1					1
	GLAZE CR																		1									1
	GRANT CR															1		1						2				4
	JONT CR															5								1				6
KINSEY CR					1																						1	

Table 6.5.2.1. Summary of the water withdrawal permits in the study area by watershed (Luckiamute and Rickreall). Shown are the source streams and permitted usage for all types of permits. AG= agriculture, CM=commercial, DI= domestic including lawn and garden, DN=domestic expanded including non-commercial garden , DO=domestic , DS=domestic & livestock , FI=fish culture, FP=fire protection , FW=fish & wildlife , GD=group domestic , I*=irrigation, livestock & domestic, ID=irrigation and domestic , IL=irrigation & livestock , IM=manufacturing , IR=irrigation, IS=supplemental irrigation, LV=livestock, MM=unspecified, MU=municipal, PW=power development, QM=quasi-municipal, RC=recreation, ST=storage , TC=temperature control, WI=wildlife, and BL=unspecified. For details on permitted uses see www.wrd.state.or.us.

Watershed	STREAM2_ NA	A G	C M	D I	D N	D O	D S	F I	F P	F W	G D	I *	I D	I L	I M	I R	I S	L V	M M	M U	P W	Q M	R C	S T	T C	W I	B L	Tot al
	LITTLE LUCKIAMU TE R			1		13										64	3	33		3	1		1	14			2	135
	LUCKIAMU TE R	3		2	1	16	4	3	3	1			3	2	4	22	1	52			1			29		1	2	365
	MARYS R					1																						1
	MAXFIELD CR			1												3	4							1		2		11
	MCTIMMO NDS CR					1		2								2		1						2				8
	N FK PEDEE CR					2										1		2						1				6
	N SANTIAM R																	4										4
	PEDEE CR				1											1		1										3
	PEEDEE CR					1										1		3										5
	PETERSON CR															5		3										8
	PLUNKETT CR		2									4		2														8
	PRICE CR								1							2								2				5
	RITNER CR			6		3										4		35										48
	ROCK CR															1												1
	S FK ASH CR															1		2										3
	SANTIAM R					1																						1

Table 6.5.2.1. Summary of the water withdrawal permits in the study area by watershed (Luckiamute and Rickreall). Shown are the source streams and permitted usage for all types of permits. AG= agriculture, CM=commercial, DI= domestic including lawn and garden, DN=domestic expanded including non-commercial garden , DO=domestic , DS=domestic & livestock , FI=fish culture, FP=fire protection , FW=fish & wildlife , GD=group domestic , I*=irrigation, livestock & domestic, ID=irrigation and domestic , IL=irrigation & livestock , IM=manufacturing , IR=irrigation, IS=supplemental irrigation, LV=livestock, MM=unspecified, MU=municipal, PW=power development, QM=quasi-municipal, RC=recreation, ST=storage , TC=temperature control, WI=wildlife, and BL=unspecified. For details on permitted uses see www.wrd.state.or.us.

Watershed	STREAM2_	A	C	D	D	D	D	F	F	F	G	I	I	I	I	I	I	M	M	P	Q	R	ST	T	W	B	Tot						
	NA	G	M	I	N	O	S	I	P	W	D	*	D	L	M	IR	S	LV	M	M	U	P	W	Q	M	R	C	ST	T	C	I	L	al
Luckiamute	SOAP CR			1		2	1	4	1			1			1	13	2						1	6		2							35
	TEAL CR			1		9		1	2							8	2	1	1	16			7	4									52
	UNN STR			2	1	11										14	4	16							3								51
	VINCENT CR			1				1							1	4		1					1	3									12
	WALKER CR															1																	1
	WAYMIRE CR					2		2								3	1	3							2								13
	WILLAMETTE R			1	2	2	1	3					2		1	22	2	4	2	10					1		1						270
	(blank)								3							20	1	8		6				1		2	5						46
	ASH CR								1							1	22		26				1	3		1							55
Rickreall	COLUMBIA R	1														31	4						1									37	
	DUCK SL															19																	19
	FERN CR															1								1									2
	HARTMAN SLOUGH															3																	3
	LITTLE LUCKIAMUTE CR															1																	1
	LUCKIAMUTE R															2																	2
	M FK ASH CR					2										1		4						3	1								11

Table 6.5.2.1. Summary of the water withdrawal permits in the study area by watershed (Luckiamute and Rickreall). Shown are the source streams and permitted usage for all types of permits. AG= agriculture, CM=commercial, DI= domestic including lawn and garden, DN=domestic expanded including non-commercial garden , DO=domestic , DS=domestic & livestock , FI=fish culture, FP=fire protection , FW=fish & wildlife , GD=group domestic , I*=irrigation, livestock & domestic, ID=irrigation and domestic , IL=irrigation & livestock , IM=manufacturing , IR=irrigation, IS=supplemental irrigation, LV=livestock, MM=unspecified, MU=municipal, PW=power development, QM=quasi-municipal, RC=recreation, ST=storage , TC=temperature control, WI=wildlife, and BL=unspecified. For details on permitted uses see www.wrd.state.or.us.

Watershed	STREAM2_ NA	A G	C M	D I	D N	D O	D S	F I	F P	F W	G D	I *	I D	I L	I M	I R	I S	L V	M M	M U	P W	Q M	R C	S T	T C	W I	B L	Tot al
	N FK ASH CR							2								5		1						5				13
	RICKREAL L CR					1																						1
	S FK ASH CR															5		8		3				7				23
	UNN STR			1		4		1								5	1	6										18
	WILLAMET TE R	1						2			1				2	74	4	1		9				3				97
	(blank)					3										6	2					2		1				14
Total		5	2	2 0	5	82	7	2 7	7	1	1	5	5	4	2	1 78	6 2	24 5	1	38	2	2	12	10 7	1	1 2	7	145 8

6.5.2.2 Water Availability

We queried the Water Availability Report System (WARS) for the Luckiamute watershed (wars.wrd.state.or.us) to determine the status of the water resources. Table XX shows water availability, at the 50% and 80% exceedance levels, for five water availability basins. We found that, in general, the watersheds won't support any additional summer and early fall water withdrawals.

Information on individual water rights can be found at http://stamp.wrd.state.or.us/apps/wr/wrinfo/wrinfo.php?search_type=FindStream and <http://www.wrd.state.or.us/>.

Table 6.5.2.3. Water availability for the Luckiamute watershed at 50% and 80% exceedance modeled streamflow by month. Permits for water storage (e.g., reservoirs) are based on the 50% exceedance values and permits for other uses are based on 80% exceedance values (see text) Data as of November 2003.					
Name/ Watershed ID	Month	50% Exceedance		80 % Exceedance	
		Total CFS Used	Available	Total CFS Used	Available
Luckiamute River > Willamette River (Mouth) 117	J	1750.0	YES	840.0	YES
	F	1720.0	YES	938.0	YES
	M	1250.0	YES	751.0	YES
	A	745.0	YES	481.0	YES
	M	366.0	YES	235.0	YES
	J	142.0	YES	81.6	YES
	J	25.9	YES	-1.9	NO
	A	-1.2	NO	-12.9	NO
	S	11.8	YES	-1.5	NO
	O	49.0	YES	19.9	YES
	N	524.0	YES	149.0	YES
	D	1670.0	YES	727.0	YES
Luckiamute River > at Soap Creek Willamette River 118	J	1450.0	YES	691.0	YES
	F	1420.0	YES	733.0	YES
	M	1040.0	YES	622.0	YES
	A	616.0	YES	402.0	YES
	M	309.0	YES	196.0	YES
	J	120.0	YES	70.9	YES
	J	25.0	YES	--	NO
	A	-1.5	NO	--	NO
	S	8.6	YES	-4.4	NO
	O	40.9	YES	12.2	YES
	N	480.0	YES	136.0	YES
	D	1390.0	YES	621.0	YES
Luckiamute River >	J	717.0	YES	348.0	YES
	F	690.0	YES	377.0	YES

Table 6.5.2.3. Water availability for the Luckiamute watershed at 50% and 80% exceedance modeled streamflow by month. Permits for water storage (e.g., reservoirs) are based on the 50% exceedance values and permits for other uses are based on 80% exceedance values (see text) Data as of November 2003.

Name/ Watershed ID	Month	50% Exceedance		80 % Exceedance	
		Total CFS Used	Available	Total CFS Used	Available
Willamette River at McTimmonds Creek 119	M	522.0	YES	303.0	YES
	A	295.0	YES	192.0	YES
	M	157.0	YES	97.9	YES
	J	62.7	YES	36.7	YES
	J	13.7	YES	--	NO
	A	-2.7	NO	--	NO
	S	-0.3	NO	-8.1	NO
	O	16.4	YES	0.6	YES
	N	247.0	YES	69.6	YES
	D	708.0	YES	312.0	YES
Luckiamute River > Willamette River at Kopplein Creek 120	J	316.0	YES	151.0	YES
	F	306.0	YES	166.0	YES
	M	246.0	YES	142.0	YES
	A	149.0	YES	93.2	YES
	M	73.7	YES	46.0	YES
	J	30.9	YES	20.0	YES
	J	11.8	YES	--	NO
	A	--	NO	--	NO
	S	--	NO	--	NO
	O	14.0	YES	--	NO
Pedee Creek > Luckiamute River (Mouth) 148	N	164.0	YES	52.9	YES
	D	321.0	YES	158.0	YES
	J	97.0	YES	32.4	YES
	F	94.0	YES	38.6	YES
	M	67.4	YES	27.9	YES
	A	26.9	YES	7.5	YES
	M	-1.1	NO	-9.5	NO
	J	1.3	YES	-4.5	NO
	J	0.1	YES	-2.9	NO
	A	----	NO	-12.9	NO
S	-2.2	NO	-8.1	NO	
O	-4.4	NO	-7.0	NO	
N	22.4	YES	-9.7	NO	
D	95.0	YES	28.5	YES	

6.5.3 INTERPRETATION AND RECOMMENDATIONS

Approximately 70% of Oregon's population lives in the Willamette Valley (Lee and Risley, 2002). Water is an important natural resource for both humans and for fish and wildlife. As such, state law regulates water use. Protection and thoughtful use of this resource are necessary to ensure that there will be enough water to meet demand. We recommend the following actions:

- Educate yourself and new residents:
 - Become familiar with water used planning in your community
 - Know where your water comes from
 - Protect sensitive aquifers and recharge areas
- Establish stream gauging stations, weather stations and rainfall gages to improve knowledge of water availability.
- Verify mapped points of water diversion.
- Map (or verify) spring and well locations.
- Document areas of ground water shortages and water quality problems from well logs.
- Begin to gather information on the location of the water table. Subsurface water flow entering streams may help to maintain cool water temperatures necessary for good salmonid habitat.
- Conserve water
 - Indoors, use low-flow showers and toilets
 - Outdoors, limit the size of irrigated landscaping; require timed sprinklers.
 - Consider rainwater collection and storage systems.

6.6 WATER QUALITY

Water quality is a term that is often used to describe many properties of bodies of water including, but not limited to temperature, nutrient concentration (most commonly nitrogen and phosphorus), pH, conductivity, alkalinity, dissolved oxygen concentration, contaminant (pollutant) concentration, and concentration of indicator bacteria. All of these factors vary in time and space within streams, rivers, lakes, and estuaries, which make them very difficult to study. Yet, water quality often limits (in biological terms) the types and abundance of organisms that live in these aquatic environments.

For this assessment, we focused on existing data sources. We used the Oregon DEQ 303(d) list, and data contained in EPA STORET database. STORET is used by DEQ to determine which stream segments are of poor water quality and will not support the designated use (see Section 6.6.2).

6.6.1 FACTORS AFFECTING WATER QUALITY

A number of factors can affect water quality including land use, land cover, terrain, presence of point and non-point pollution sources, etc. Because water quality is affected by so many factors, it is often thought of as an integrator of watershed condition. Larger stream reaches and receiving bodies of water (i.e., estuaries and lakes) are often assessed for 'cumulative effects.' That is, the influence of 'stressors' or factors that adversely affect water quality throughout the entire watershed.

6.6.1.1 Point Source Pollution

The 1972 Federal Water Pollution Control Act defined two sources of pollution: point and nonpoint. **Point sources** of pollution can be clearly identified; examples include discharges from industry and sewage treatment plants. Such discharges often enter the receiving waters *via* a discharge pipe. All point sources discharging into navigable waters are regulated by the National Pollutant Discharge Elimination System (NPDES). In Oregon, the Department of Environmental Quality is responsible for implementing components of the NPDES program, such as storm water discharge permits.

The purpose of the NPDES Program is to protect human health and the environment. By point sources, EPA means discrete conveyances such as pipes or man made ditches. All facilities (excluding individual households) must obtain permits if their discharges go directly to surface waters. Examples of pollutants that may threaten public health and the nation's waters are: human wastes, ground-up food from sink disposals, laundry and bath waters, toxic chemicals, oil and grease, metals, and pesticides (EPA, <http://www.epa.gov/owm/npdes.htm>).

Point sources of pollution include wastewater treatment plants and other effluent discharges. The Clean Water Act requires that all point sources discharging pollutants into waters of the United States must obtain an NPDES permit. This includes storm water discharges associated with "industrial activity," according to a fact sheet put out by ODEQ. Industrial activity is defined as having the industry listed by EPA or having storm or snow melt leaving the site through a point source (pipe, culvert, ditch, basin, channel, etc.) and reaching surface waters directly or through storm drainage. Some construction activities are also included.

We found six active NPDES permit holders in the study area (Table 6.6.1.1). Records for each facility were examined online. All facilities were found to be operating within the conditions of their permits. No prior violations were recorded.

Table 6.6.1.1. Active NPDES permits found within the study area. Shown are the NPDES ID, facility name, county, permit issue and expiration dates, and facility description. Source was EPA web site.					
NPDES ID	Facility Name	County	Permit Issue Date	Permit Expire Date	Description
OR0043630	VALLEY LANDFILLS INC	BENTON	FEB-13-1998	DEC-31-2002	REFUSE SYSTEMS
ORG253505	FRANKLIN SWEED INCORPORATED	POLK	NOV-05-1996	JUL-31-2001	GRAY AND DUCTILE IRON FOUNDRIES
OR0020613	CITY OF MONMOUTH WASTE WATER TREATMENT FACILITY	POLK	AUG-10-1994	MAY-31-1999	SEWERAGE SYSTEMS
OR0032701	FALLS CITY, CITY OF	POLK	MAR-08-1991	FEB-28-1996	SEWERAGE SYSTEMS
OR0020443	INDEPENDENCE, CITY OF	POLK	AUG-10-1994	MAY-31-1999	SEWERAGE SYSTEMS
ORG703500	BOISE CASCADE CORPORATION	POLK	APR-14-1997	JUL-31-2001	SOFTWOOD VENEER AND PLYWOOD

6.6.1.2 Non-Point Source Pollution

Nonpoint sources of pollution may have no readily identifiable source, or may originate from broad areas rather than discrete points. Examples are pesticides entering streams from aerial spraying; run-off from urban, construction, and agricultural activities; animal wastes entering streams from pastures; and septic tank seepage. Nonpoint source pollution can enter the receiving waters via overland or underground flow. It is much more difficult to identify and manage non-point sources of pollution than point sources.

No GIS data were available on non-point pollution sources, and therefore we were not able to prioritize 7th field watersheds on this basis. However, both the Luckiamute River and Soap Creek appear on the state 303(d) list for low dissolved oxygen and bacterial concentrations (see Section 6.6.2); since many stream miles are listed both of these water quality impairments may be indicative of non-point source pollution. Part of good watershed management includes awareness of these pollution sources. We recommend that local watershed groups work towards increasing awareness of nonpoint pollution sources, and take action to reduce these pollution sources. Examples of actions that can reduce pollutants entering streams from surface water runoff include riparian fencing, riparian plantings, grazing management and pasture rotation, and education for responsible pesticide use.

6.6.1.3 Water Temperature

Stream temperature is of concern within the study area. A number of water temperature monitoring projects have been conducted or are ongoing within the study area (Table 4.1.4). The South Fork Berry Creek is on the 303(d) due to stream temperatures that exceed water quality standards.

Stream temperature is important for several reasons. First, temperature directly affects the amount of dissolved oxygen that water contains and, therefore, the productivity of the stream. Second, aquatic organisms have varying tolerances to temperature: salmon, in particular, are sensitive to warm temperatures. According to a fact sheet published by DEQ (http://www.deq.state.or.us/pubs/water/Stream_Temperature.pdf) Oregon salmonids require water temperatures to be 10° C for spawning and 17.8° C for all other life stages. Oregon DEQ temperature standards are based on a 7-day moving average of the high temperatures in a stream. There are also many indirect effects of stream temperature on the nature of streams. For example, temperature affects the viscosity of water; therefore cold water travels a little slower and transports more suspended particulates than warm water (e.g., silt sinks twice as fast at 23° C than at 0° C: (Hynes, 1970)).

Temperature fluctuates on daily and seasonal time scales. In aquatic ecosystems, this variability makes it difficult to interpret instantaneous measurements (discrete in time and space), which are often recorded by water quality monitoring teams. Because of this variability temperature data loggers are often used to make measurements at frequent, repeated intervals in streams. Models are often used to integrate these measurements over space.

To determine the current trends in water quality we queried the LASAR water quality database for stations where water temperature had been measured (Table 6.6.3.1). We found that only six stations had measurements made with the past three years. Figure 6.6.13 show temperature from three stations. Unfortunately, existing temperature data were not useful in prioritizing 7th field watersheds in this assessment.




Figure 6.6.1.3. Water temperature from three stations in the Luckiamute / Ash Creek Study area. These are the monitoring stations for which recent temperature data are recorded in the LASAR database.

We recommend that LWC use the available water temperature data at the stream reach and basin planning scale to prioritize project sites. Data gathered during this assessment can be combined with water temperature data to provide powerful tools for action planning. For example, where a monitoring program shows a consistently high water temperature, AHI data, DOQs or local knowledge should be investigated to determine where in the watershed stream bank shading may be poor and riparian vegetation may be lacking. Riparian plantings and riparian fencing can then be planned for appropriate sites. We also recommend that stream temperature models be considered. Models can be developed that integrate temperature over space and time, based on a relatively few number of measurements or based on emerging technology, such as, FLIR imagery.

6.6.2 WATER QUALITY EVALUATION
303(d) Listed Streams

The 1972 Federal Water Pollution Control Act (amended as The Clean Water Act in 1977) established broad water quality goals for the nation’s fishable and swimmable waters. The Oregon Department of Environmental Quality (ODEQ) is one of the agencies that monitor water quality in the State of Oregon. ODEQ is required by the federal Clean Water Act to maintain a list of steam segments that do not meet water quality standards, the so-called 303(d) list. Water bodies that do not meet water quality standards are said to be water quality limited or impaired. The term, “water quality limited”, refers to a limitation in a beneficial use of that water body. Beneficial uses of state waters, as defined by the Oregon Legislature (ORS 468.710) include: domestic, municipal, irrigation, power development, industrial, mining, recreation, wildlife and fish uses, and pollution abatement. Water quality standards, levels or concentrations of water quality variables, such as fecal coliform bacteria, temperature, or dissolved oxygen concentration, have been established to classify state waters as "supporting", "partially-supporting", or "not-supporting" certain beneficial uses.

The 303(d) list is used as a first step in locating water quality-impaired reaches, as described in the OWEB Watershed Assessment Manual. The 303(d) list does not include all streams that are impaired by high temperatures, sedimentation, fecal coliform, or other factors; however, all streams appearing on the 303(d) list have been assessed. This may reflect the methods used to designate 303(d) streams (*i.e.*, larger rivers may receive more scrutiny during the designation process) as well as actual differences in water quality. Streams suspected to be water quality impaired for which data have not been collected appear on ODEQ’s “Water Bodies of Concern” list.

We obtained database files and GIS coverages of Oregon's 2002 List of Water Quality Limited Water bodies (the "303[d] list") from the ODEQ website (Map 12). In 2002, there were 13,300 miles of Oregon streams listed on the 303(d) list including three streams in the study area (Table 6.6.2).

Table 6.6.2. Water quality limited stream in the study area. These streams appear on ODEQ’s 303(d) list.						
Record ID	Water body Name	River Mile	Parameter	Season	List Date	Listing Status
6054	Luckiamute River	0 to 31.7	Fecal Coliform	Winter/Spring/Fall	1998	303(d) List
8523	Soap Creek	0 to 16.8	Dissolved Oxygen	October 1 - May 31	2002	303(d) List

Table 6.6.2. Water quality limited stream in the study area. These streams appear on ODEQ's 303(d) list.						
Record ID	Water body Name	River Mile	Parameter	Season	List Date	Listing Status
8791	South Fork Berry Creek	0 to 2.1	Temperature	Summer	2002	303(d) List

Almost 32 miles of the Luckiamute River appear on the 303(d) list. The Luckiamute River was listed in 1998 for fecal coliform bacteria concentrations that exceeded the water quality standards for water contact recreation (the beneficial use). The river is listed for winter, spring and fall. Fecal coliform bacteria are themselves an indicator of the potential human health risk due to waterborne pathogens.

Almost 17 miles of Soap Creek are listed for not meeting the dissolved oxygen concentration water quality standard from October through May. Soap Creek was listed in 2002 and salmonid spawning was the beneficial use that was impaired.

Finally, about two miles of the South Fork of Berry Creek were listed for temperature in 2002. Stream water temperature exceeded the state water quality standards (17.8°C) for salmon fish rearing. Salmonid fish rearing and anadromous fish passage are the beneficial uses that were affected.

Compare Map 12 with Maps 9-11. These are areas where water quality is likely to have a large impact on salmonid populations. Recall from Section 6.7 that water quality can also restrict the movement of salmon and act as a barrier. We recommend that water quality locations be established on these reaches.

In a relatively new approach to managing water quality, Total Maximum Daily Loads (TMDL) are being developed. USEPA requires the state to develop TMDL that take in to account pollution from all sources that a water body can receive including discharges from industry, sewage treatment facilities, and runoff from agricultural, forests, and urban areas. Although EPA has not approved TMDLS in the Luckiamute basin, the Upper Willamette was due to have TMDL (for temperature, bacteria and mercury) developed in 2003 and are now being reviewed. We recommend that LWC keep abreast of and participate in the TMDL process.

6.6.2.1 High Quality Waters

The LWC requested information on areas of high water quality in the Luckiamute/ Ash Creek study area. We checked Oregon's Waterwatch (www.waterwatch.org), America's Scenic Byways (www.byways.org), and Oregon Wild and Scenic Rivers Program (www.oregonwaters.org) for information on high quality waterways. We did not find any information on high quality waterways in the study area.

6.6.3 WATER QUALITY MONITORING

6.6.3.1 EPA's STORET and ODEQ's LASER

Water quality data, measured by federal, state and private groups, is available via two online databases, STORET and LASAR. The STORET (short for STOrage and RETrieval) database is a repository for water quality, biological, and physical data. STORET contains raw biological,

chemical, and physical data on surface and ground water collected by federal, state and local agencies, Indian Tribes, volunteer groups, academics, and others. Data collected from all 50 States, territories, and jurisdictions of the U.S., along with portions of Canada and Mexico, are stored in the system. If water quality was measured, it generally ends up in the STORET database.

Currently, STORET data are available as two separate databases, divided according to when data were originally supplied to EPA. The older of the two databases is called the STORET Legacy Data Center (LDC for short), and the more current is called Modernized STORET. Water quality observations made prior to 1999 are stored in the LDC database. Both data sets are available via the Internet (<http://www.epa.gov/storet/>).

LASAR (Laboratory Analytical Storage and Retrieval Database) was developed and maintained by Oregon Department of Environmental Quality (ODEQ). Generally, collected water quality data appears in both STORET and LASAR. The following is a brief description of how the data are organized in STORET and LASAR. Individual water quality measurements, called parameters, are given unique parameter codes. Within the STORET database parameters are grouped into 18 major categories (group codes) which include administrative, bacteriological, biological, dissolved oxygen concentration, flow, general inorganic, general organic, metal, nitrogen, oxygen demand, pesticide, phosphorus, physical, radiological, solid, temperature, miscellaneous, and other. Measurements are made at STORET stations, each identified by a unique number. Data can be retrieved online by 4th field HUC, by station, or by parameter number or group codes.

We queried both STORET (December 2003) and LASAR (July 2003) for water quality information collected within the study area. Table 6.6.3.1 summarizes water quality stations for which data were collected.

Table 6.6.3.1. Range of dates for which water quality measurements were made. Shown are station ID numbers. Data were obtained from LASAR in July 2003.

1970-1980	1981-1990	1991-2000		2001-present
10347	10348	11895	12456	10658
10844	14425	12433	12457	10659
11292	14426	12434	12458	11111
11293	14427	12435	14429	11113
11294	14428	12436	14435	11114
11295	14430	12437	14436	11118
11296	14431	12438	14438	
11297	14432	12439	14442	
11298	14433	12440	14444	
11299	14434	12441	14445	
11300	14437	12442	14446	
11301	14439	12443	14449	
11302	14440	12444	14450	
11109	14441	12445	14451	
11110	14443	12446	14452	

Table 6.6.3.1. Range of dates for which water quality measurements were made. Shown are station ID numbers. Data were obtained from LASAR in July 2003.

1970-1980	1981-1990	1991-2000		2001-present
11112	14447	12447	14453	
11115	14448	12448	14454	
11117		12449	14455	
11319		12450	14457	
14424		12451	14458	
		12452	14459	
		12453	14460	
		12454	14461	
		12455	23866	

The following water quality monitoring stations were examined but contained no data records: 11894, 11896, 12402, 12459, 16479, 16480, 1116, 14067, 14608, 14609, 14456, 16479, and 16480.

In general, we found that water quality was measured infrequently and not in enough locations to be of use in prioritizing 7th field watersheds. Only six water quality monitoring stations had measurements taken within the last three years. These stations were: Luckiamute River at Lower Bridge (ID 10658), Luckiamute River at Helmick St Park (ID 10659), Luckiamute River at Hoskins (ID 11111), Soap Creek at Corvallis Rd (Suver: ID 11113), Little Luckiamute River at Elkins Road (ID 11114), and Teal Creek at Gardner Rd (Falls City: ID 11118). Fecal coliform bacteria were measured at most of these stations; unfortunately, the measurements were made infrequently and at irregular intervals. For example, three of the sites had only 5-10 measurements made from the mid 1970's to 2002. The remaining two sites had between 16 and 20 measurements made between 1968 and the mid 1980's. These data are not suitable for looking at trends within the study area. We provide summaries of the STORET data and sampling locations to assist the Luckiamute Watershed Council in their Action Planning.

6.7 FISH BARRIERS

Fish barriers have been important to Oregon's citizens since before Oregon was a state. In 1948, the new Oregon Territory constitution prohibited the obstruction of salmon streams; if a project did obstruct a salmon stream, it required the construction of fish passage facilities. Oregon's first game laws, passed in 1872, included requirements for fishways to take precedence over dams. Current laws give the Oregon Department of Fish and Wildlife authority to require maintenance of fish passage at all man-made in-channel obstructions in streams where fish are present (ORS 498.268 and ORS 509.605 through 509.645). These laws make dam owners and operators responsible for installing and maintaining adequate fish passage facilities, with some exceptions (<http://www.dfw.state.or.us/odfwhtml/infocntrfish/management/fishpassage.txt>).

Barriers to fish passage can be physical, chemical or behavioral. There is also a temporal component to barriers: landslides and debris flows can temporarily block fish passage until the stream cuts through or around the obstacle. Sometimes permanent natural blockages can form (e.g.,

as a result of volcanoes or glaciers) and some fish runs become landlocked. Most frequently, fish barriers are thought of as the artificial structures in stream channels that may restrict or eliminate the ability of fish to move up and downstream. For example, culverts that create water velocities exceeding the swimming ability of the fish (especially for juvenile fish), or dams. However, obstacles other than physical structures can also block fish passage. Stream reaches with chronically low dissolved oxygen concentrations, high temperatures, or toxic contaminants can also impede fish passage (see Section 6.6.2 and Map 12).

6.7.1 CULVERTS

Several sources of culvert data exist. We planned to examine the culvert data available on streamnet and from Benton and Polk Co. The streamnet data were not complete. We obtained the data from Benton County but not from Polk County. We recommend that LWC acquire and review the culvert data. These data can be augmented by culvert surveys: we recommend that the LWC located and map culverts on main roads.

6.7.2 DAMS

We gathered available information on dams from ODFW. There are several dams that are recorded for the Luckiamute/ Ash Creek study area. These dams should be field checked to ensure that they have not been removed.

Table 6.7.2a. Dams listed in the ODFW dams database.		
Dam Name	Owner	Maximum Storage
Unnamed dam	Unknown	Unknown
Unnamed dam	Unknown	Unknown
Unnamed dam	Unknown	Unknown
Mulkey, Gylan Reservoir Dam	Gylan Mulkey	50.00
Kennel Reservoir Dam	Earl Kennel	160.00
Unnamed dam	Sam Oberg	Unknown
Emory Moore Dam	James Heggemeir	166.00
Unnamed diversion dam	Unknown	Unknown
Unnamed diversion dam	Unknown	Unknown
Unnamed dam	Unknown	Unknown
Unnamed dam	Unknown	Unknown
Whispering Winds Dam (Lake of the Winds Dam)	Girl Scout Camp	100.00
Unnamed dam	Unknown	Unknown

Since 1983 the Northwest Power Planning Council has directed studies of existing salmonid habitat. In 1988, the Council concluded that: 1) the studies had identified fish and wildlife resources of critical importance to the region; 2) mitigation techniques cannot assure that all adverse impacts of hydroelectric development on these fish and wildlife populations will be mitigated; 3) even small

hydroelectric projects may have unacceptable individual and cumulative impacts on these resources; and 4) protecting these resources and habitats from hydroelectric development is consistent with an adequate, efficient, economical, and reliable power supply. Consequently, the Council also has considered alternative means of habitat protection.

As a result, the Council has designated certain river reaches as "protected areas." Protected areas are stream reaches where the Council believes hydroelectric development would have unacceptable risks of loss to fish and wildlife species of concern, their productive capacity, or their habitat. River reaches to be protected are those reaches or portions of reaches listed on the "Protected Areas List" adopted by the Council on August 10, 1988. Table 6.7.2b shows river reaches considered and those appearing on the Protected Areas List. No dams exist on the Luckiamute system (Wevers *et al.*, 1992).

Table 6.7.2b. Protection status adopted by the Northwest Power Planning Council in 1988. Shown are status (1=Anadromous and Resident Fish or Wildlife; 2=Anadromous; and 3=Unprotected), stream, start reach and end reach and number of miles. Source was Streamnet.org, December 2003.				
Status	Stream	Start Reach	End Reach	Reach Length (Mi)
1	Little Luckiamute R	Waymire Cr	Berry Cr	2.5
2	Clayton Cr	Mouth	Headwaters	3
2	Little Luckiamute R	Berry Cr	Dutch Cr	0.1
		Black Rock Cr	Headwaters	0.5
		Cooper Cr	Fern Cr	6
		Dutch Cr	Sams Cr	2.1
		Farley Cr	Waymire Cr	1.2
		Fern Cr	Teal Cr	1.3
		Mouth	Cooper Cr	1.5
		Sams Cr	Black Rock Cr	1.2
		Teal Cr	Farley Cr	1
2	Luckiamute R	Beaver Cr	Boulder Cr	0.5
		Bonner Cr	Cougar Cr	9.4
		Boulder Cr	Headwaters	0.5
		Cougar Cr	Rock Pit Cr	1.5
		Jont Cr	Mctimmonds Cr	7.8
		Little Luckiamute R	Jont Cr	2
		Maxfield Cr	Price Cr	1.2
		Mctimmonds Cr	Pedee Cr	2.5
		Miller Cr	Beaver Cr	0.5
		Mouth	Soap Cr	2.5
		Pedee Cr	Ritner Cr	1
		Price Cr	Vincent Cr	2
		Ritner Cr	Maxfield Cr	3
		Rock Pit Cr	Wolf Cr	0.5

Table 6.7.2b. Protection status adopted by the Northwest Power Planning Council in 1988. Shown are status (1=Anadromous and Resident Fish or Wildlife; 2=Anadromous; and 3=Unprotected), stream, start reach and end reach and number of miles. Source was Streamnet.org, December 2003.

Status	Stream	Start Reach	End Reach	Reach Length (Mi)
		Soap Cr	Little Luckiamute R	15.5
		Vincent Cr	Bonner Cr	1.4
		Wolf Cr	Miller Cr	0.4
2	Maxfield Cr	Mouth	Headwaters	2.8
2	Pedee Cr	Mouth	Pedee Cr, S Fk	2.5
		Pedee Cr, S Fk	Pedee Cr, N Fk	1.7
2	Pedee Cr, N Fk	Mouth	Headwaters	2
2	Pedee Cr, S Fk	Mouth	Headwaters	2
2	Ritner Cr	Mouth	Clayton Cr	2.2
2	Soap Cr	Berry Cr	Baker Cr	1.6
		Mouth	Berry Cr	5.5
2	Teal Cr	Grant Cr	Headwaters	3.5
		Mouth	Grant Cr	1.5
2	Waymire Cr	Mouth	Headwaters	2
2	Willamette R	Ash Cr	Luckiamute R	12
		Luckiamute R	Santiam R	0.4
3	Ash Cr	Ash Cr, M Fk	Ash Cr, N Fk	0.1
		Mouth	Ash Cr, M Fk	1.2
3	Ash Cr, N Fk	Mouth	Headwaters	1.2
3	Ash Cr, S Fk	Mouth	Headwaters	1.7
3	Baker Cr	Mouth	Headwaters	0
3	Beaver Cr	Mouth	Headwaters	0
3	Berry Cr	Berry Cr, S Fk	Berry Cr, M Fk	0.5
		Mouth	Headwaters	0
			Peterson Cr	2
		Peterson Cr	Berry Cr, S Fk	2.5
3	Black Rock Cr	Mouth	Headwaters	0
3	Bonner Cr	Mouth	Headwaters	0
3	Boulder Cr	Mouth	Headwaters	0
3	Clayton Cr	Mouth	Headwaters	1
3	Cooper Cr	Mouth	Headwaters	0
3	Cougar Cr	Mouth	Headwaters	0
3	Dutch Cr	Mouth	Headwaters	0
3	Farley Cr	Mouth	Headwaters	0
3	Fern Cr	Mouth	Headwaters	0
3	Grant Cr	Mouth	Headwaters	0
3	Jont Cr	Fuller Cr	Headwaters	0

Table 6.7.2b. Protection status adopted by the Northwest Power Planning Council in 1988. Shown are status (1=Anadromous and Resident Fish or Wildlife; 2=Anadromous; and 3=Unprotected), stream, start reach and end reach and number of miles. Source was Streamnet.org, December 2003.

Status	Stream	Start Reach	End Reach	Reach Length (Mi)
		Mouth	Fuller Cr	0
3	Little Luckiamute R	Black Rock Cr	Headwaters	3
3	Maxfield Cr	Mouth	Headwaters	4.7
3	Mctimmonds Cr	Mouth	Headwaters	0
3	Miller Cr	Mouth	Headwaters	0
3	Pedee Cr	Pedee Cr, N Fk	Headwaters	2.7
3	Pedee Cr, N Fk	Mouth	Headwaters	2.5
3	Pedee Cr, S Fk	Mouth	Headwaters	1.5
3	Price Cr	Mouth	Woods Cr	5.2
		Woods Cr	Headwaters	6
3	Ritner Cr	Clayton Cr	Sheythe Cr	1
		Sheythe Cr	Headwaters	3.8
3	Rock Pit Cr	Mouth	Headwaters	0
3	Sams Cr	Mouth	Headwaters	0
3	Sheythe Cr	Mouth	Headwaters	4
3	Soap Cr	Baker Cr	Headwaters	10.5
		Berry Cr	Baker Cr	8.1
3	Teal Cr	Grant Cr	Headwaters	2.4
3	Vincent Cr	Mouth	Headwaters	4
3	Waymire Cr	Mouth	Headwaters	1.5
3	Wolf Cr	Mouth	Headwaters	0
3	Woods Cr	Mouth	Headwaters	5

6.7.3 NATURAL BARRIERS: RAPIDS, FALLS

There are a number of barriers documented in the Luckiamute system on StreamNet (www.streamnet.org). A reservoir acts as a barrier at mile 3 on Price Creek. There are unnamed cascades on Ritner Creek (rm 5), and the Little Luckiamute at river mile (rm) 20 and 21. A culvert acts as a barrier on the South Fork of Pedee Creek at rm 2. There are falls at rm 1 on Berry Creek, rm 0 on Grant Creek, rm 0 on the North Fork of Teal Creek, rm 0 on the South Fork of Teal Creek, at rm 13, 19 and 20 on the Little Luckiamute. The barrier on Teal Creek has isolated populations of cutthroat trout ((Licata *et al.*, 1998). We recommend that barriers be field check and observations recorded.

6.8 CHANNEL MODIFICATION ASSESSMENT

The only source of channel modification data were the AHI surveys. Since the AHI surveys covered so little of the study area, we did summarize the data. Channel modifications are visible from the ground in many places. We recommend that the LWC develop a channel modification survey to

collect this information. Ideas that have been used in other watersheds include a day where citizens and land owners walk streams along their property and record observations and take photographs. Results can be displayed in a public place, such as a library or school. Alternatively, aerial photographs can be reviewed and channel modifications can be recorded. Such an analysis was beyond the scope of this study.

6.9 INTERPRETATION AND RECOMMENDATIONS

We recommend conducting a survey of physical and behavioral barriers to fish passage. Culvert data should be collected from county, state and private timber groups. Information describing the diameter, drop of outfall, pool below, gradient, road condition, and ditch conditions should be organized for each culvert. Consider following ODFW and ODF culvert survey guidelines if new surveys are to be conducted.

When developing a water quality monitoring plan, consider stream reaches where high water temperature and low dissolved oxygen concentrations may act as a fish barrier.

- Set up a systematic water quality monitoring program with strategically located sample stations. Set up a monitoring program to answer specific questions and to develop baseline information. For example, sample in areas known to have spawning salmon, downstream of rare species management areas, urban areas, and intensively managed forests. Know how the data will be used before they are collected.
- Expand continuous stream temperature monitoring and using collected data in a stream temperature model (GIS-based) to interpolate (spatially) between sampling points.
- Establish stream gauging stations, weather stations and rainfall gages to improve knowledge of water availability.
- Participate in the TMDL process
- Evaluate first order streams for riparian condition (shade) and potential for LWD delivery.
- Locate areas where there are buildings in the riparian zone and determine if monitoring is warranted at that location; educate landowner about water quality.
- Verify points of water diversions.
- Map (or verify) spring and well locations.
- Document areas of ground water shortages and water quality problems from well logs.
- Begin to gather information on the location of the water table. Subsurface water flow entering streams may help to maintain cool water temperatures necessary for good salmonid habitat.
- Map locations of potential water contamination sources, *i.e.*, underground storage tanks and agricultural chemical storage areas.
- Prevent livestock from entering streams

7 SALMONID HABITAT ANALYSIS

7.1.1 DISTRIBUTIONS

A number of factors can regulate and control the distribution of organisms in their habitats. The science of ecology is concerned with identifying the factors that control and regulate the abundance and distribution of organisms. At any one time there is exactly one factor that limits the growth of a biological population. This is called the limiting factor. Different species have different factors that limit their populations and a single population can be limited by different factors at different times. It is difficult to actually measure limiting factors in nature; however, an understanding of the potentially limiting factors is absolutely essential to successfully manage biological resources.

Consider salmon. Anadromous salmonids spend part of the life cycle in streams, estuaries, and the sea. Therefore, a salmonid's habitat consists of stream, estuaries and the ocean **plus** all the factors that structure stream, estuary and oceanic ecosystems. In addition to physical habitat constraints, biological interactions between fish and between fish and other organisms, including disease causing organisms, can also affect the distribution of salmon. It is the complex life history and all of the possible interactions between salmon and their fresh-brackish-salt water environments, and between salmon and other organisms, which make the management and recovery of depressed salmon populations such a difficult problem.

Ideally, natural resource managers would identify the factor that limits salmon population growth and supply more of that limiting factor. For example, large wood is added to streams because natural resource managers believe that salmon populations are limited by a lack of in-stream complexity. Adding more wood, the suspected limiting factor, should result in more salmon. It is true that current in-stream complexity is dramatically different than historic accounts (this is supported by information presented in Sections 4.4.2.3 & 4.4.3); however, it is unlikely that the same factor (in this case, in-stream complexity) limits all salmon populations. For this reason, we recommend that LWC develop a diverse suite of restoration strategies and that they work with fisheries managers to develop the scientific underpinnings of the factors that truly limit salmonid populations.

A first step in developing an understanding of salmon habitat requirements is to develop a list of the factors that are believed to control and regulate salmon distribution in the Luckiamute/ Ash Creek study area. We searched for information describing the current distribution of salmon species in the study area. During a July 2003 meeting with the LWC technical team, a short list was developed of the most important factors (according to LWC). This list included: shade, stream gradient, pool & riffle information, temperature, flows, substrate, nutrients, carbon, and stream complexity.

7.1.1.1 Dataset Discussion

The first step to understanding what controls the abundance and distribution of any population is to know its current distribution.

ODFW Distribution Maps and Salmonid Core Areas

According to the Oregon Plan, "Core Areas are reaches or watersheds within individual coastal basins that are judged to be of critical importance to the sustenance of salmon populations that

inhabit those basins. Core Areas contain habitat needed to sustain populations. Furthermore, Core Areas provide a source for repopulating habitats as restoration programs are implemented.” Core areas were identified by a Scientific Panel assembled to create and review the Oregon Plan. Therefore, these areas are based on their best professional judgment. Core areas should be considered high priority areas for watershed protection and enhancement activities.

Since core areas were established for coastal populations of salmon, there are no core areas in the Luckiamute / Ash Creek study area. The mapping of core areas built on previous efforts undertaken by groups including: FEMAT Key Watersheds (selected by federal biologists as part of the Presidents Forest Plan; all located on federal lands); AFS Aquatic Diversity Areas (selected by committee of members from the Oregon Chapter of the American Fisheries Society), and DSL Essential Salmonid Habitat; and ODFW Source Watersheds (both selected by Oregon Department of Fish and Wildlife).

ODF Fish Limits Maps

As a part of its role in regulating timber harvest activities on Oregon lands, the Oregon Department of Forestry maintains maps of fish use in streams. These maps show the known or estimated upstream limits of game fish presence in many coastal streams. All game fish are considered, including resident cutthroat trout.

7.1.2 MULTI-FACTOR ANALYSES OF SALMONID HABITAT

We used a multi-factor approach to examine patterns in stream conditions within the Luckiamute / Ash Creek study area. Geographic information systems are ideal tools for simultaneously querying multiple spatial data sets to answer specific questions about the watershed. It is important to keep in mind that GIS is simply a data manipulation tool. Like any other tool, it can be used or abused: results must always be viewed critically. Results of the multi-factor analysis are predominantly affected by the quality of each data set used in the analysis.

As mentioned in earlier sections of this report, salmon populations are influenced by a wide variety of factors, both inside and outside of the Luckiamute and Ash Creek watershed study area, and operating at multiple spatial and temporal scales. Understanding the factors that control the abundance and distribution of salmon populations is a complex problem. Nevertheless, general benchmarks have been established for physical and biological factors by ODFW and others for instream habitat (Table 7.1.2a).

At the request of the LWC, we evaluated existing AHI data for the Luckiamute / Ash Creek study areas using two different approaches. In the first approach, we evaluated each of the 12 - 7th field watersheds according to criteria established by ODFW. We were limited by available data, *i.e.*, those stream reaches that were surveyed by AHI field teams. Each 7th field watershed was evaluated separately using several of the ODFW benchmarks selected from Table 7.1.2a. Recall, that only 12.2% of the streams in the study area were surveyed by AHI teams (see Section 6.1.1.2). In all cases, AHI survey extents did not cover the entire stream (1:100K) network in any individual 7th field watershed. In addition, stream data were not recorded for all stream reaches by field teams. Therefore, the level of sampling effort was not standardized between 7th field watersheds.

Nonetheless, general habitat quality patterns for the areas surveyed can be summarized. Since most of the Luckiamute / Ash Creek study area has not been surveyed, the following results should not be considered to be representative of the study area. Instead, we provide these summaries to demonstrate what could be done if appropriate data were available and to provide a starting place for action planning discussions.

Table 7.1.2a. Benchmarks recommended in the OWEB Watershed Assessment Manual (Watershed Professionals Network, 1999) are from the Oregon Department of Fish and Wildlife.		
Stream characteristic	Undesirable	Desirable
Pools		
Pool area (percent of total stream area)	<10	>35
Distance between pools (# of channel widths)	>20	5-8
Residual pool depth (meters)		
Small streams (<7m width)	<0.2	>0.5
Medium streams (≥7m & <15m width)		
Low gradient (slope <3%)	<0.3	>0.6
High gradient (slope >3%)	<0.5	>1.0
Large streams (≥15m width)	<0.8	>1.5
Complex pools/km (pools w/wood complexity>3)	<1.0	>2.5
Riffles		
Width:Depth ratio (Western Oregon)	>30	<15
Substrate		
Gravel substrate (% area)	<15	>35
Silt+sand+organic substrates (combined % area)		
Volcanic parent material	>15	<8
Sedimentary parent material	>20	<10
Channel gradient <1.5%	>25	<12
Shade		
Shade (reach average %)		
Stream width <12m (western Oregon)	<60	>70
Stream width >12m (western Oregon)	<50	>60
Woody debris		
Large woody debris (15cm X 3m minimum size)		
# of pieces/100m stream length	<10	>20
Volume/100m stream length (cubic m)	<20	>30
'Key' pieces (>60cm X 10m) per 100m stream length	<1	>3
Riparian conifers		
Riparian conifers within 30m of stream		
Number >20in dbh/1000 ft stream length	<150	>300
Number >35in dbh/1000 ft stream length	<75	>200
Source: (Watershed Professionals Network, 1999)		

Table 7.1.2b summarizes the ODFW habitat benchmarks for pools in each of the 12 7th field watersheds. Of all the watersheds, only Lower Pedee Ck (#17090003060401) ranks high in terms of desirable pool characteristics although, like the other basins, none of its stream reaches meet ODFW benchmark criteria for distance between pools. This indicates that, for these areas, habitat quality could be improved by adding structural complexity to the streams that would increase pool formation.

Table 7.1.2b: Evaluation of in-stream aquatic habitat using ODFW habitat benchmarks for 12 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are total length (ft) and proportion of AHI surveyed stream for each 7th field HUC falling into undesirable (U) and desirable (D) categories for selected stream characteristics.

Watershed Name/ 7th Field HUC	Total Length (ft)	Pool area		Complex pools		Distance between pools	
		% U	% D	% U	% D	% U	% D
Upper Luckiamute 17090003060101	51462.2	14.7	55.7	100.0	0.0	14.7	0.0
Miller Creek 17090003060102	38635.4	27.4	72.6	100.0	0.0	49.0	0.0
Wolf Creek 17090003060201	39843.5	19.0	40.1	100.0	0.0	23.1	0.0
Cougar Creek 17090003060202	25845.1	28.3	7.0	100.0	0.0	20.2	0.0
Lower Pedee CK 17090003060401	38628.7	0.0	89.4	0.0	64.3	0.0	0.0
Upper Pedee CK 17090003060402	20139.6	38.1	0.2	11.9	42.4	49.3	0.0
Upper Little Luckiamute 17090003060601	59225.6	43.2	2.7	58.6	5.0	5.9	0.0
Black Rock CK 17090003060603	11405.3	0.0	0.0	100.0	0.0	0.0	0.0
Socialist Valley 17090003060701	14259.1	0.0	0.0	35.8	64.2	0.0	0.0
Upper Soap CK 17090003061101	12437.8	0.0	100.0	100.0	0.0	0.0	0.0
Middle Soap CK 17090003061102	19360.0	0.0	100.0	100.0	0.0	0.0	0.0
Rifle Range 17090003061103	7357.5	0.0	100.0	100.0	0.0	0.0	0.0

We also evaluated each of the 7th field watersheds for Large Wood using the ODFW habitat benchmarks for key pieces of large wood, number of large wood pieces and large wood volume (Table 16.1.2c). Results show that overall, large wood is scarce in the stream reaches surveyed. Only two of the watersheds have any stream reaches that meet the desired criteria for key pieces of large wood. Several of the watersheds have reaches that meet desirable benchmarks for the number of pieces and large wood volume, but over all the majority of habitat surveyed falls into the undesirable category. Finally, Upper Pedee Ck (7090003060402) stands out as having the highest proportion of stream reaches meeting the desirable criteria for large wood.

Once source of large woody debris is the riparian zone. We summarized the number of conifers and shade in the riparian zone. Unfortunately, none of surveyed stream reaches met the ODFW-benchmarks for riparian conifers most of the streams were unshaded by trees within 12m of the

stream edge (Table 7.1.2d). This indicates the restoration strategy developed by LWC should include riparian plantings as well as supplying wood from some other source.

We also evaluated stream gradient and substrate data from the AHI data set. Unlike previous examples, many of the 7th field watersheds had many stream reaches that met the benchmark for stream gradient and percent gravel substrate. Fewer watersheds met criteria for silt, sand and organics (Table 7.1.2.e). The Upper Little Luckiamute (17090003060601) and Socialist Valley (17090003060701) are among the top watersheds surveyed. Results suggest that stream gradient is suitable for salmon habitat and that instream structure may be needed to capture and sort gravel and sediments.

Table 7.1.2c: Evaluation of in-stream aquatic habitat using ODFW habitat benchmarks for 12 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are total length (ft) and proportion of AHI surveyed stream falling into undesirable (U) and desirable (D) categories for selected stream characteristics.

Watershed Name/ 7th Field HUC	LWD key pieces		Number of LWD pieces		LWD volume	
	% U	% D	% U	% D	% U	% D
Upper Luckiamute 17090003060101	95.8	0.0	45.6	0.0	45.6	18.9
Miller Creek 17090003060102	91.0	0.0	65.9	1.3	70.1	10.4
Wolf Creek 17090003060201	60.2	0.0	54.9	25.8	31.4	35.1
Cougar Creek 17090003060202	80.2	0.0	69.5	0.0	69.5	19.8
Lower Pedee CK 17090003060401	78.3	10.6	82.5	10.6	89.4	10.6
Upper Pedee CK 17090003060402	45.7	11.9	45.7	38.1	45.7	54.3
Upper Little Luckiamute 17090003060601	26.5	0.0	40.5	35.0	59.9	20.3
Black Rock CK 17090003060603	0.4	0.0	0.4	0.0	100.0	0.0
Socialist Valley 17090003060701	100.0	0.0	35.8	64.2	100.0	0.0
Upper Soap CK 17090003061101	100.0	0.0	100.0	0.0	100.0	0.0
Middle Soap CK 17090003061102	100.0	0.0	100.0	0.0	100.0	0.0
Rifle Range 17090003061103	100.0	0.0	100.0	0.0	100.0	0.0

Table 7.1.2d: Evaluation of in-stream aquatic habitat using ODFW habitat benchmarks for 12 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are total length (ft) and proportion of AHI surveyed stream falling into undesirable (U) and desirable (D) categories for selected stream characteristics

Watershed Name/ 7th Field HUC	Riparian conifers >20 in		Riparian conifers >35 in.		Shade <=12m stream width		Shade >12m stream width	
	U	D	U	D	U	D	U	D
Upper Luckiamute 17090003060101	100.0	0.0	100.0	0.0	0.0	0.0	0.0	74.9
Miller Creek 17090003060102	100.0	0.0	100.0	0.0	0.0	0.0	0.0	91.9
Wolf Creek 17090003060201	100.0	0.0	100.0	0.0	0.0	0.0	9.4	72.2
Cougar Creek 17090003060202	80.2	0.0	80.2	0.0	0.0	60.7	0.0	39.3
Lower Pedee CK 17090003060401	100.0	0.0	100.0	0.0	0.0	0.0	18.0	53.2
Upper Pedee CK 17090003060402	100.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0
Upper Little Luckiamute 17090003060601	80.7	0.0	100.0	0.0	0.0	0.0	0.0	94.1
Black Rock CK 17090003060603	100.0	0.0	100.0	0.0	0.0	0.0	0.0	99.6
Socialist Valley 17090003060701	100.0	0.0	100.0	0.0	0.0	0.0	0.0	64.2
Upper Soap CK 17090003061101	100.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0
Middle Soap CK 17090003061102	100.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0
Rifle Range 17090003061103	100.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0

Table 7.1.2e: Evaluation of in-stream aquatic habitat using ODFW habitat benchmarks for 12 7th field watersheds in the Luckiamute and Ash Creek study area. Shown are total length (ft) and proportion of AHI surveyed stream falling into undesirable (U) and desirable (D) categories for selected stream characteristics

Watershed Name/ 7th Field HUC	Gradient		Percent gravel substrate		Silt, sand, and organics	
	U	D	U	D	U	D
Upper Luckiamute 17090003060101	0.0	85.3	0.0	39.7	100.0	0.0
Miller Creek 17090003060102	0.0	100.0	0.0	47.5	100.0	0.0
Wolf Creek	0.0	81.0	17.4	26.1	77.1	22.9

Table 7.1.2e: Evaluation of in-stream aquatic habitat using ODFW habitat benchmarks for 12 7 th field watersheds in the Luckiamute and Ash Creek study area. Shown are total length (ft) and proportion of AHI surveyed stream falling into undesirable (U) and desirable (D) categories for selected stream characteristics						
Watershed Name/ 7th Field HUC	Gradient		Percent gravel substrate		Silt, sand, and organics	
	U	D	U	D	U	D
17090003060201						
Cougar Creek 17090003060202	0.0	79.8	20.1	0.0	26.3	73.7
Lower Pedee CK 17090003060401	0.0	100.0	0.0	21.5	100.0	0.0
Upper Pedee CK 17090003060402	0.0	100.0	0.0	0.0	100.0	0.0
Upper Little Luckiamute 17090003060601	0.0	96.7	0.0	23.6	2.7	94.0
Black Rock CK 17090003060603	0.0	100.0	0.0	0.4	0.0	100.0
Socialist Valley 17090003060701	0.0	100.0	0.0	35.8	0.0	100.0
Upper Soap CK 17090003061101	0.0	100.0	0.0	100.0	100.0	0.0
Middle Soap CK 17090003061102	0.0	100.0	0.0	100.0	100.0	0.0
Rifle Range 17090003061103	0.0	100.0	0.0	100.0	100.0	0.0

In the second example, we used criteria established for steelhead by Burke et al. (in prep) to rank stream reaches from the AHI data set. These criteria were developed from a more recent survey of the scientific literature review than the ODFW benchmarks presented in the OWEB manual. In this example, we used criteria developed for a particular life stage of steelhead salmon, spawning habitat. Since this analysis used the AHI data the same caveats apply. We urge caution in interpreting these results.

At the request of LWC, we evaluated ODFW-defined stream reaches within each of the 12 7th field watersheds which were surveyed by AHI field teams. Each stream reach was evaluated separately according the criteria listed below.

Substrate:

Fines (good ≤ 10%; fair 10%- 20%; poor >20%)

Gravel (good ≥30%; fair 15%-30%; poor < 15%)

Cobbles (good ≥10% and ≤ 30%; fair 30% - 60%; and poor < 10% or > 60%)

Pools:

Tailouts (good 40% - 60%; fair 20% -40%; and poor < 20% or > 60%)

Residual Depth (good $\geq 0.2\text{m}$ and poor = no pools)

Temperature:

Temperature (good 6-12.5°C; fair 4-6°C and 12.5-16°C; and <4°C or >16°C)

Table 7.1.2.f shows the results for each stream reach surveyed. Some of the summary information is expressed for each habitat unit (a subset of a stream reach according to the ODFW method) and some of the information is expressed for the entire stream reach. For variables that were recorded for each habitat unit, we evaluated each habitat unit according to the criteria proposed by Burke *et al.* (in prep). Numbers in Table 7.1.2f are the number of habitat units meeting the 'good', 'fair' and 'poor' criteria. For summaries that are expressed for the entire reach, we recorded 'good', 'fair' or 'poor' for the entire reach.

TABLE 7.1.2.f: Evaluation of Steelhead Spawning Habitat: Stream segments of AHI surveyed streams ranked according to criteria developed by (Burke *et al.*, in prep). Shown are stream reach ID, length (ft), and the number of stream segments ranked as 'good', 'fair', and 'poor' steelhead spawning habitat. Also shown are reach-level ranks for pool frequency, residual pool depth and maximum temperature.

Stream	Reach ID	Length (ft)	Fines ^A (# habitats/reach)			Gravel ^A (# habitats/reach)			Cobble ^A (# habitats/reach)			Pool Frequency ^B	Residual Pool Depth ^B	Temp. (max) ^B
			Good	Fair	Poor	Good	Fair	Poor	Good	Fair	Poor			
Little Luckiamute River	123287844891401	9153.06	14	1		5	9	1	8	7		Poor	Good	Fair
	123287844891402	5147.21	11	2		9	3	1	2	9	2	Fair	Good	Fair
	123287844891403	15257.89	30			8	13	9	24	4	2	Poor	Good	Good
	123287844891404	22705.54	56		2	19	32	7	14	41	3	Poor	Good	Good
	123287844891405	2969.34	23	3	2	26	2		11	16	1	Poor	Good	Fair
	123287844891406	1289.53	3			1	2			3		Poor	Good	Fair
	123287844891407	1380.23	8			6	2		3	5		Poor	Good	Good
	123287844891408	1956.46	12	2	1	15			8	6	1	Poor	Good	Fair
	123287844891409	1577.30										Fair	Good	Good
	123556944867701	2152.56	3	1		2	1	1	2	2		Poor	Good	Fair
	123556944867702	4125.89	13		2	3	6	6	6	5	4	Poor	Good	Fair
	123556944867703	1336.05	4				2	2	1	1	2	Poor	Good	Fair
	123556944867704	5407.97	11		2	7	2	4	10	2	1	Poor	Good	Good
	123556944867705	1935.38	8			4	1	3	5		3	Poor	Good	Fair
123564544867001	8495.66	21	1		18	2	2	14	1	7	Poor	Good	Fair	
Luckiamute River	123148044755901	2236.24	10			6	3	1	5	4	1	Poor	Good	Poor
	123148044755902	1796.71	2			1	1			2		Good	Good	Fair
	123148044755903	11653.14	21				11	10	5	3	13	Fair	Good	Poor
	123148044755904	7027.36	1					1	1			Fair	Good	Poor

TABLE 7.1.2.f: Evaluation of Steelhead Spawning Habitat: Stream segments of AHI surveyed streams ranked according to criteria developed by (Burke *et al.*, in prep). Shown are stream reach ID, length (ft), and the number of stream segments ranked as 'good', 'fair', and 'poor' steelhead spawning habitat. Also shown are reach-level ranks for pool frequency, residual pool depth and maximum temperature.

Stream	Reach ID	Length (ft)	Fines ^A (# habitats/reach)			Gravel ^A (# habitats/reach)			Cobble ^A (# habitats/reach)			Pool Frequency ^B	Residual Pool Depth ^B	Temp. (max) ^B
			Good	Fair	Poor	Good	Fair	Poor	Good	Fair	Poor			
Luckiamute River	123148044755905	8795.87	21			13	4	4	7	9	5	Good	Good	Poor
	123148044755906	15918.30	26			21	5		5	16	5	Fair	Good	Poor
	123148044755907	12878.55	12			11	1		8	3	1	Poor	Good	Fair
	123565844758201	2769.81	2			2			1	1		Poor	Good	Good
	123565844758202	2078.69										Poor	Good	Good
	123565844758203	5109.77										Poor	Good	Good
	123568644762701	2313.62										Poor	Good	Good
	123568644762702	3254.21	7			5	2		3	4		Poor	Good	Good
	123568644762703	3729.78	1		1	2				2		Good	Good	Good
	123568644762704	2111.11		1		1			1			Poor	Good	Good
	123585844772501	9351.79	8	2	5	11	3	1	9	5	1	Poor	Good	Good
	123589544776901	3985.03	19	2	2	17	1	5	1	1	21	Poor	Good	Poor
	123589544776902	4895.20	7			7			5		2	Poor	Good	Fair
	123593144793601	2379.57	25	7	3	32	1	2	27	7	1	Fair	Good	Fair
	123593144793602	7550.57	3	1		3	1		2	2		Poor	Good	Fair
123593544776601	11724.18	18			16	2		11	2	5	Poor	Good	Poor	
123593544776602	7526.66	2	1		2	1		2		1	Good	Good	Fair	
123596744788901	5398.63	1	5	4	7	1	2			10	Poor	Good	Fair	
123596744788902	2180.78	1		3	4					4	Good	Good	Fair	
Luckiamute River	123600444776901	2859.25	18	2		20			13	6	1	Good	Good	Fair
	123600444776902	7338.72			1	1			1			Poor	Good	Good
	123608244774201	2440.31	3		1	3	1		3	1		Poor	Good	Good
	123608244774202	824.09										Poor	Poor	Good

TABLE 7.1.2.f: Evaluation of Steelhead Spawning Habitat: Stream segments of AHI surveyed streams ranked according to criteria developed by (Burke *et al.*, in prep). Shown are stream reach ID, length (ft), and the number of stream segments ranked as 'good', 'fair', and 'poor' steelhead spawning habitat. Also shown are reach-level ranks for pool frequency, residual pool depth and maximum temperature.

Stream	Reach ID	Length (ft)	Fines ^A (# habitats/reach)			Gravel ^A (# habitats/reach)			Cobble ^A (# habitats/reach)			Pool Frequency ^B	Residual Pool Depth ^B	Temp. (max) ^B
			Good	Fair	Poor	Good	Fair	Poor	Good	Fair	Poor			
	123610444765001	1643.38	3	2	1	6			3		3	Fair	Good	Good
	123624444790701	520.92	2	2		4			2		2	Poor	Good	Fair
	123624444790702	3494.09	6	4		9	1				10	Poor	Good	Fair
Pedee Creek	123432744740001	2665.45	4	5	2	10	1		5		6	Good	Good	Fair
	123432744740002	5654.95	15	7	8	26	2	2	3	1	26	Poor	Good	Fair
	123432744740003	6952.29	12	12	2	25	1		9	10	7	Poor	Good	Fair
	123447844771101	5758.87	14			2	11	1	1	12	1	Fair	Good	Fair
	123447844771102	4269.30	2			2			1	1		Fair	Good	Fair
	123447844771103	4102.77	4	2	3	5	3	1	7		2	Poor	Good	Good
	123447844771201	5473.68	12	6	1	13	4	2	9	7	3	Poor	Good	Good
	123447844771202	3783.81	14			1	12	1	5	8	1	Good	Good	Fair
	123447844771203	7409.76	22	6		24	2	2	13	11	4	Fair	Good	Fair
	123447844771204	2507.99										Poor	Good	Good
	123449244794201	1768.12	9			1	5	3	3	3	3	Poor	Good	Fair
	123449244794202	3264.68	9	1		9	1		4		6	Poor	Good	Poor
	123449244794203	2767.56	6	2	3	7	2	2	7	1	3	Poor	Good	Fair
123449244794204	2389.07										Poor	Good	Poor	
Soap Creek	123163044730501	33934.21	169	9	8	174	5	7	21	1	164	Poor	Good	Poor
	123163044730502	5221.11	46	2		48			39	3	6	Good	Good	Poor

^A= data summarized from habitat unit AHI data; ^B=data summarized from reach-level AHI data

In order to combine these various statistics into one value, we developed a numeric index. The Index in the following manner: for criteria that were expressed as the number of habitat units per stream reach, we multiplied 'good' values by 2 and 'fair' values by 1 and 'poor' values by 0; and for criteria that we expressed at the reach level, we added 10 for 'good', 5 for 'fair' and 0 for 'poor'. These values were added for each stream reach. This resulted in a single number that indicates the over all condition of the stream reach. Since reaches vary in length, the number is not scaled to the area surveyed: it is intended to show the relative rank of each stream reach in comparison with the others. The higher the number the better the conditions are, overall.

Table 7.1.2g identifies the 'best' stream reaches for each of the major river basins.

Table 7.1.2g. Ranked list of ODFW-defined stream reaches for each of the major river basins for which AHL data were available. Shown are stream reach ID and rank score. See text for a description of how rank scores were assigned			
ODFW Stream Reach	score	ODFW Stream Reach	score
Luckiamute River		Little Luckiamute River	
123593144793601	198	123564544867001	271
123148044755906	135	123556944867703	161
123600444776901	125	123287844891403	156
123148044755905	105	123287844891406	125
123593544776601	104	123287844891401	93
123589544776901	88	123287844891407	86
123585844772501	86	123556944867704	80
123148044755907	81	123556944867701	73
123148044755903	76	123287844891405	70
123148044755901	59	123287844891408	61
123568644762702	56	123287844891409	50
123589544776902	53	123556944867705	33
123624444790702	50	123556944867702	28
123610444765001	46	123287844891404	28
123608244774201	40	123287844891402	20
123596744788901	37	Pedee Creek	
123593144793602	35	123447844771203	152
123624444790701	33	123432744740003	130
123565844758201	31	123432744740002	113
123593544776602	29	123447844771201	105
123568644762703	28	123447844771202	75
123568644762704	25	123447844771101	72
123596744788902	25	123449244794203	60
123148044755902	24	123432744740001	59
123600444776902	24	123447844771103	57
123565844758202	20	123449244794202	56
123565844758203	20	123449244794201	49

123568644762701	20	123447844771102	26
123608244774202	20	123447844771204	20
123148044755904	14	123449244794204	10
		Soap Creek	
		123163044730501	753
		123163044730502	281

7.1.3 MULTI-FACTOR SALMONID HABITAT ANALYSES: SYNTHESIS

In general, the factors that control the abundance and distribution of salmonids in the Luckiamute / Ash Creek watersheds are not well know. The distribution of salmonid species is known in a broad sense. Although there is a sense of the factors that may affect the distribution and abundance of salmonids in the study area, data are lacking.

For the areas surveyed, in-stream structure is lacking. Evidence for this comes from the lack of pool complexity, large woody debris, and substrates. There is also evidence that riparian shade (and potential for large wood to enter streams) is of concern. From the previous section, we saw that current salmonid distributions coincide with water quality limited streams. Nevertheless, we did identify 7th field watersheds and stream reaches where one or more environmental variables were found to be desirable.

Additional data is needed to better manage salmon in the Luckiamute/ Ash Creek study area. We recommend that fish populations be measured to get quantitative population measurements. From population number, the success of restoration actions can be evaluated. We recommend that benchmarks or reference conditions be established from areas within the watershed identified in this section or from nearby areas. Future restoration actions can be modeled after these reference sites. We recognize that some of the watersheds/ stream reaches identified as good habitat may not be accessible to salmon because of barriers. These areas, however, can still be used as areas to illustrate the desirable condition of the ODFW benchmarks.

7.2 INTERPRETATION AND RECOMMENDATIONS

- Map active floodplains and wetland areas. Collect data from landowners on flood frequency, areas of inundation, alternate stream channels and backwater wetlands.
- Map areas of dynamic (frequently changing) stream channels.
- Map locations of channel modifications.
- Map locations where streams are entrenched.
- Map locations of exposed bedrock along streams.
- Map locations of algal blooms, indicators of nutrient enrichment and low dissolved oxygen concentration.

- Use the results of this report to prioritize areas in which AHL surveys need to be conducted or updated. To improve spatial accuracy of AHL surveys, measure habitat unit lengths with hip chains from landmarks that are visible on the DOQ photographs or the USGS topographic base maps. Use GPS if possible. Calibrate observers to maximize spatial accuracy. Ensure that data are quickly processed and incorporated into the MCWC GIS at an appropriate spatial scale.
- Map the locations of exotic plants.
- Map the locations of beaver dams. Review the AHL data for locations of beaver dams and beaver activity (in the AHL comment columns). Consider beaver dam locations when planning riparian plantings, especially conifers.
- Work with ODFW and others to develop reliable estimates of the populations and distribution (including fish limit maps) for species of concern, such as salmon, lamprey, and mussels. Volunteers can be used to expand agency surveys provided that established protocols are followed. The lack of data on the distribution and abundance of aquatic organisms is a major impediment to developing a successful watershed enhancement strategy.
- Design data collection strategies that include biological sampling. For example, water quality monitoring data should include sampling for benthic macroinvertebrates, which can be good indicators of water quality and environmental change.

8 SURFACE EROSION

8.1.1 SOILS

Under some circumstances, soils can move across the landscape and into the stream network where suspended sediments can dramatically affect the quality of salmonid habitat. Circumstances that foster soil erosion include any actions that remove vegetation (which acts to stabilize soils), or any actions that lead to an increased frequency of mass wasting events. Detailed information on soils provided here can be used to plan actions to minimize the effect on soils prone to erosion.

8.1.2 EROSION BY LAND USE

We obtained soils information for each county in the Luckiamute/ Ash Creek study area. Then we identified erodible soil units, downloaded Map Unit Interpretation Records (MUIR) from NRCS or OSU, and assigned attributes to the soils coverage. We divided the study area into forested and non-forested areas using the EC90 Land cover layer. We used the wind erosion parameter from the SSURGO data set ('wderosn') to indicate the risk of forested area soils to erosion (except for tree farms). We used the 'K-factor' from the SSURGO data set to group agricultural soils into low (<0.20), moderate (0.20 to 0.40), and high (>0.40) soil erosion risk classes as specified in the OWEB manual (Watershed Professionals Network, 1999). We did not split soil polygons between forested and non-forested areas, rather, we assigned the class to which ever land cover class occurred at the center of each polygon. We did this because there was not a decisive dividing line between forested and non-forested areas. We did encounter some soil polygons in the forested areas that did not have a value recorded for 'wderosn.' In these cases, the K-factor values were used as a substitute.

See information in Section 4.4.5.3 on gravel pits and quarries, which can act as sediment sources.

8.1.3 ROADS

Roads can contribute sediments to stream especially improperly constructed or abandoned roads. The LWC was interested in learning the status of abandoned in the watershed. We did acquire and use a roads layer from the BLM; however, BLM personnel recommended that road attribute data not be used because of its low accuracy (Michelle Davis, pers. comm. 7/23/2003). We also checked with the timber companies but could not get their road data. The county data sets do have improved and unimproved data in them, but the counties have their data at different levels of detail (Polk is more detailed), so the data sets were not combined for this analysis. Therefore, we were not able to look at the impact of abandoned and unimproved roads on sediment loading.

In addition to unimproved or abandoned roads, roads passing over areas of steep slopes can fail, or act as chronic sources of sediments to streams. We ranked 7th field watersheds by the length of roads passing near streams.

We used the DEM and roads layers to rank 7th field watersheds by the length of roads that each had passing within 200 ft of a stream. Roads passing over or near streams can contribute sediments directly to streams during rain events.

Table 8.1.3.1. List of 7 th field watersheds ranked by the length of road passing within 200 ft of a stream.		
Watershed ID	Name	Total Length of Road (ft) within 200ft of a stream
3061101	Upper Soap Creek	113,978.2
3060304	Maxfield Creek	106,170.8
3060203	Hoskins	84,936.0
3060204	Vincent Creek	72,199.9
3060401	Lower Pedee Creek	69,855.1
3060701	Socialist Valley	68,001.0
3060102	Miller Creek	65,614.4
3060802	Lower Teal Creek	65,183.6
3060305	Bump Creek	62,936.9
3060404	Ritner Creek	62,158.1
3060303	Price Creek	58,915.8
3060101	Upper Luckiamute	57,582.7
3060603	Black Rock Creek	55,388.9
3060502	McTimmonds	55,255.7
3060402	Upper Pedee Creek	52,405.8
3060201	Wolf Creek	47,125.0
3060504	Jont Creek	43,647.2
3060901	Fern Creek	41,660.3
3060702	Falls City	38,907.4

Table 8.1.3.1. List of 7 th field watersheds ranked by the length of road passing within 200 ft of a stream.		
Watershed ID	Name	Total Length of Road (ft) within 200ft of a stream
3060301	Plunkett Creek	38,142.0
7020602	Middle North Fork Ash Creek	37,835.7
3060202	Cougar Creek	36,376.8
3060302	Woods Creek	35,914.8
3060503	Middle Luckiamute	35,265.7
3061201	Upper Berry Creek	33,252.8
7020605	Upper South Fork Ash Creek	32,907.8
3061102	Middle Soap Creek	30,121.2
7020601	Upper North Fork Ash Creek	29,327.7
3060601	Upper Little Luckiamute	29,057.7
3060903	Cooper Creek	28,928.1
3060704	Bridgeport	28,164.1
3061203	Peterson Creek	27,155.2
3060403	Clayton Creek	26,025.3
3060703	Waymire Creek	25,868.7
7020604	Middle Fork Ash Creek	23,610.2
3060501	Ira Hooker	21,961.6
3060803	Grant Creek	21,139.0
3060801	Upper Teal Creek	20,012.4
3061302	Palestine	19,862.1
3060902	Lower Little Luckiamute	19,218.6
3060602	Cold Springs	18,160.9
3061202	Staats Creek	14,243.8
3061303	Springhill	14,073.6
3061103	Rifle Range	10,773.3
3061004	Parker	9,678.7
3061005	Luckiamute Landing	9,292.9
3061003	Helmick	8,958.6
7020603	Lower North Fork Ash Creek	8,918.4
7020606	Lower South Fork Ash Creek	7,249.8
7020503	Harman Slough	6,728.7
7020502	Duck Slough	6,500.3
3061301	E.E. Wilson	6,215.5
3061001	Simpson	5,929.0
3061204	Lower Berry Creek	3,547.6
3061002	Zumwalt	3,384.3
7020404	Buena Vista	3,170.6

8.2 STREAMBANK EROSION

Stream bank erosion can be a significant source of sediments entering streams. Bank erosion can cause sediment loading, which can cover gravel beds and make them unsuitable for salmonid spawning. Excessive fine sediments may also reduce the quality of in-stream habitat for other species such as lamprey, freshwater mussels and macroinvertebrates. The sediment input from streambank erosion can also provide gravel, which is needed for salmon spawning beds.

Observations made by Aquatic Habitat Inventory crews concerning actively eroding stream banks are recorded. We queried the AHI database and retrieved the following records (Table 8.2.1). Keep in mind that the AHI data coverage in the study was limited and dated. These data should not be considered representative of the study area. We recommend that the LWC map areas where stream banks are eroding as part of its future monitoring program.

Table 8.2.1. Stream reaches with actively eroding banks as identified by AHI crews.

Stream Reach	Stream	Reach Number	Percent Bank Erosion	Length (ft)
123148044755901	LUCKIAMUTE RIVER	1	5.3	2236.2
123148044755902	LUCKIAMUTE RIVER	2	0.0	1796.7
123148044755903	LUCKIAMUTE RIVER	3	4.2	11653.1
123148044755904	LUCKIAMUTE RIVER	4	0.8	7027.4
123148044755905	LUCKIAMUTE RIVER	5	14.1	8795.9
123148044755906	LUCKIAMUTE RIVER	6	19.7	15918.3
123148044755907	LUCKIAMUTE RIVER	7	8.9	12878.6
123163044730501	SOAP CREEK	1	67.3	33934.2
123163044730502	SOAP CREEK	2	27.8	5221.1
123287844891401	LITTLE LUCKIAMUTE RIVER	1	8.2	9153.1
123287844891402	LITTLE LUCKIAMUTE RIVER	2	18.5	5147.2
123287844891403	LITTLE LUCKIAMUTE RIVER	3	10.0	15257.9
123287844891404	LITTLE LUCKIAMUTE RIVER	4	4.3	22705.5
123287844891405	LITTLE LUCKIAMUTE RIVER	5	34.4	2969.3
123287844891406	LITTLE LUCKIAMUTE RIVER	6	10.2	1289.5
123287844891407	LITTLE LUCKIAMUTE RIVER	7	18.4	1380.2
123287844891408	LITTLE LUCKIAMUTE RIVER	8	71.2	1956.5
123287844891409	LITTLE LUCKIAMUTE RIVER	9	22.5	1577.3
123432744740001	PEDEE CREEK	1	25.0	2665.5
123432744740002	PEDEE CREEK	2	28.0	5655.0
123432744740003	PEDEE CREEK	3	26.0	6952.3
123447844771101	SOUTH FORK PEDEE CREEK	1	14.0	5758.9
123447844771102	SOUTH FORK PEDEE CREEK	2	13.0	4269.3
123447844771103	SOUTH FORK PEDEE CREEK	3	4.0	4102.8
123447844771201	NORTH FORK PEDEE CREEK	1	25.0	5473.7

Table 8.2.1. Stream reaches with actively eroding banks as identified by AHI crews.				
Stream Reach	Stream	Reach Number	Percent Bank Erosion	Length (ft)
123447844771202	NORTH FORK PEDEE CREEK	2	8.0	3783.8
123447844771203	NORTH FORK PEDEE CREEK	3	26.0	7409.8
123447844771204	NORTH FORK PEDEE CREEK	4	20.0	2508.
123449244794201	PEDEE CREEK TRIBUTARY	1	5.0	1768.1
123449244794202	PEDEE CREEK TRIBUTARY	2	22.0	3264.7
123449244794203	PEDEE CREEK TRIBUTARY	3	12.0	2767.6
123449244794204	PEDEE CREEK TRIBUTARY	4	12.0	2389.1
123556944867701	CAMP CREEK	1	1.6	2152.6
123556944867702	CAMP CREEK	2	5.0	4125.9
123556944867703	CAMP CREEK	3	0.0	1336.1
123556944867704	CAMP CREEK	4	0.0	5408.0
123556944867705	CAMP CREEK	5	0.3	1935.4
123564544867001	LOST CREEK	1	34.6	8495.7
123565844758201	COUGAR CREEK	1	27.7	2769.8
123565844758202	COUGAR CREEK	2	16.9	2078.7
123565844758203	COUGAR CREEK	3	16.0	5109.8
123568644762701	SLICK CREEK	1	1.2	2313.6
123568644762702	SLICK CREEK	2	5.0	3254.2
123568644762703	SLICK CREEK	3	7.2	3729.8
123568644762704	SLICK CREEK	4	1.4	2111.1
123585844772501	ROCK PIT CREEK	1	11.8	9351.8
123589544776901	WOLF CREEK	1	71.5	3985.0
123589544776902	WOLF CREEK	2	47.8	4895.2
123593144793601	BOULDER CREEK	1	31.7	2379.6
123593144793602	BOULDER CREEK	2	2.5	7550.6
123593544776601	LUCKIAMUTE RIVER TRIBUTARY 1	1	20.7	11724.2
123593544776602	LUCKIAMUTE RIVER TRIBUTARY 1	2	0.9	7526.7
123596744788901	BEAVER CREEK	1	7.2	5398.6
123596744788902	BEAVER CREEK	2	0.8	2180.8
123600444776901	MILLER CREEK	1	15.6	2859.3
123600444776902	MILLER CREEK	2	2.2	7338.7
123608244774201	MILLER CREEK TRIBUTARY 1	1	18.0	2440.3
123608244774202	MILLER CREEK TRIBUTARY 1	2	9.8	824.1
123610444765001	MILLER CREEK TRIBUTARY 2	1	16.3	1643.4
123624444790701	LUCKIAMUTE RIVER TRIBUTARY 2	1	6.7	520.9
123624444790702	LUCKIAMUTE RIVER TRIBUTARY 2	2	37.3	3494.1

8.3 LANDSLIDES

Areas of the Luckiamute / Ash Creek study area are very dynamic especially in the Coast Range. Steep slopes and high amounts of precipitation are generally responsible for mass wasting (e.g., landslides and debris torrents) events throughout the region. Even the earliest accounts of the region's explorers describe large areas of landslides and debris torrents. Thus, Oregon's Coast Range has been susceptible to mass wasting prior to the time of European settlement. Mass wasting is a natural process; it is the frequency and magnitude of events that are of concern. Many factors can contribute to an increased frequency of mass wasting events including, land use practices, road building, development, etc.

Mass wasting adds sediments (both fine and coarse) and organic material to the stream network. These natural stream components are neither good nor bad in themselves; it is the frequency, magnitude and duration of mass wasting events that may have undesired consequences on in-stream conditions, especially on salmonid habitat. After all, organisms like Pacific Northwest salmonids have evolved in these rapidly changing landscapes and they are adapted to the 'natural' (background) patterns of mass wasting.

8.3.1 ODF DEBRIS FLOW HAZARD MAPS

According to information available on the ODF web site, Western Oregon Debris Hazard Maps were prepared to depict areas that are subject to naturally occurring debris flows. They include initiation sites and paths of potential debris flows. These are coarse scale risk assessment maps and should not be used without on-the-ground verification. These maps were developed from the 30-m DEM and lithology data layers. Streams were represented by USGS digital raster graphic data. These maps were also developed using available historic information on debris flow from a variety of sources (e.g., ODF, USFS, DOGAMI, BLM and ODOT). These maps did not account for patterns in rainfall.

Briefly, the ODF debris hazard maps assign a risk category to 2-4 acres parcels based on steepness and lithology. Steep areas that occur on Tye (and similar) geologic formations are rated higher (i.e., having a higher chance of sliding). Past landslide occurrence in an area resulted in a higher risk category being assigned to that area. ODF plans to develop additional guidance based on this work.

We summarized the areas of landslide risk from the ODF debris flow hazard maps for our study area (Table 8.3.1).

Table 8.3.1. List of ranked 7 th field watersheds according to debris flow hazard. Shown are the watershed ID, name, the number of acres in the high and moderate risk categories (defined by ODF) and the total acres.				
Watershed ID	Name	High	Moderate	Grand Total
3060202	Cougar Creek	719.3	3,368.6	4,087.9
3060101	Upper Luckiamute	244.7	3,258.6	3,503.3
3060102	Miller Creek	156.0	3,056.8	3,212.8
3060601	Upper Little Luckiamute	228.2	2,444.7	2,672.8
3060304	Maxfield Creek	81.9	2,435.8	2,517.7

Table 8.3.1. List of ranked 7th field watersheds according to debris flow hazard. Shown are the watershed ID, name, the number of acres in the high and moderate risk categories (defined by ODF) and the total acres.

Watershed ID	Name	High	Moderate	Grand Total
3060203	Hoskins	268.0	2,231.0	2,499.0
3061101	Upper Soap Creek	115.2	2,321.2	2,436.3
3060404	Ritner Creek	109.2	2,325.3	2,434.5
3060402	Upper Pedee Creek	169.2	2,165.4	2,334.6
3060201	Wolf Creek	513.7	1,763.0	2,276.7
3060303	Price Creek	110.8	1,532.1	1,642.9
3060701	Socialist Valley	106.9	1,494.0	1,601.0
3060602	Cold Springs	287.9	1,268.8	1,556.7
3060302	Woods Creek	108.9	1,381.1	1,490.0
3060401	Lower Pedee Creek	52.5	1,293.4	1,345.8
3060603	Black Rock Creek	96.9	994.2	1,091.1
3060204	Vincent Creek	2.4	1,052.3	1,054.7
3060802	Lower Teal Creek	53.0	959.4	1,012.3
3061102	Middle Soap Creek		882.0	882.0
3060801	Upper Teal Creek	39.8	834.5	874.3
3060301	Plunkett Creek	39.5	805.8	845.3
3060502	McTimmonds		711.3	711.3
3060403	Clayton Creek	37.9	547.8	585.7
3060702	Falls City	0.2	564.5	564.7
3061201	Upper Berry Creek		553.6	553.6
3060501	Ira Hooker		454.8	454.8
3061103	Rifle Range		324.9	324.9
3060305	Bump Creek		322.4	322.4
3060504	Jont Creek		318.9	318.9
3060503	Middle Luckiamute		288.1	288.1
3060803	Grant Creek		262.8	262.8
3060703	Waymire Creek	0.4	165.0	165.4
3061301	E.E. Wilson		129.5	129.5
3061302	Palestine		90.2	90.2
3061005	Luckiamute Landing		50.0	50.0
3060704	Bridgeport		28.7	28.7
3061001	Simpson		24.5	24.5
7020601	Upper North Fork Ash Creek		21.4	21.4
7020604	Middle Fork Ash Creek		21.2	21.2
7020404	Buena Vista		20.7	20.7
3061004	Parker		20.6	20.6
3061003	Helmick		17.1	17.1
3061204	Lower Berry Creek		15.9	15.9
7020602	Middle North Fork Ash Creek		0.4	0.4
3060901	Fern Creek		0.2	0.2

8.4 INTERPRETATION AND RECOMMENDATIONS

- Acquire or develop a complete roads layer at a consistent spatial scale of 1:24,000 or better. Differentiate between paved and non-paved roads.
- Recommend using the Riparian layer to located specific places (buildings) and bare areas where erosion may be a problem.
- Fence areas where livestock can enter streams.
- Evaluate roads located in high sediment risk watersheds for sediment control, if necessary.
- Map road failures and the condition of roads that pass through riparian areas.
- Map roads that may confine streams.
- Map culvert locations and collect information on culvert features, including degree of blockage and if culverts are fish barriers. Use a standardized data sheet to collect this information, similar to the one prepared by ODFW.
- During or after heavy rainfall events, record locations where surface flow runs directly along roadways and into streams. These roads can be major sediment sources of streams.

8.5 DATA COLLECTION AND MONITORING RECOMMENDATIONS

Will be pulled from text and repeated here.

8.6 RECOMMENDATIONS FOR FUTURE ANALYSES

Will be pulled from text and repeated here.

8.7 WATERSHED ENHANCEMENT RECOMMENDATIONS

Will be pulled from text and repeated here.

8.7.1 ECOSYSTEM CONTEXT

Declining salmonid population trends have been apparent for several decades. During the past 60 years or so, natural resource managers have relied on fish harvest restrictions, hatcheries, and habitat enhancement/improvement as management tools to bolster moribund populations. The declines have continued. It is likely that our knowledge of the ecological process that maintain salmonid populations is incomplete. A watershed approach involves putting the valued resources, in this case, salmon, into an ecological perspective. This watershed assessment completes the first step in developing a watershed-based restoration plan. Using knowledge gained from this report, it may be possible to apply some of the old management tools in ways that are more effective, or to develop completely new tools.

In a recent workshop that focused on habitat restoration in the Columbia River, guidelines were established by estuarine ecologist Dan Bottom to assist managers in restoring Columbia River salmonid populations (Bottom, 2001). These guidelines are also applicable for estuaries and watersheds in the MidCoast region.

1. Knowledge of the ecosystem is incomplete; therefore, restoration actions are largely experimental. Careful attention must be paid to the evaluation of restoration actions (through experimentation or monitoring).
2. Selection of restoration/enhancement sites should be for ecological reasons, rather than simply opportunistic.
3. Identify and conserve (or preserve) the high quality habitat that currently exists.
4. Where restoration is necessary, select historically important, high quality habitats (e.g., wetlands) or areas known to be important for sensitive life history stages of salmon (e.g., oligohaline-brackish water transition zones).
5. Give priority to passive restoration, rather than highly engineering solutions (active restoration).
6. Finally, adopt a landscape / watershed perspective. Avoid unlinked 'parcel-based' or 'postage stamp' restoration projects.

In this assessment, we used GIS to examine patterns in the factors believed to affect salmonid populations. We recommend that the LWC use and refine this information to assist the council in locating and assessing all restoration projects.

The importance of relatively small, locally adapted populations of salmonids in stabilizing 'salmon runs' (i.e., decreasing fluctuations in those runs) is just beginning to be appreciated. Current thinking is that small populations of salmon are adapted to the unique local conditions of individual stream reaches. Environmental conditions like food availability, stream substrate, water temperature, flow (and flow pattern), etc. vary from stream reach to stream reach so that no two stream reaches are identical. The salmon that are best suited for conditions at a particular stream reach contribute more offspring to that local population than others do. Over time, the genetic makeup of that local population will become slightly different from the genetic makeup of nearby populations, even within the same watershed.

Historically, there were many small populations within a larger watershed, each adapted to its local conditions and varying slightly from one another. Differences may also have resulted in slightly different spawning and migration behaviors (timing). All of these populations existed within very dynamic coastal watersheds. From time to time, cataclysmic (i.e., landslides, floods, etc.) or biological (i.e., competition, disease) events would eliminate (or dramatically reduce) some of the locally adapted populations. However, if the change was not too great, other populations within the same watershed would not be affected. Consequently, the overall production of salmon for that watershed would remain fairly constant over time. In other words, the stability of the larger watershed's salmon population largely depended upon the diversity of the locally adapted populations.

Currently, genetic diversity in coastal salmonid populations is believed to be low. This may be due to the disappearance of many of the locally adapted populations or perhaps to environmental alterations that have all but eliminated environmental variability at the stream reach level. In the dynamic coastal environment, it is believed that the current salmon populations do not have the resiliency to quickly rebound after disturbance (natural or man-made) like the more diverse, historical populations. Consequently, salmon runs are observed to fluctuate widely. A successful restoration strategy will involve rebuilding the genetic diversity of salmonid populations. This involves creating and maintaining the conditions to which salmon populations can locally adapt.

- Consult the GIS to determine if areas for planned projects or land use changes have potentially high erosion and/or landslide risk. Consider scheduling actions that disturb vegetation in these areas for times of low precipitation to avoid disturbing soils. Plan on leaving wide vegetated buffer strips to trap eroding sediments.
- Consult the results of the analyses in this assessment when planning watershed enhancement activities. Use the stream-reach level data provided with this report (along with local knowledge and additional information like land ownership, landowner willingness, adjacent land use, existing anadromous migration barriers, etc.) to help pick specific sites for such actions.
- Supplement recommendations with local knowledge and more detailed site-specific information when choosing final locations for projects.

REFERENCES

- Adamus, P. R., B. Anderson and R. Garono. in press. Model-based predictions of habitat quality and terrestrial vertebrate species composition in 72 watersheds of the Willamette River Basin. Salem, Oregon, Report to the Technical Advisory Committee of the Willamette Restoration Initiative.
- Aikens, C. M. 1986. *Archeology of Oregon*, U.S. Department of Interior, Bureau of Land Management.
- Aikens, C. M. 1992. *Archeology of Oregon*. Portland, OR, U.S. Department of the Interior. Bureau of Land Management.
- Aikens, C. M. 1993. *Archeology of Oregon*. Portland, OR, U.S. Department of the Interior. Bureau of Land Management.
- Barnhart, H. E. 1915. Survey of the North Luckiamute Valley, Polk County, Oregon. Historical, Social, and Economic. Master of Arts. Eugene, OR, Oregon State University.
- Benson, B. 2003. General Manager, Valley Landfills. Corvallis, OR.
- Botkin, D. B., K. Cummins, T. Dunne, H. Regier, M. Sobel and L. M. Talbot. 1993. Status and Future of Anadromous Fish of Western Oregon and Northern California: Rationale for a New Approach. Santa Barbara, CA, The Center for the Study of the Environment.
- Bureau of Land Management. 1882. Cadastral Surveys. Cadastral Surveys of the Willamette Valley, Bureau of Land Management.
- Bureau of Land Management. 1991. Fire Occurrence (1980-1989), Oregon/Washington BLM. **2003**.
- Bureau of Land Management. 2003. Ground Transportation Roads and Trails, Bureau of Land Management. **2003**.
- Burke, J. L., K. K. Jones and J. M. Dambacher. in prep. HabRate: A Stream Habitat Evaluation Methodology for Assessing Potential Production of Salmon and Steelhead in the Middle Deschutes River Basin. Portland, OR, Oregon Department of Fish and Wildlife: 78.
- Clark, J. L. 1999. Effects of Urbanization on Streamflow in Three Basins in the Pacific Northwest. Department of Geology. Portland, OR, Portland State University.
- Compton, J. E., M. R. Church, S. T. Larned and W. E. Hogsett. In Press. "Nitrogen export from forested watersheds in the Oregon Coast Range: The role of N₂-fixing red alder." *Ecosystems*.
- Franklin, J. F. and C. T. Dyrness. 1988. *Natural Vegetation of Oregon and Washington*. Corvallis, OR, Oregon State University Press.
- Frantz, W. C. and C. Brandon. 1976. Timber up the Luckiamute. Monmouth, OR.
- Garono, R. and L. Brophy. 2001. MidCoast Sixth Field Watershed Assessment Final Report: Prepared for the MidCoast Watersheds Council, July 2001. Corvallis, OR, Earth Design Consultants, Inc., Green Point Consulting.
- Hynes, H. B. N. 1970. *The Ecology of Running Waters*. Liverpool, UK., Liverpool University Press.
- Independent Multidisciplinary Science Team. 1999. Defining and Evaluating Recovery of OCN Coho Salmon Stocks: Implications for rebuilding stocks under the Oregon Plan for Salmonids and Watersheds. Technical Report 1999-2 to the Oregon Plan for Salmon and Watersheds. Salem, OR, Governor's Natural Resources Office.
- Institute for a Sustainable Environment. 1999. Landuse and Landcover ca. 1990, Institute for a Sustainable Environment, University of Oregon GIS Research Lab. **2003**.
- Johnson, D. H. and T. A. O'Neil. 2001. *Wildlife-Habitat Relationships in Oregon and Washington*. Corvallis, OR, Oregon State University Press.

- Lee, K. K. and J. C. Risley. 2002. Estimates of Ground-Water Recharge, Base Flow, and Stream Reach Gains and Losses in the Willamette River Basin, Oregon. Portland, USGS: 52.
- Licata, G., T. Tomczyk, G. Humbard, A. Haynes, D. Bergen, R. Exeter, E. Frazier and P. Hawe. 1998. Rowel Creek/Mill Creek/Rickreall Creek/Luckiamute River Watershed Analysis. Salem, OR, Mary's Peak Resource Area, Salem District Bureau of Land Management.
- Long, C. J. and C. Whitlockm. 2002. "Fire and vegetation history from the coastal rain forest of the western Oregon coast range." *Quaternary Research* **58**(3): 215-225.
- Mattson, K. and A. Gallagher. 2001. Rickreall Watershed Assessment. Dallas, OR, Ecosystems Northwest.
- McArthur, L. A. 1992. *Oregon Geographic Names*. Portland, OR, Oregon Historical Society Press.
- Melancon, P. A. 1999. A GIS Based Watershed Analysis System for Tillamook Bay, Oregon. Engineering. Austin, TX, University of Texas (Austin): 361.
- Miller, D., K. Burnett and K. Christiansen. 2001. all4hastreams (CLAMS streams layer), Corvallis Forestry Sciences Laboratory- CLAMS project. **2003**.
- Moore, K., K. K. Jones and J. M. Dambacher. 2002. Methods for Stream Habitat Surveys, Aquatic Inventories Project, Natural Production Program: Oregon Department of Fish and Wildlife. Corvallis, OR, Oregon Department of Fish and Wildlife: 59.
- Newton, S. W. 1971. *Early History of Independence Oregon*. Salem, OR, Panther Printing Company.
- Omernik, J. M. 1987. "Ecoregions of the conterminous United States. Map (scale 1:7,500,000)." *Annals of the Association of American Geographers* **77**(1): 118-125.
- Oregon Climate Service. 2003. Oregon Climate Service, Oregon Climate Service, Oregon State University. **2003**.
- Oregon Department of Land Conservation and Development. 2003. Goals/Rules/Laws, Oregon Department of Land Conservation and Development. **2003**.
- Oregon State University Animal Sciences Department. 2003. Oregon State University Beef Ranches, Oregon State University Animal Sciences Department. **2003**.
- Oregon State University College of Forestry. 1990.
- Oregon Water Resources Department and Oregon Department of Land Conservation and Development. 2002. Ground Water Supplies In the Willamette Basin. 2002. Salem, OR, Oregon Water Resources Department
- Oregon Department of Land Conservation and Development: 53.
- Rogers, A. 2003. Cultural Resources Manager, Peavy Arboretum. Corvallis, Oregon. Pers. Comm.
- Ruby, R. H. and J. A. Brown. 1992. *A Guide to the Indian Tribes of the Pacific Northwest*. Norman and London, University of Oklahoma Press.
- Sexson, C. 2003. Soap Creek Beef Barns. Corvallis, Oregon. Pers. Comm.
- Simenstad, C. A., M. Dethier, C. Levings and D. Hay. 1997. The Land-Margin Interface of Coastal Temperate Rain Forest Ecosystems: Shaping the Nature of Coastal Interactions. *The Rain Forests of Home: Profile of a North American Bioregion*. P. Schoonmaker, B. von Hagen and E. Wolf. Covelo, CA USA, Island Press: 149-187.
- Taft, O. W. and S. M. Haig. 2003. "Historical Wetlands in Oregon's Willamette Valley: Implication for Restoration of Winter Waterbird Habitat." *Wetlands* **23**(1).
- Taylor, S. B., B. E. Dutton and P. E. Poston. 2003. Field Guide to the Luckiamute River Watershed, Upper Willamette Basin: An Integrated Environmental Study for K-12 Educators. Monmouth, OR, Western Oregon University: 14.

- Teensma, P. D. A., J. T. Rienstra and M. A. Yeiter. 1991. Preliminary Reconstruction and Analysis of Change in Forest Stand Age Classes of the Oregon Coast Range from 1850 to 1940. Portland, Oregon, USDI Bureau of Land Management.
- Theurer, A. 2003. Kings Valley: The Middle Years, Benton County Historical Museum.
- Titus, J. H., J. A. Christy, D. VanderSchaaf, J. S. Kagan and E. R. Alverson. 1996. Native wetland and riparian plant communities in the Willamette Valley, Oregon. Portland, OR, Oregon Natural Heritage Program and The Nature Conservancy.
- U.S. Bureau of Land Management. 1991. Fire Occurrence (1980-1989), Oregon/Washington BLM. **2003.**
- U.S. Bureau of Land Management. 1996. Fire History 1850/1890/1920/1940, Oregon/Washington. **2003.**
- U.S. Census Bureau. 2002. U.S. Gazetteer: 2000 and 1990, U.S. Census Bureau. **2003.**
- U.S. Census Bureau. 2003. U.S. Census Bureau Quick Facts. **2003.**
- U.S. Census Bureau Population Division. 1995. Population of Counties by Decennial Census: 1900 to 1990, Richard L. Forstall. **2003.**
- University of Virginia Geospatial and Statistical Data Center. 1998. United States Historical Census Data Browser, University of Virginia. **2003.**
- USDA Natural Resources Conservation Service. 2000. SSURGO database for Benton and Polk Counties, Oregon, USDA Natural Resources Conservation Service. **2003.**
- Walstad, J. D. 1990. *Natural and Prescribed Fire in Pacific Northwest Forests*. Corvallis, Oregon, Oregon State University Press.
- Watershed Professionals Network. 1999. Oregon Watershed Assessment Manual. Salem, Oregon, Governor's Watershed Enhancement Board.
- Wemple. 1994. *This needs a citation from Garono and Brophy 2001.*
- Wevers, M. J., D. Nemeth, J. Haxton and S. Mamoyac. 1992. Coast Range Subbasin Fish Management Plan. Portland, OR, Oregon Department of Fish and Wildlife: 94.
- Whitlock, C. and M. A. Knox. 2002. Prehistoric Burning in the Pacific Northwest: Human versus Climatic Influences. *Fire, Native Peoples, and the Natural Landscape*. T. R. Vale. Washington, D.C., Island Press: 195-231.
- Whitlock, C. and M. A. Knox. 2002. Prehistoric Burning in the Pacific Northwest: Human versus Climatic Influences. *Fire, Native Peoples, and the Natural Landscape*. T. R. Vale. Washington, D.C., Island Press: 195-231.
- Whitlock, C. and M. A. Knox. in press. Prehistoric Burning in the Pacific Northwest: Human versus Climatic Influences. *Fires in Western Wilderness*. T. R. Vale. Washington, D.C., Island Press.
- Williams, G. W. 2002. References on the American Indian use of fire in ecosystems, USDA Forest Service.
- Woods, A., S. Bryce, J. Omernik and e. al. 2000. Level III and IV Ecoregion Descriptions for Oregon, US EPA/The Nature Conservancy. **2003.**